

Entropy stability and well-balancedness of space-time DG for the shallow water equations with bottom topography

A. Hiltebrand and S. Mishra

Research Report No. 2015-13
April 2015

Seminar für Angewandte Mathematik
Eidgenössische Technische Hochschule
CH-8092 Zürich
Switzerland

ENTROPY STABILITY AND WELL-BALANCEDNESS OF SPACE-TIME DG FOR THE SHALLOW WATER EQUATIONS WITH BOTTOM TOPOGRAPHY

ANDREAS HILTEBRAND AND SIDDHARTHA MISHRA

Seminar for Applied Mathematics (SAM)
Department of Mathematics
ETH Zurich
8092 Zurich, Switzerland

ABSTRACT. We describe a shock-capturing streamline diffusion space-time discontinuous Galerkin (DG) method to discretize the shallow water equations with variable bottom topography. This method, based on the entropy variables as degrees of freedom, is shown to be energy stable as well as well-balanced with respect to the lake at rest steady state. We present numerical experiments illustrating the numerical method.

1. Introduction. The shallow water equations model many phenomena of interest to meteorology and oceanography. In one space-dimension, the shallow water equations with variable bottom topography are given by,

$$\begin{aligned} h_t + (hu)_x &= 0, \\ (hu)_t + \left(\frac{1}{2}gh^2 + hu^2\right)_x &= -ghb_x. \end{aligned} \tag{1}$$

Here h is the water height, u is the depth-averaged velocity, g the gravitational constant, and b is the bottom topography. The two-dimensional version of the equations is provided in the appendix.

The shallow water equations with bottom topography are an example of nonlinear systems of balance laws of the form,

$$\mathbf{U}_t + \sum_{k=1}^d \mathbf{F}^k(\mathbf{U})_{x_k} = \mathbf{S}(\mathbf{U}) \tag{2}$$

by defining

$$\mathbf{U} = \begin{pmatrix} h \\ hu \end{pmatrix}, \quad \mathbf{F} = \begin{pmatrix} hu \\ hu^2 + \frac{1}{2}gh^2 \end{pmatrix}, \quad \mathbf{S} = \begin{pmatrix} 0 \\ -ghb_x \end{pmatrix}.$$

Here and in the following, we drop the index k in the one-dimensional case for simplicity.

It is well-known that solutions of balance laws (or conservation laws, when the source term $\mathbf{S}(\mathbf{U}) \equiv 0$) can be discontinuous due to the appearance of shock waves (hydraulic jumps for shallow water equations), even when the initial data

2010 *Mathematics Subject Classification.* Primary: 65M60, 35L60.

Key words and phrases. Well-balancedness, entropy stability, space-time DG, shallow water equations, bottom topography.

Both authors were partially supported by ERC STG NN. 306279 SPARCCLLE.

are smooth. Hence, the solutions to such equations are sought in the weak sense [3]. However, weak solutions are not unique and one needs to impose additional admissibility criteria, in the form of entropy conditions, to recover uniqueness [3]. In general, entropy conditions are of the form,

$$S(\mathbf{U})_t + \sum_{k=1}^d Q^k(\mathbf{U})_{x_k} \leq 0, \quad (3)$$

with the above *entropy inequality* being satisfied in the sense of distributions.

In the specific case of shallow water equations with variable bottom topography, it is well-known that the total energy,

$$S(\mathbf{U}) = \frac{1}{2}hu^2 + \frac{1}{2}gh^2 + ghb, \quad Q(\mathbf{U}) = \frac{1}{2}hu^3 + gh^2u + ghbu,$$

serves as the entropy with the energy flux Q .

Furthermore, the *entropy variables* $\mathbf{V} = S_{\mathbf{U}}(\mathbf{U})$, with the specific form,

$$\mathbf{V} = \begin{pmatrix} g(h+b) - \frac{1}{2}u^2 \\ u \end{pmatrix},$$

play an important role in the entropy stability analysis. In particular, we can rewrite (2) in entropy variables as

$$\mathbf{U}(\mathbf{V})_t + \sum_{k=1}^d \mathbf{F}^k(\mathbf{V})_{x_k} = \mathbf{S}(\mathbf{V}) \quad (4)$$

using the notation $\mathbf{F}^k(\mathbf{V}) = \mathbf{F}^k(\mathbf{U}(\mathbf{V}))$ and $\mathbf{S}(\mathbf{V}) = \mathbf{S}(\mathbf{U}(\mathbf{V}))$ for simplicity.

1.1. Numerical methods. The design of efficient numerical methods for approximating systems of conservation (balance) laws is fairly mature. Among the popular discretization frameworks are the finite volume (conservative finite difference) methods, based on (approximate) Riemann solvers [12]. High-order of spatial accuracy can be attained by employing suitable non-oscillatory piecewise polynomial reconstructions such as TVD, ENO and WENO. An attractive alternative is provided by the discontinuous Galerkin method.

In the specific case of shallow water equations, a further challenge is provided by the fact that most situations of interest are modeled as small perturbations of a steady state, the so-called *lake (ocean) at rest* state [13] given by,

$$h + b = \text{const}, \quad u = 0. \quad (5)$$

Therefore, one needs to design numerical methods that *preserve* this steady state. If a numerical method only approximates this steady state to truncation error, then it will not be able to resolve the very small amplitude waves (perturbations of steady state) that are of interest, except on very fine meshes. Hence, the goal has been to design *well-balanced schemes*, i.e. numerical approximations of the shallow water equations that preserve a discrete version of the lake at rest steady state [13].

A large number of well-balanced schemes have been designed for the shallow-water equations over the past decade or so. A (very incomplete) list of references is [1, 9, 10, 5, 2, 11, 14] and references therein. The basic idea behind most of these papers is to modify the numerical fluxes by a hydrostatic reconstruction and introduce a source discretization to balance the flux difference.

Unfortunately, very few rigorous stability results exist in the context of well-balanced schemes for the shallow water equations with variable bottom topography.

Given the fact that energy stability appears to be the most natural stability framework, it is natural to seek a well-balanced scheme that is also entropy stable. To the best of our knowledge, this issue is first tackled in [4], where the authors designed an energy stable scheme for the shallow water equations with bottom topography. Additionally, this scheme was also well-balanced with respect to the lake at rest as well as moving steady states. However, this scheme suffers from two limitations. Only the semi-discrete version of it was shown to be energy stable. Furthermore, the multi-dimensional version of the scheme was restricted to Cartesian grids.

The main aim of this paper is to design a scheme for the shallow-water equations with bottom topography that is

- Fully discrete.
- Able to handle unstructured grids (in several space dimensions).
- Energy stable.
- Well-balanced with respect to the lake at rest steady state.

To this end, we will design a shock-capturing space-time discontinuous Galerkin (DG) method to approximate the shallow-water equations with bottom topography. The scheme will be an extension of a space-time DG method for conservation laws, proposed in [6, 7] and based on earlier work [8] and references therein. The main novelty of the paper will be to handle the bottom topography source terms such that the resulting scheme continues to be energy stable and is in addition, well-balanced.

2. The space-time DG formulation. The aim of this section is to present the entropy stable space-time DG method for the shallow water equations with variable bottom topography. We start with the description of the mesh.

2.1. The mesh. At the n -th time level t^n , we denote the time step as $\Delta t^n = t^{n+1} - t^n$ and the update time interval as $I^n = [t^n, t^{n+1})$. For simplicity, we assume that the spatial domain $\Omega \subset \mathbb{R}^d$ is polyhedral and divide it into a triangulation \mathcal{T} , i.e. a set of open convex polyhedra $K \subset \mathbb{R}^d$ with plane faces. Furthermore, we assume mesh regularity [8] and quasiuniformity. For a generic element (cell) K , we denote

$$\Delta x_K = \text{diam}(K) \quad (\text{element width}),$$

$$\mathcal{N}(K) = \{K' \in \mathcal{T} : K' \neq K \text{ and } \text{meas}_{d-1}(\overline{K} \cap \overline{K}') > 0\} \quad (\text{neighbouring cells}).$$

The *mesh width* of the triangulation is $\Delta x(\mathcal{T}) = \max_K \Delta x_K$. A generic space-time element is the prism:

$$K \times I^n.$$

We also assume that there exists an (arbitrarily large) constant $C > 0$ such that

$$(1/C)\Delta x \leq \Delta t^n \leq C\Delta x,$$

for all time levels n .

2.2. The variational formulation. The formulation reads: find $\mathbf{V}^{\Delta x} \in \mathcal{V}_p$ such that

$$\mathcal{B}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) := \mathcal{B}_{\text{DG}}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) + \mathcal{B}_{\text{SD}}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) + \mathcal{B}_{\text{SC}}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) = 0 \quad (6)$$

for all $\mathbf{W}^{\Delta x} \in \mathcal{V}_p$, where \mathcal{V}_p denotes the space of piecewise polynomials associated with the space-time mesh with total degree at most p . The variational formulations consists of three quasilinear forms, which we will describe in the following.

2.3. The DG quasilinear form. The form \mathcal{B}_{DG} is given by

$$\begin{aligned}
\mathcal{B}_{\text{DG}}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) &= - \sum_{n,K} \int_{I^n} \int_K \left(\langle \mathbf{U}(\mathbf{V}^{\Delta x}), \mathbf{W}_t^{\Delta x} \rangle + \sum_{k=1}^d \langle \mathbf{F}^k(\mathbf{V}^{\Delta x}), \mathbf{W}_{x_k}^{\Delta x} \rangle + \langle \mathbf{S}(\mathbf{V}^{\Delta x}), \mathbf{W}^{\Delta x} \rangle \right) dx dt \\
&+ \sum_{n,K} \int_K \langle \mathbb{U}(\mathbf{V}_{n+1,-}^{\Delta x}, \mathbf{V}_{n+1,+}^{\Delta x}), \mathbf{W}_{n+1,-}^{\Delta x} \rangle dx - \sum_{n,K} \int_K \langle \mathbb{U}(\mathbf{V}_{n,-}^{\Delta x}, \mathbf{V}_{n,+}^{\Delta x}), \mathbf{W}_{n,+}^{\Delta x} \rangle dx \\
&+ \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \left\langle \mathbb{F}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}; \nu_{KK'}) + \sum_{k=1}^d \mathbf{B}^k(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) \nu_{KK'}^k, \mathbf{W}_{K,-}^{\Delta x} \right\rangle d\sigma(x) dt,
\end{aligned} \tag{7}$$

where the sums are taken over the time steps $n = 0, \dots, N-1$ and all the cells $K \in \mathcal{T}$ in the mesh. The last sum is also taken over all neighbouring cells $K' \in \mathcal{N}(K)$ of the cell K . In addition, we have employed the notation

$$\begin{aligned}
\mathbf{W}_{n,\pm}(x) &= \mathbf{W}(x, t_{\pm}^n), \\
\partial_{KK'} &= \overline{K} \cap \overline{K'}, \\
\nu_{KK'} &= \text{Unit normal for edge } KK' \text{ pointing outwards from element } K, \\
\mathbf{W}_{K,\pm}(x, t) &= \lim_{h \rightarrow 0+} \mathbf{W}(x \pm h\nu, t), \quad x \in \partial_{KK'},
\end{aligned}$$

for all $\mathbf{W} \in \mathcal{V}_p$.

The additional boundary terms for the shallow water equations with bottom topography are given by

$$\mathbf{B}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) = \begin{pmatrix} 0 \\ \frac{1}{2} g \bar{h} [\![b]\!] \end{pmatrix},$$

where for any quantity q , $\bar{q} = \frac{1}{2}(q_- + q_+)$ and $[\![q]\!] = q_+ - q_-$ denotes the mean and the jump, respectively.

Upwind fluxes are used for the temporal numerical fluxes

$$\mathbb{U}(\mathbf{V}_-, \mathbf{V}_+) = \mathbf{U}(\mathbf{V}_-).$$

This ensures causality and further allows to perform marching in time.

We use an *entropy-stable* numerical flux given by

$$\mathbb{F}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}; \nu_{KK'}) = \sum_{k=1}^d \mathbb{F}^{k,*}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) \nu_{KK'}^k - \frac{1}{2} \mathbf{D}(\mathbf{V}_{K,+}^{\Delta x} - \mathbf{V}_{K,-}^{\Delta x})$$

with $\mathbf{D} = \mathbf{D}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}; \nu_{KK'})$, which consists of an entropy-conservative flux and a diffusion operator.

Note that the diffusion is added in terms of entropy variables. This is important both for the entropy stability and the well-balancedness of the scheme.

More precisely, we will use a Rusanov type of diffusion, which is given by

$$\mathbf{D}(\mathbf{V}_-, \mathbf{V}_+; \nu) = \max\{\lambda_{\max}(\mathbf{V}_-; \nu), \lambda_{\max}(\mathbf{V}_+; \nu)\} \mathbf{U}_{\mathbf{V}} \left(\frac{1}{2}(\mathbf{V}_- + \mathbf{V}_+) \right),$$

where $\lambda_{\max}(\mathbf{U}; \nu)$ is the maximal wave speed in the direction of ν .

An entropy-conservative flux for the shallow water equations is given by (see [4])

$$\mathbb{F}^*(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) = \left(\bar{h}\bar{u}, \bar{h}\bar{u}^2 + \frac{1}{2}g\bar{h}^2 \right).$$

Inserting the numerical fluxes into the form \mathcal{B}_{DG} (7), we obtain

$$\begin{aligned} \mathcal{B}_{\text{DG}}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) &= - \sum_{n,K} \int_{I^n} \int_K \left(\langle \mathbf{U}(\mathbf{V}^{\Delta x}), \mathbf{W}_t^{\Delta x} \rangle + \sum_{k=1}^d \langle \mathbf{F}^k(\mathbf{V}^{\Delta x}), \mathbf{W}_{x_k}^{\Delta x} \rangle + \langle \mathbf{S}(\mathbf{V}^{\Delta x}), \mathbf{W}^{\Delta x} \rangle \right) dx dt \\ &\quad + \sum_{n,K} \int_K \langle \mathbf{U}(\mathbf{V}_{n+1,-}^{\Delta x}), \mathbf{W}_{n+1,-}^{\Delta x} \rangle dx - \sum_{n,K} \int_K \langle \mathbf{U}(\mathbf{V}_{n,-}^{\Delta x}), \mathbf{W}_{n,+}^{\Delta x} \rangle dx \\ &\quad + \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \left(\sum_{k=1}^d \langle \mathbb{F}^{k,*}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}), \mathbf{W}_{K,-}^{\Delta x} \rangle \nu_{KK'}^k \right) d\sigma(x) dt \\ &\quad - \frac{1}{2} \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \langle \mathbf{D}(\mathbf{V}_{K,+}^{\Delta x} - \mathbf{V}_{K,-}^{\Delta x}), \mathbf{W}_{K,-}^{\Delta x} \rangle d\sigma(x) dt \\ &\quad + \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \left(\sum_{k=1}^d \langle \mathbf{B}^k(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}), \mathbf{W}_{K,-}^{\Delta x} \rangle \nu_{KK'}^k \right) d\sigma(x) dt. \end{aligned} \quad (8)$$

2.4. Streamline diffusion operator. Following [7], we need a streamline diffusion operator to minimize oscillations that might arise in the pure space-time DG formulation. However, the streamline diffusion operator of [7] needs to be adapted to balance laws. The equation residual (or intra-element residual) is now

$$\text{Res} := \mathbf{U}(\mathbf{V}^{\Delta x})_t + \sum_{k=1}^d \mathbf{F}^k(\mathbf{V}^{\Delta x})_{x_k} - \mathbf{S}(\mathbf{V}^{\Delta x}). \quad (9)$$

The streamline diffusion operator is then given by

$$\mathcal{B}_{\text{SD}}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) = \sum_{n,K} \int_{I^n} \int_K \langle \mathcal{L}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}), \mathbf{D}^{\text{SD}} \text{Res} \rangle dx dt, \quad (10)$$

where $\mathcal{L}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x})$ denotes a linearised form of the equation in the following sense:

- $\mathcal{L}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x})$ is linear in $\mathbf{W}^{\Delta x}$
- $\mathcal{L}(\mathbf{V}^{\Delta x}, \mathbf{V}^{\Delta x}) = \text{Res}$

For conservation laws this operator can be chosen as

$$\mathcal{L}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) = \mathbf{U}_{\mathbf{V}}(\mathbf{V}^{\Delta x}) \mathbf{W}_t^{\Delta x} + \sum_{k=1}^d \mathbf{F}_{\mathbf{V}}^k(\mathbf{V}^{\Delta x}) \mathbf{W}_{x_k}^{\Delta x}.$$

The difficulty for balance laws is to include the source term in an appropriate manner without destroying the linearity. For the shallow water equations with bottom topography, it can be chosen as

$$\mathcal{L}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) = \mathbf{U}_{\mathbf{V}}(\mathbf{V}^{\Delta x}) \mathbf{W}_t^{\Delta x} + \sum_{k=1}^d \mathbf{F}_{\mathbf{V}}^k(\mathbf{V}^{\Delta x}) \mathbf{W}_{x_k}^{\Delta x} + \mathbf{G}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) \quad (11)$$

with

$$\mathbf{G}\left(\begin{pmatrix} v_1 \\ v_2 \end{pmatrix}, \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}\right) = \begin{pmatrix} -b_x w_2 \\ -b_x v_2 w_2 \end{pmatrix}.$$

This is a consequence of the following. The residual satisfies

$$\text{Res} = \mathbf{U}_{\mathbf{V}}(\mathbf{V}^{\Delta x})\mathbf{V}_t^{\Delta x} + \mathbf{F}_{\mathbf{V}}(\mathbf{V}^{\Delta x})\mathbf{V}_x^{\Delta x} + \mathbf{F}_{\mathbf{U}}(\mathbf{U}(\mathbf{V}^{\Delta x}))\mathbf{U}_x(\mathbf{V}^{\Delta x}, x) - \mathbf{S}(\mathbf{V}^{\Delta x})$$

and we have

$$\mathbf{F}_{\mathbf{U}} = \begin{pmatrix} 0 & 1 \\ gh - u^2 & 2u \end{pmatrix}, \quad \mathbf{U}_x(\mathbf{V}^{\Delta x}, x) = \begin{pmatrix} -b_x \\ -b_x u \end{pmatrix}, \quad \mathbf{S} = \begin{pmatrix} 0 \\ -ghb_x \end{pmatrix}.$$

This implies

$$\mathbf{F}_{\mathbf{U}}(\mathbf{U}(\mathbf{V}^{\Delta x}))\mathbf{U}_x(\mathbf{V}^{\Delta x}, x) - \mathbf{S}(\mathbf{V}^{\Delta x}) = \begin{pmatrix} -b_x u \\ -b_x u^2 \end{pmatrix}$$

and thus, \mathcal{L} satisfies the assumptions.

The scaling matrix in the streamline diffusion operator

$$\mathbf{D}^{\text{SD}} := C^{\text{SD}} \Delta t^n \mathbf{U}_{\mathbf{V}}^{-1}(\mathbf{V}^{\Delta x}) \quad (12)$$

remains unchanged. Here, C^{SD} is a positive constant.

2.5. Shock-capturing operator. As in [7], we further need a shock-capturing operator to stabilize any possible oscillations. The shock-capturing operator of [7] need to be modified for the shallow water equations. It is given by

$$\begin{aligned} \mathcal{B}_{\text{SC}}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) &= \sum_{n,K} \int_{I^n} \int_K D_{n,K}^{\text{SC}} \left(\langle \mathbf{W}_t^{\Delta x}, \widetilde{\mathbf{U}}_{\mathbf{V}} \mathbf{V}_t^{\Delta x} \rangle + \sum_{k=1}^d \frac{\Delta x_K^2}{(\Delta t^n)^2} \langle \mathbf{W}_{x_k}^{\Delta x}, \widetilde{\mathbf{U}}_{\mathbf{V}} \mathbf{V}_{x_k}^{\Delta x} \rangle \right) dx dt, \end{aligned} \quad (13a)$$

with $\widetilde{\mathbf{U}}_{\mathbf{V}} = \mathbf{U}_{\mathbf{V}}(\widetilde{\mathbf{V}}_{n,K})$ for brevity and

$$\widetilde{\mathbf{V}}_{n,K} = \frac{1}{\text{meas}(I^n \times K)} \int_{I^n} \int_K \mathbf{V}^{\Delta x}(x, t) dx dt.$$

being the cell average. The scaling factor is

$$D_{n,K}^{\text{SC}} = \frac{\Delta t^n C^{\text{SC}} \overline{\text{Res}}_{n,K} + (\Delta t^n)^{\frac{1}{2}} \bar{C}^{\text{SC}} \overline{\text{BRes}}_{n,K}}{\sqrt{\int_{I^n} \int_K \left(\langle \mathbf{V}_t^{\Delta x}, \widetilde{\mathbf{U}}_{\mathbf{V}} \mathbf{V}_t^{\Delta x} \rangle + \sum_{k=1}^d \frac{\Delta x_K^2}{(\Delta t^n)^2} \langle \mathbf{V}_{x_k}^{\Delta x}, \widetilde{\mathbf{U}}_{\mathbf{V}} \mathbf{V}_{x_k}^{\Delta x} \rangle \right) dx dt} + \epsilon}, \quad (13b)$$

with $\epsilon := |K|^{\frac{1}{2}} (\Delta t^n)^{\frac{-1}{2}} \left(\frac{\Delta x}{\text{diam}(\Omega)} \right)^{\theta}$, $\theta \geq 1/2$ (chosen as 1), and C^{SC} and \bar{C}^{SC} being positive constants. The scaling factor relies on the integrated intra-element residual

$$\overline{\text{Res}}_{n,K} := \sqrt{\int_{I^n} \int_K \langle \text{Res}, \mathbf{U}_{\mathbf{V}}^{-1}(\mathbf{V}^{\Delta x}) \text{Res} \rangle dx dt} \quad (13c)$$

and on the integrated boundary residual

$$\begin{aligned} \overline{\text{BRes}}_{n,K} := & \left(\int_K \|\mathbf{U}(\mathbf{V}_{n,-}^{\Delta x}) - \mathbf{U}(\mathbf{V}_{n,+}^{\Delta x})\|_{\mathbf{U}_{\mathbf{V}}^{-1}(\mathbf{V}_{n,+}^{\Delta x})}^2 dx \right. \\ & + \sum_{K'} \int_{I^n} \int_{\partial_{KK'}} \frac{\Delta t^n}{\Delta x_K} \left\| \mathbb{F}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}; \nu_{KK'}) - \sum_{k=1}^d \mathbf{F}^k(\mathbf{V}_{K,-}^{\Delta x}) \nu_{KK'}^k \right. \\ & \left. \left. + \sum_{k=1}^d \mathbf{B}^k(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) \nu_{KK'}^k \right\|_{\mathbf{U}_{\mathbf{V}}^{-1}(\mathbf{V}_{K,-}^{\Delta x})}^2 d\sigma dt \right)^{\frac{1}{2}}, \end{aligned} \quad (13d)$$

which needed to be adapted at the spatial boundaries.

3. Entropy stability. We will show that the scheme is entropy-stable even in the presence of source terms due to the bottom topography.

Theorem 3.1. *Consider the space-time DG scheme (6) for the shallow water equations with non-constant bottom topography (1). For simplicity, assume that the exact and approximate solutions have compact support inside the spatial domain Ω . Then, the resulting scheme is entropy-stable, i.e. the approximate solutions $\mathbf{U}^{\Delta x} = \mathbf{U}(\mathbf{V}^{\Delta x})$ satisfy*

$$\int_{\Omega} S(\mathbf{U}^{\Delta x}(x, t_-^N)) dx \leq \int_{\Omega} S(\mathbf{U}^{\Delta x}(x, t_-^0)) dx. \quad (14)$$

Proof. Let us use $\tilde{\cdot}$ to denote the quantities used without bottom topography. In this case the entropy is given by

$$\tilde{S}(\mathbf{U}) = \frac{1}{2} h u^2 + \frac{1}{2} g h^2,$$

with the entropy variables being

$$\tilde{\mathbf{V}} = \begin{pmatrix} gh - \frac{1}{2} u^2 \\ u \end{pmatrix}.$$

From the entropy stability proof we know (compare [7, 6])

$$\begin{aligned} & \tilde{\mathcal{B}}_{\text{DG}}(\mathbf{V}^{\Delta x}, \tilde{\mathbf{V}}^{\Delta x}) + \frac{1}{2} \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \left\langle \tilde{\mathbf{V}}_{K,-}^{\Delta x}, \mathbf{D}(\mathbf{V}_{K,+}^{\Delta x} - \mathbf{V}_{K,-}^{\Delta x}) \right\rangle d\sigma dt \\ & \geq \int_{\Omega} \tilde{S}(\mathbf{U}^{\Delta x}(x, t_-^N)) dx - \int_{\Omega} \tilde{S}(\mathbf{U}^{\Delta x}(x, t_-^0)) dx. \end{aligned}$$

We will show

$$\begin{aligned} & \mathcal{B}_{\text{DG}}(\mathbf{V}^{\Delta x}, \mathbf{V}^{\Delta x}) - \tilde{\mathcal{B}}_{\text{DG}}(\mathbf{V}^{\Delta x}, \tilde{\mathbf{V}}^{\Delta x}) - \frac{1}{2} \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \left\langle \tilde{\mathbf{V}}_{K,-}^{\Delta x}, \mathbf{D}(\mathbf{V}_{K,+}^{\Delta x} - \mathbf{V}_{K,-}^{\Delta x}) \right\rangle d\sigma dt \\ & \geq \int_{\Omega} S(\mathbf{U}^{\Delta x}(x, t_-^N)) dx - \int_{\Omega} S(\mathbf{U}^{\Delta x}(x, t_-^0)) dx - \int_{\Omega} \tilde{S}(\mathbf{U}^{\Delta x}(x, t_-^N)) dx + \int_{\Omega} \tilde{S}(\mathbf{U}^{\Delta x}(x, t_-^0)) dx. \end{aligned} \quad (15)$$

This implies

$$\mathcal{B}_{\text{DG}}(\mathbf{V}^{\Delta x}, \mathbf{V}^{\Delta x}) \geq \int_{\Omega} S(\mathbf{U}^{\Delta x}(x, t_-^N)) dx - \int_{\Omega} S(\mathbf{U}^{\Delta x}(x, t_-^0)) dx \quad (16)$$

The difference in the spatial interior part is given by

$$\begin{aligned}
& \sum_{n,K} \int_{I^n} \int_K -\langle \mathbf{S}(\mathbf{V}^{\Delta x}), \mathbf{V}^{\Delta x} \rangle dx dt - \sum_{n,K} \int_{I^n} \int_K \langle \mathbf{F}(\mathbf{V}^{\Delta x}), (\mathbf{V}^{\Delta x} - \tilde{\mathbf{V}}^{\Delta x})_x \rangle dx dt \\
&= \sum_{n,K} \int_{I^n} \int_K \left(\left\langle \begin{pmatrix} 0 \\ -ghb_x \end{pmatrix}, \begin{pmatrix} g(h+b) - \frac{1}{2}u^2 \\ u \end{pmatrix} \right\rangle - \left\langle \begin{pmatrix} hu \\ hu^2 + \frac{1}{2}gh^2 \end{pmatrix}, \begin{pmatrix} gb \\ 0 \end{pmatrix}_x \right\rangle \right) dx dt \\
&= \sum_{n,K} \int_{I^n} \int_K (ghb_x u - hugb_x) dx dt = 0
\end{aligned} \tag{17}$$

The difference in the spatial boundary part without diffusion is given by

$$\begin{aligned}
& \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \langle B(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}), \mathbf{V}_{K,-}^{\Delta x} \rangle \nu_{KK'} d\sigma dt \\
&+ \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \left\langle \mathbb{F}^*(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}; \nu_{KK'}), (\mathbf{V}_{K,-}^{\Delta x} - \tilde{\mathbf{V}}_{K,-}^{\Delta x}) \right\rangle d\sigma dt \\
&= \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \left(\left\langle \begin{pmatrix} 0 \\ \frac{1}{2}g\bar{h}[b] \end{pmatrix}, \begin{pmatrix} g(h_- + b_-) - \frac{1}{2}u_-^2 \\ u_- \end{pmatrix} \right\rangle \right. \\
&\quad \left. + \left\langle \begin{pmatrix} \bar{h}\bar{u} \\ \bar{h}\bar{u}^2 + \frac{1}{2}g\bar{h}^2 \end{pmatrix}, \begin{pmatrix} gb_- \\ 0 \end{pmatrix} \right\rangle \right) \nu_{KK'} d\sigma dt \\
&= \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \left(\frac{1}{2}g\bar{h}[b]u_- + \bar{h}\bar{u}gb_- \right) \nu_{KK'} d\sigma dt \\
&= \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \left(\frac{1}{2}g\bar{h}[b]\bar{u} - \frac{1}{2}\bar{h}\bar{u}g[b] \right) \nu_{KK'} d\sigma dt = 0
\end{aligned} \tag{18}$$

For the diffusion part we can proceed as in the case of the conservation law to obtain

$$\begin{aligned}
& -\frac{1}{2} \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \langle \mathbf{V}_{K,-}^{\Delta x}, \mathbf{D}(\mathbf{V}_{K,+}^{\Delta x} - \mathbf{V}_{K,-}^{\Delta x}) \rangle d\sigma dt \\
& \geq \frac{1}{4} \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \langle \mathbf{V}_{K,+}^{\Delta x} - \mathbf{V}_{K,-}^{\Delta x}, \mathbf{D}(\mathbf{V}_{K,+}^{\Delta x} - \mathbf{V}_{K,-}^{\Delta x}) \rangle d\sigma dt \geq 0
\end{aligned} \tag{19}$$

The difference in the temporal interior part is given by

$$\begin{aligned}
& -\sum_{n,K} \int_{I^n} \int_K \left\langle \mathbf{U}^{\Delta x}, (\mathbf{V}^{\Delta x} - \tilde{\mathbf{V}}^{\Delta x})_t \right\rangle dx dt \\
&= -\sum_{n,K} \int_{I^n} \int_K \left\langle \mathbf{U}^{\Delta x}, \underbrace{\begin{pmatrix} gb \\ 0 \end{pmatrix}_t}_{=0} \right\rangle dx dt = 0
\end{aligned} \tag{20}$$

The difference in the temporal boundary part is given by

$$\begin{aligned}
& \sum_{n,K} \int_K \left\langle \mathbf{U}_{n+1,-}^{\Delta x}, (\mathbf{V}_{n+1,-}^{\Delta x} - \tilde{\mathbf{V}}_{n+1,-}^{\Delta x}) \right\rangle dx - \sum_{n,K} \int_K \left\langle \mathbf{U}_{n,-}^{\Delta x}, (\mathbf{V}_{n,+}^{\Delta x} - \tilde{\mathbf{V}}_{n,+}^{\Delta x}) \right\rangle dx \\
&= \sum_{n,K} \int_K \left\langle \begin{pmatrix} h \\ hu \end{pmatrix}_{n+1,-}, \begin{pmatrix} gb \\ 0 \end{pmatrix}_{n+1,-} \right\rangle dx - \sum_{n,K} \int_K \left\langle \begin{pmatrix} h \\ hu \end{pmatrix}_{n,-}, \begin{pmatrix} gb \\ 0 \end{pmatrix}_{n,+} \right\rangle dx \\
&= \sum_{n,K} \int_K h_{n+1,-} gb_{n+1} dx - \sum_{n,K} \int_K h_{n,-} gb_n dx \\
&= \sum_K \int_K h_{N,-} gb_N dx - \sum_K \int_K h_{0,-} gb_0 dx \\
&= \int_{\Omega} S(\mathbf{U}^{\Delta x}(x, t_-^N)) dx - \int_{\Omega} S(\mathbf{U}^{\Delta x}(x, t_-^0)) dx \\
&\quad - \int_{\Omega} \tilde{S}(\mathbf{U}^{\Delta x}(x, t_-^N)) dx + \int_{\Omega} \tilde{S}(\mathbf{U}^{\Delta x}(x, t_-^0)) dx.
\end{aligned} \tag{21}$$

Combining the equations (17), (18), (19), (20), and (21), we obtain (15). For the streamline diffusion and the shock-capturing term, we can proceed analogously as for conservation laws to obtain

$$\mathcal{B}_{\text{SD}}(\mathbf{V}^{\Delta x}, \mathbf{V}^{\Delta x}) \geq 0 \tag{22}$$

and

$$\mathcal{B}_{\text{SC}}(\mathbf{V}^{\Delta x}, \mathbf{V}^{\Delta x}) \geq 0. \tag{23}$$

Combining (6), (16), (22), and (23), we obtain the stability estimate (14). \square

4. Well-balancedness. We will show that the space-time DG scheme is well-balanced.

Theorem 4.1. *Consider the space-time DG scheme (6) for the shallow water equations (1). The resulting scheme is well-balanced, i.e. it can preserve the “lake at rest” state exactly.*

Proof. We will concentrate on the one-dimensional case, as the proof can be extended to the two-dimensional case in a straightforward manner. Let us consider initial conditions, such that the “lake at rest” condition is satisfied:

$$u_0 = 0, \quad h_0 + b = \text{const.}$$

We consider

$$\mathbf{V}^{\Delta x} = \begin{pmatrix} g(h_0 + b) \\ 0 \end{pmatrix} \quad \forall t \tag{24}$$

and we will show that $\mathcal{B}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) = 0$ is satisfied for all $\mathbf{W}^{\Delta x} \in \mathcal{V}_p$.

We will use the notation $\mathbf{U}^{\Delta x} = \mathbf{U}(\mathbf{V}^{\Delta x}) = (h, hu)^T$ for simplicity. Let us start with the temporal part of the DG operator

$$\begin{aligned}
& - \sum_{n,K} \int_{I^n} \int_K \langle \mathbf{U}^{\Delta x}, \mathbf{W}_t^{\Delta x} \rangle dx dt \\
& + \sum_{n,K} \int_K \langle \mathbf{U}_{n+1,-}^{\Delta x}, \mathbf{W}_{n+1,-}^{\Delta x} \rangle dx - \sum_{n,K} \int_K \langle \mathbf{U}_{n,-}^{\Delta x}, \mathbf{W}_{n,+}^{\Delta x} \rangle dx \\
& = \sum_{n,K} \int_{I^n} \int_K \langle \mathbf{U}_t^{\Delta x}, \mathbf{W}^{\Delta x} \rangle dx dt \\
& \quad - \sum_{n,K} \int_K \langle \mathbf{U}_{n+1,-}^{\Delta x}, \mathbf{W}_{n+1,-}^{\Delta x} \rangle dx + \sum_{n,K} \int_K \langle \mathbf{U}_{n,+}^{\Delta x}, \mathbf{W}_{n,+}^{\Delta x} \rangle dx \\
& \quad + \sum_{n,K} \int_K \langle \mathbf{U}_{n+1,-}^{\Delta x}, \mathbf{W}_{n+1,-}^{\Delta x} \rangle dx - \sum_{n,K} \int_K \langle \mathbf{U}_{n,-}^{\Delta x}, \mathbf{W}_{n,+}^{\Delta x} \rangle dx \quad (\text{int. by parts}) \\
& = \sum_{n,K} \int_{I^n} \int_K \underbrace{\langle \mathbf{U}_t^{\Delta x}, \mathbf{W}^{\Delta x} \rangle}_{=0} dx dt + \sum_{n,K} \int_K \underbrace{\langle \mathbf{U}_{n,+}^{\Delta x} - \mathbf{U}_{n,-}^{\Delta x}, \mathbf{W}_{n,+}^{\Delta x} \rangle}_{=0} dx \\
& = 0 \quad (\text{by (24)}). \tag{25}
\end{aligned}$$

Next, we consider the spatial parts of the DG operator. We need to integrate by parts

$$\begin{aligned}
& - \sum_{n,K} \int_{I^n} \int_K \sum_{k=1}^d \langle \mathbf{F}^k(\mathbf{V}^{\Delta x}), \mathbf{W}_{x_k}^{\Delta x} \rangle dx dt \\
& = \sum_{n,K} \int_{I^n} \int_K \sum_{k=1}^d \langle \mathbf{F}^k(\mathbf{V}^{\Delta x})_{x_k}, \mathbf{W}^{\Delta x} \rangle dx dt \\
& \quad - \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \left\langle \sum_{k=1}^d \mathbf{F}^k(\mathbf{V}_{K,-}^{\Delta x}) \nu_{KK'}^k, \mathbf{W}_{K,-}^{\Delta x} \right\rangle d\sigma dt. \tag{26}
\end{aligned}$$

We start with the resulting interior part and observe that

$$\begin{aligned}
& \mathbf{F}(\mathbf{V}^{\Delta x})_x - \mathbf{S}(\mathbf{V}^{\Delta x}) \\
& = \mathbf{F}_{\mathbf{U}}(\mathbf{U}^{\Delta x}) \mathbf{U}_x^{\Delta x} - \mathbf{S}(\mathbf{V}^{\Delta x}) = \begin{pmatrix} 0 & 1 \\ gh - u^2 & 2u \end{pmatrix} \begin{pmatrix} h_x \\ (hu)_x \end{pmatrix} + \begin{pmatrix} 0 \\ ghb_x \end{pmatrix} \\
& = \begin{pmatrix} 0 & 1 \\ gh & 0 \end{pmatrix} \begin{pmatrix} h_x \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ ghb_x \end{pmatrix} = \begin{pmatrix} 0 \\ gh(h+b)_x \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \tag{27}
\end{aligned}$$

by (24). This trivially implies

$$\sum_{n,K} \int_{I^n} \int_K \left(\sum_{k=1}^d \langle \mathbf{F}^k(\mathbf{V}^{\Delta x})_{x_k}, \mathbf{W}^{\Delta x} \rangle - \langle \mathbf{S}(\mathbf{V}(\mathbf{V}^{\Delta x})), \mathbf{W}^{\Delta x} \rangle \right) dx dt = 0. \tag{28}$$

Now we consider the boundary fluxes without the diffusion part and obtain

$$\begin{aligned}
& (\mathbb{F}^*(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) - \mathbf{F}(\mathbf{V}_{K,-}^{\Delta x}) + \mathbf{B}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}))\nu_{KK'} \\
&= \left(\left(\frac{\bar{h}\bar{u}}{\bar{h}\bar{u}^2 + \frac{1}{2}g\bar{h}^2} \right) - \left(\frac{h_-u_-}{h_-u_-^2 + \frac{1}{2}gh_-^2} \right) + \left(\frac{0}{\frac{1}{2}g\bar{h}[[b]]} \right) \right) \nu_{KK'} \\
&= \left(\left(\frac{0}{\frac{1}{2}g\bar{h}^2} \right) - \left(\frac{0}{\frac{1}{2}gh_-^2} \right) + \left(\frac{0}{\frac{1}{2}g\bar{h}[[b]]} \right) \right) \nu_{KK'} \quad (\text{by (24)}) \\
&= \left(\frac{0}{\frac{1}{2}g(\bar{h}^2 - h_-^2 + \bar{h}[[b]])} \right) \nu_{KK'}
\end{aligned}$$

Let us add and subtract $\bar{h}[[h]]$ to obtain

$$\bar{h}^2 - h_-^2 + \bar{h}[[b]] = \bar{h}^2 - h_-^2 - \underbrace{\bar{h}[[h]]}_{=0} + \bar{h}[[h+b]] = \bar{h}^2 - h_-^2 - \bar{h}[[h]].$$

A short calculation confirms that $\bar{h}^2 - h_-^2 - \bar{h}[[h]]$ actually vanishes:

$$\begin{aligned}
\bar{h}^2 - h_-^2 - \bar{h}[[h]] &= \frac{1}{2}h_+^2 + \frac{1}{2}h_-^2 - h_-^2 - \frac{1}{2}(h_- + h_+)(h_+ - h_-) \\
&= \frac{1}{2}h_+^2 + \frac{1}{2}h_-^2 - h_-^2 - \frac{1}{2}h_+^2 + \frac{1}{2}h_-^2 = 0.
\end{aligned}$$

This results in

$$(\mathbb{F}^*(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) - \mathbf{F}(\mathbf{V}_{K,-}^{\Delta x}) + \mathbf{B}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}))\nu_{KK'} = 0 \quad (29)$$

and

$$\begin{aligned}
& \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \left(\left\langle \mathbb{F}^*(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) - \mathbf{F}(\mathbf{V}_{K,-}^{\Delta x}) \right. \right. \\
& \quad \left. \left. + \mathbf{B}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x})\nu_{KK'}, \mathbf{W}_{K,-}^{\Delta x} \right\rangle \right) d\sigma dt = 0.
\end{aligned} \quad (30)$$

Now we consider the diffusion part of the flux:

$$\mathbf{D}(\mathbf{V}_{K,+}^{\Delta x} - \mathbf{V}_{K,-}^{\Delta x})$$

Note that

$$\mathbf{V}_{K,+}^{\Delta x} - \mathbf{V}_{K,-}^{\Delta x} = \begin{pmatrix} g[[h+b]] - \frac{1}{2}[[u^2]] \\ [[u]] \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (\text{by (24)}).$$

This yields

$$\mathbf{D}(\mathbf{V}_{K,+}^{\Delta x} - \mathbf{V}_{K,-}^{\Delta x}) = 0 \quad (31)$$

and

$$-\frac{1}{2} \sum_{n,K,K'} \int_{I^n} \int_{\partial_{KK'}} \langle \mathbf{D}(\mathbf{V}_{K,+}^{\Delta x} - \mathbf{V}_{K,-}^{\Delta x}), \mathbf{W}_{K,-}^{\Delta x} \rangle d\sigma dt = 0. \quad (32)$$

Combining (25), (28), (30), and (32) we obtain that the DG form is satisfied, i.e.

$$\mathcal{B}_{\text{DG}}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) = 0. \quad (33)$$

For the streamline and shock-capturing terms, first note that the interior residual vanishes by (27):

$$\text{Res} = \mathbf{U}_t^{\Delta x} + \sum_{k=1}^d \mathbf{F}^k(\mathbf{V}^{\Delta x})_{x_k} - \mathbf{S}(\mathbf{V}^{\Delta x}) = 0.$$

Similarly, also the boundary residual vanishes by (29) and (31):

$$\begin{aligned} \overline{\text{BRes}}_{n,K} = & \left(\int_K \|\mathbf{U}_{n,-}^{\Delta x} - \mathbf{U}_{n,+}^{\Delta x}\|_{\mathbf{U}_V^{-1}(\mathbf{V}_{n,+}^{\Delta x})}^2 dx \right. \\ & + \sum_{K'} \int_{I^n} \int_{\partial_{KK'}} \frac{\Delta t^n}{\Delta x_K} \|(\mathbb{F}^*(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) - \mathbf{F}(\mathbf{V}_{K,-}^{\Delta x}) \\ & \quad \left. + \mathbf{B}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}))\nu_{KK'}\|_{\mathbf{U}_V^{-1}(\mathbf{V}_{K,-}^{\Delta x})}^2 d\sigma dt \right)^{\frac{1}{2}} = 0. \end{aligned}$$

This trivially implies

$$\mathcal{B}_{\text{SD}}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) = 0$$

and

$$\mathcal{B}_{\text{SC}}(\mathbf{V}^{\Delta x}, \mathbf{W}^{\Delta x}) = 0.$$

We conclude that the scheme is indeed well-balanced. \square

Remark 1. The proof relies on an integration by parts in the spatial terms. If numerical quadrature is used, this introduces an additional error and the resulting scheme will in general not be well-balanced. Therefore, rather than implementing (8) directly, one should implement it after an integration by parts is performed as in (26). This will guarantee that the scheme is well-balanced even if numerical quadrature is used.

5. Numerical Experiments. We present several numerical experiments to demonstrate the shock-capturing streamline diffusion space-time DG scheme. In all experiments, we have chosen to represent the bottom topography exactly. The constant in the scheme are set to $C^{\text{SD}} = 10$, $C^{\text{SC}} = 1$ and $\bar{C}^{\text{SC}} = 0$. A value of $g = 9.812$ is used for the gravitational constant in all the experiments.

5.1. Lake at rest. Let us consider the following problem taken from [4]. The bottom topography is given

$$b(x) = \begin{cases} \frac{4-(x-10)^2}{20} & \text{if } |x-10| < 2 \\ 0 & \text{otherwise} \end{cases} \quad (34)$$

The initial condition are “lake at rest”, i.e. $u = 0$ and $h + b = 1$. We compute on the domain $[0, 20]$ and calculate up to the final time $t = 10$.

The bottom topography and the computed surface height of one example run are shown in Figure 1.

The well-balanced property of the scheme is best demonstrated by tabulating relative L^1 errors of the total height $h + b$ (Table 1) with respect to the resolution as well as the polynomial order.

The results show that scheme preserves the exact steady state up to machine precision. The error is slightly bigger for higher polynomial degrees. This is to be expected as the conditioning becomes slightly worse with a higher polynomial degree (or increased resolution).




FIGURE 1. Water level $h + b$ and bottom topography b for the lake at rest for $p = 2$ and $N_c = 80$.

N_c	$p = 0$	$p = 1$	$p = 2$
20	1.1e-17	3.3e-16	3.3e-15
40	5.6e-18	1.9e-15	1.6e-15
80	2.8e-18	5.0e-15	5.5e-15
160	4.2e-18	1.2e-14	1.3e-14
320	6.9e-18	2.5e-14	1.6e-14

TABLE 1. Relative L^1 errors of the water surface $h + b$ for the lake at rest case.

5.2. Perturbed lake at rest. The main interest is not in steady state itself (as it is known in advance), but rather waves, representing small perturbations of the lake at rest. Therefore, we perturb the lake of rest initial condition of the last experiment by choosing (as in [4])

$$h(x) = \begin{cases} 1.01 - b(x) & \text{if } |x - 6| < 1/4 \\ 1 - b(x) & \text{otherwise} \end{cases} \quad (35)$$


The bottom topography and the initial velocity are unchanged, i.e. $u = 0$ and b is given by (34). The exact solution consists of a left-going and a right-going wave. We evolve the flow only up to $t = 1.5$ such that the waves have not yet left the domain. Figure 2 shows the water surface height $h + b$ for piecewise constant, linear, and quadratic functions. For all polynomial degrees, the waves can be clearly identified without any spurious numerical artifacts or additional waves. The waves are quite diffused in the case of $p = 0$. However, the accuracy improves to a great extent by considering higher polynomial degrees.

5.3. Two-dimensional lake at rest. We consider the following bottom topography


$$b(x) = 0.8 \exp(-5(x - 0.9)^2 - 50(y - 0.5)^2) \quad (36)$$

in the domain $[0, 2] \times [0, 1]$. The bottom topography is illustrated in Figure 3. The initial condition is given by the lake at rest state $u = 0$ and $h + b = 1$.

We compute up to the final time $t = 1$. We start on a triangular mesh with 178 cells and perform uniform refinements. Table 2 shows the resulting relative L^1

FIGURE 2. $h + b$ for the perturbed lake at rest.

errors of $h + b$ for different polynomial degrees. The error is again on the order of machine precision, which demonstrates the well-balancedness of the scheme also in the two-dimensional case.

FIGURE 3. Bottom topography b for the lake at rest case.

5.4. Perturbed two-dimensional lake at rest. The experiment in the last subsection is perturbed by adding 0.01 to h in the region $0.1 \leq x \leq 0.2$. We use reflecting wall boundary conditions at the top and the bottom boundary. The left and right boundaries are specified as non-reflecting boundaries. Thus, the initial bump separates into two waves, a left-going and a right-going one. The leftgoing wave hits the left boundary at about $t = 0.03$ and leaves the domain without any

N_c	$p = 0$	$p = 1$	$p = 2$
178	3.3e-17	2.3e-15	1.6e-15
712	3.6e-17	4.0e-15	4.0e-15
2848	3.7e-17	1.8e-14	1.4e-14

TABLE 2. Relative L^1 errors of the water surface $h + b$ for the two-dimensional lake at rest case.

reflections. The rightgoing wave is slowed down in the middle by the bottom topography and this then creates a complicated pattern. Figure 4 shows the flow at various times for piecewise constant and piecewise linear functions. The results for piecewise quadratic functions can be found in Figure 5. The results for $p = 0$ are very smeared, but still qualitatively the flow is captured. The results for higher degrees are much more accurate, even on a coarse mesh. This experiment clearly demonstrates the ability of the proposed space-time DG method to compute small perturbations of the lake at rest steady state, in a stable and accurate manner.




FIGURE 4. $h + b$ for the perturbed lake at rest. The number of cells is 182272 and 45568 for $p = 0$ and $p = 1$, respectively.




FIGURE 5. $h+b$ for the perturbed lake at rest with 11392 cells and $p = 2$.

6. Conclusion. We extend the shock-capturing streamline diffusion space-time DG method of [6, 7] to approximate the shallow water equations with variable bottom topography. The proposed schemes are energy stable, even for the fully discrete case. Furthermore, we show that the proposed schemes are *well-balanced* i.e. they preserve the lake at rest steady state. The key design elements of the scheme are the use of entropy (energy) variables as degrees of freedom, energy conservative numerical fluxes and discretizations of the bottom topography and the design of suitable streamline diffusion and shock-capturing operators.

Numerical experiments that demonstrate the well-balancing of the schemes and the stability and accuracy in resolving small perturbations of the lake at rest steady state are presented. They illustrate that the schemes, particularly with piecewise linear and piecewise quadratic basis functions, perform very well.

The proposed schemes are implicit in time and involve the solution of non-linear algebraic equations in every time step. However, the schemes are unconditionally energy stable. Hence, very large time steps can be chosen. This flexibility is particularly advantageous while dealing with problems in oceanography, where gravity waves can lead to excessively small time steps for explicit methods. The schemes uses unstructured grids in two space dimensions. This is a marked advantage, when dealing with flows in domains with complicated boundaries such as shorelines. Furthermore, space-time DG schemes lend themselves to adaptive algorithms (particularly goal oriented adaptive algorithms) in a natural manner, see [6]. Hence, these schemes could form an attractive alternative to standard well-balanced schemes for shallow water flows in complex domains that involve multiple time scales. Such extensions will be considered in the future.

7. Appendix - two-dimensional formulation. The shallow water equations in two spatial dimensions are given by

$$\begin{aligned} h_t + (hu)_x &= 0, \\ (hu)_t + \left(\frac{1}{2}gh^2 + hu^2\right)_x + (huv)_y &= -ghb_x, \\ (hv)_t + (huv)_x + \left(\frac{1}{2}gh^2 + hv^2\right)_y &= -ghb_y. \end{aligned} \quad (37)$$

This can be written in the form (2) by defining

$$\mathbf{U} = \begin{pmatrix} h \\ hu \\ hv \end{pmatrix}, \quad \mathbf{F}^1 = \begin{pmatrix} hu \\ hu^2 + \frac{1}{2}gh^2 \\ huv \end{pmatrix}, \quad \mathbf{F}^2 = \begin{pmatrix} hu \\ huv \\ hv^2 + \frac{1}{2}gh^2 \end{pmatrix}, \quad \mathbf{S} = \begin{pmatrix} 0 \\ -ghb_x \\ -ghb_y \end{pmatrix}.$$

An entropy pair is given by

$$\begin{aligned} S(\mathbf{U}) &= \frac{1}{2}hu^2 + \frac{1}{2}hv^2 + \frac{1}{2}gh^2 + ghb, \\ Q^1(\mathbf{U}) &= \frac{1}{2}hu^3 + \frac{1}{2}huv^2 + gh^2u + ghbu, \\ Q^2(\mathbf{U}) &= \frac{1}{2}hu^2v + \frac{1}{2}hv^3 + gh^2v + ghbv, \end{aligned}$$

which leads to the entropy variables

$$\mathbf{V} = \begin{pmatrix} g(h+b) - \frac{1}{2}u^2 - \frac{1}{2}v^2 \\ u \\ v \end{pmatrix}.$$

Entropy-conservative fluxes are given by (see [4])

$$\mathbb{F}^{1,*}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) = \begin{pmatrix} \bar{h}\bar{u} \\ \bar{h}\bar{u}^2 + \frac{1}{2}g\bar{h}^2 \\ \bar{h}\bar{u}\bar{v} \end{pmatrix}, \quad \mathbb{F}^{2,*}(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) = \begin{pmatrix} \bar{h}\bar{u} \\ \bar{h}\bar{u}\bar{v} \\ \bar{h}\bar{u}^2 + \frac{1}{2}g\bar{h}^2 \end{pmatrix}.$$

The additional boundary terms are given by

$$\mathbf{B}^1(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) = \begin{pmatrix} 0 \\ \frac{1}{2}g\bar{h}\llbracket b \rrbracket \\ 0 \end{pmatrix}, \quad \mathbf{B}^2(\mathbf{V}_{K,-}^{\Delta x}, \mathbf{V}_{K,+}^{\Delta x}) = \begin{pmatrix} 0 \\ 0 \\ \frac{1}{2}g\bar{h}\llbracket b \rrbracket \end{pmatrix}.$$

The function \mathbf{G} in the linearised operator \mathcal{L} in the streamline diffusion operator (11) can be chosen as

$$\mathbf{G} \left(\begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}, \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} \right) = (-b_x w_2 - b_y w_3) \begin{pmatrix} 1 \\ v_2 \\ v_3 \end{pmatrix}.$$

This can be seen as follows. The residual satisfies

$$\begin{aligned} \text{Res} &= \mathbf{U}_{\mathbf{V}}(\mathbf{V}^{\Delta x}) \mathbf{V}_t^{\Delta x} \\ &+ \sum_{k=1}^2 (\mathbf{F}_{\mathbf{V}}^k(\mathbf{V}^{\Delta x}) \mathbf{V}_{x_k}^{\Delta x} + \mathbf{F}_{\mathbf{U}}^k(\mathbf{U}(\mathbf{V}^{\Delta x})) \mathbf{U}_{x_k}(\mathbf{V}^{\Delta x}, x)) - \mathbf{S}(\mathbf{V}^{\Delta x}) \end{aligned}$$

and

$$\mathbf{F}_{\mathbf{U}}^1 = \begin{pmatrix} 0 & 1 & 0 \\ gh - u^2 & 2u & 0 \\ -uv & v & u \end{pmatrix}, \quad \mathbf{U}_x(\mathbf{V}^{\Delta x}, x) = \begin{pmatrix} -b_x \\ -b_x u \\ -b_x v \end{pmatrix},$$

$$\mathbf{F}_{\mathbf{U}}^2 = \begin{pmatrix} 0 & 0 & 1 \\ -uv & v & u \\ gh - v^2 & 0 & 2v \end{pmatrix}, \quad \mathbf{U}_y(\mathbf{V}^{\Delta x}, x) = \begin{pmatrix} -b_y \\ -b_y u \\ -b_y v \end{pmatrix}, \quad \mathbf{S} = \begin{pmatrix} 0 \\ -ghb_x \\ -ghb_y \end{pmatrix}.$$

Therefore,

$$\sum_{k=1}^2 \mathbf{F}_{\mathbf{U}}^k(\mathbf{U}(\mathbf{V}^{\Delta x})) \mathbf{U}_{x_k}(\mathbf{V}^{\Delta x}, x) - \mathbf{S}(\mathbf{V}^{\Delta x}) = (-b_x u - b_y v) \begin{pmatrix} 1 \\ u \\ v \end{pmatrix}$$

and then \mathcal{L} satisfies the assumptions.

References

- [1] E. Audusse, F. Bouchut, M. O. Bristeau, R. Klien and B. Perthame. A fast and stable well-balanced scheme with hydrostatic reconstruction for shallow water flows. *SIAM. Jl. Sci. Comp.*, 25 (6), 2004, 2050 - 2065.
- [2] M. Castro, J. M. Gallardo, C. Parés. High order finite volume schemes based on reconstruction of states for solving hyperbolic systems with non-conservative products. *Math. Comp.*, 75, 2006, 1103-1134.
- [3] C. Dafermos. Hyperbolic conservation laws in continuum physics. Springer, Berlin, 2000.
- [4] U. S. Fjordholm, S. Mishra and E. Tadmor, Well-balanced and energy stable schemes for the shallow water equations with discontinuous topography, *Journal of Computational Physics*, **230** (2011), 5587 – 5609.
- [5] J. M. Greenberg and A. Y. LeRoux. A well-balanced scheme for numerical processing of source terms in hyperbolic equations. *SIAM J. Numer. Anal.*, 33, 1996, 1-16.
- [6] A. Hildebrand, *Entropy-stable discontinuous Galerkin finite element methods with streamline diffusion and shock-capturing for hyperbolic systems of conservation laws.*, PhD thesis, ETH Zurich, 2014, No. 22279.
- [7] A. Hildebrand and S. Mishra, Entropy stable shock capturing space-time discontinuous Galerkin schemes for systems of conservation laws, *Numerische Mathematik*, **126** (2014), 103–151.
- [8] J. Jaffre, C. Johnson and A. Szepessy, Convergence of the discontinuous Galerkin finite element method for hyperbolic conservation laws, *Math. Model. Meth. Appl. Sci.*, **5** (1995), 367–386.
- [9] S. Jin. A steady state capturing method for hyperbolic systems with geometrical source terms. *Math. Model. Numer. Anal.*, 35, 2001, 631-646.
- [10] S. Jin and X. Wen. An efficient method for computing hyperbolic systems with geometrical source terms having concentrations. *J. Comput. Math.*, 22, 2004, 230-249.
- [11] A. Kurganov and D. Levy. Central-upwind schemes for the St. Vernant system. *Math. Model. Num. Anal.*, 36, 2002, 397-425.
- [12] R. J. LeVeque. Finite volume methods for hyperbolic problems. *Cambridge university press*, Cambridge, 2002.
- [13] R. J. LeVeque. Balancing source terms and flux gradients in high-resolution Godunov methods: The quasi-steady wave-propagation algorithm. *J. Comput. Phys.*, 146, 346-365, 1998.
- [14] S. Noelle, N. Pankratz, G. Puppo and J. Natvig. Well-balanced finite volume schemes of arbitrary order of accuracy for shallow water flows. *J. Comput. Phys.*, 213, 474-499, 2006.

E-mail address: andreas.hildebrand@sam.math.ethz.ch

E-mail address: smishra@sam.math.ethz.ch

Recent Research Reports

Nr.	Authors/Title
2015-03	P. Chen and A. Quarteroni and G. Rozza Reduced order methods for uncertainty quantification problems
2015-04	S. Larsson and Ch. Schwab Compressive Space-Time Galerkin Discretizations of Parabolic Partial Differential Equations
2015-05	S. May New spacetime discontinuous Galerkin methods for solving convection-diffusion systems
2015-06	H. Heumann and R. Hiptmair and C. Pagliantini Stabilized Galerkin for Transient Advection of Differential Forms
2015-07	J. Dick and F.Y. Kuo and Q.T. Le Gia and Ch. Schwab Fast QMC matrix-vector multiplication
2015-08	P. Chen and Ch. Schwab Adaptive Sparse Grid Model Order Reduction for Fast Bayesian Estimation and Inversion
2015-09	J.-L. Bouchot and B. Bykowski and H. Rauhut and Ch. Schwab Compressed Sensing Petrov-Galerkin Approximations for Parametric PDEs
2015-10	A. Jentzen and P. Pusnik Strong convergence rates for an explicit numerical approximation method for stochastic evolution equations with