

Project:

(version of January 8, 2008)

Eigenvalue Problems for the Curl Operator with non Classical Boundary Conditions

Researcher(s): Prof. Dr. R. Hiptmair, SAM, ETH Zürich
Prof. S. Tordeux, INSA Toulouse, France
Prof. R. Kotuiga, Boston University, USA

Funding: no external funding

Duration: ongoing project – started in 2004

The Green's formula for the **curl** operator

$$\int_D \mathbf{curl} \mathbf{u} \cdot \mathbf{v} - \mathbf{curl} \mathbf{v} \cdot \mathbf{u} \, dx = \int_{\partial D} (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{n} \, dS ,$$

illustrates that this first order operator is in fact symmetric once suitable boundary conditions are imposed. Starting from this observation we examine the **curl** operator as an unbounded operator in $\mathbf{L}^2(D)$ and ask which boundary conditions will render this operator self-adjoint.

This question is important in connection with so-called stable force-free magnetic fields, that is, current distributions that are not affected by their own induced magnetic fields. Stable force-free magnetic fields have to satisfy

$$\exists \alpha \in \mathbb{R} \setminus \{0\} : \quad \mathbf{curl} \mathbf{H} = \alpha \mathbf{H} . \quad (1)$$

This is an eigenvalue problem for the **curl** operator.

More precisely, we seek self-adjoint extensions $\mathbf{curl}_s : D_s \subset \mathbf{L}^2(D) \mapsto \mathbf{L}^2(D)$ of the minimal closed **curl** operator with domain $\mathbf{H}_0(\mathbf{curl}, D)$. We adopt the calculus of differential forms on domains D in 3-dimensional space. In this framework, **curl** is treated as operator $\star d$ on 1-forms on D . Here \star is a Hodge operator arising from a metric on D , and d denotes the exterior derivative.

We resort to the connection between self-adjoint extensions of symmetric operators and symplectic geometry. In the case of **curl**, every self-adjoint extension is described as a complete Lagrangian subspace of the trace space $\mathbf{H}(\mathbf{curl}, D)/\mathbf{H}_0(\mathbf{curl}, D)$ of 1-forms on ∂D equipped with the symplectic product

$$[\omega, \eta] := \int_{\partial D} \omega \wedge \eta .$$

It is important to notice that this symplectic pairing is completely metric free.

We identify two important classes of self-adjoint extensions of **curl** through their defining boundary conditions.

1. operators defined on 1-forms with closed traces

$$\mathcal{D}_s := \left\{ \mathbf{u} \in \mathbf{H}(\mathbf{curl}, D) : \mathbf{curl}_\Gamma(\gamma_t \mathbf{u}) = 0 \text{ on } \partial D, \int_\gamma \mathbf{u} \cdot d\vec{s} = 0 \forall \gamma \in \Gamma \right\} ,$$

2. **curl** operators acting on 1-forms with co-closed traces

$$\mathcal{D}_s := \{ \mathbf{u} \in \mathbf{H}(\mathbf{curl}, D) : \operatorname{div}_\Gamma(\gamma_t \mathbf{u}) = 0 \text{ on } \partial D, \int_\gamma \mathbf{u} \cdot d\vec{s} = 0 \forall \gamma \in \Gamma \}.$$

In each case Γ comprises a subset of g generators (1-cycles on ∂D) of the homology group $\mathbb{H}_1(\partial D)$ of order $2g$.

We show that all these self-adjoint **curl** operators possess a complete orthonormal eigen-system. This connects our research with (1).

Related publications.

Report to appear in 2008.