

Symmetric Methods on Manifolds


Florian Landis

Overview

- **Symmetrize Projection Methods**
- Analyze symmetric projection methods
- Simulation: Solve pendulum equations using symmetric projection methods
- **Symmetrize Local Coordinates methods**
- Find two symmetric Lie group methods


Projection Methods and Symmetry

Standard projection methods are not symmetric, because of asymmetric use of projections.



Projection Methods and Symmetry


Standard projection methods are not symmetric, because of asymmetric use of projections.



We look for an algorithm using projections that preserves the symmetry of the basic integrator:


Projection Methods and Symmetry

Standard projection methods are not symmetric, because of asymmetric use of projections.



We look for an algorithm using projections that preserves the symmetry of the basic integrator:

Use: “inverse projection” - *symmetric* integration step - projection



Symmetric Projection Methods

If Φ_h is a symmetric method applied to $\dot{y} = f(y)$, define:

Algorithm 1 (Symmetric Projection Method (SPM)) Assume that $y_n \in \mathcal{M}$. One step $y_n \mapsto y_{n+1}$ is defined as follows:

- $\tilde{y}_n = y_n + G(y_n)^T \mu$ (perturbation step);
- $\tilde{y}_{n+1} = \Phi_h(\tilde{y}_n)$;
- $y_{n+1} = \tilde{y}_{n+1} + G(y_{n+1})^T \mu$ with $g(y_{n+1}) = 0$ (projection step);

where $G(y) = g'(y)$ and the manifold \mathcal{M} is given by $g(y) = 0$.

Symmetric Projection Methods

If Φ_h is a symmetric method applied to $\dot{y} = f(y)$, define:

Algorithm 1 (Symmetric Projection Method (SPM)) Assume that $y_n \in \mathcal{M}$. One step $y_n \mapsto y_{n+1}$ is defined as follows:

- $\tilde{y}_n = y_n + G(y_n)^T \mu$ (perturbation step);
- $\tilde{y}_{n+1} = \Phi_h(\tilde{y}_n)$;
- $y_{n+1} = \tilde{y}_{n+1} + G(y_{n+1})^T \mu$ with $g(y_{n+1}) = 0$ (projection step);

where $G(y) = g'(y)$ and the manifold \mathcal{M} is given by $g(y) = 0$.

Note: μ is determined implicitly by $g(y_{n+1}) = 0$ and

$$\mu_{\text{perturbation}} = \mu_{\text{projection}}.$$

Existence of Numerical Solution

The vector μ and the numerical approximation y_{n+1} are implicitly defined by

$$F(h, y_{n+1}, \mu) = \begin{pmatrix} y_{n+1} - \Phi_h(y_n + G(y_n^T \mu)) - G(y_{n+1})^T \mu \\ g(y_{n+1}) \end{pmatrix} = 0.$$

Since $F(0, y_n, 0) = 0$ and since

$$\frac{\partial F}{\partial (y_{n+1}, \mu)}(0, y_n, 0) = \begin{pmatrix} I & -2G(y_n)^T \\ G(y_n) & 0 \end{pmatrix}$$

is invertible (provided that $G(y_n)$ has full rank), an application of the implicit function theorem proves the existence of the numerical solution for sufficiently small step size h .

The implicit function theorem

Theorem 0 (Implicit Function Theorem) *Let $f(x, y)$ be a function from a neighborhood of (a, b) into a neighborhood of $f(a, b)$. Define $f_a(y) := f(a, y)$ and assume that $df_a|_b$ is an isomorphism.*

Then, there exist neighborhoods U of a and V of $f(a, b)$ and a unique function $\varphi(x, z)$ from $U \times V$ into a neighborhood of b , such that $z = f(x, \varphi(x, z))$. Furthermore

$$d\varphi = \left(\frac{\partial f}{\partial y} \right)^{-1} \left[dz - \frac{\partial f}{\partial x} dx \right].$$

Order of the SPM

For a study of the local error we let $y_n := y(t_n)$ be a value on the exact solution $y(t)$ of

$$\dot{y} = f(y), \quad f(y) \in T_y\mathcal{M}.$$

If the basic method Φ_h is of order p , i.e., if $y(t_n + h) - \Phi_h(y(t_n)) = \mathcal{O}(h^{p+1})$, we have

$$F(h, y(t_{n+1}), 0) = \mathcal{O}(h^{p+1}).$$

Order of the SPM

For a study of the local error we let $y_n := y(t_n)$ be a value on the exact solution $y(t)$ of

$$\dot{y} = f(y), \quad f(y) \in T_y \mathcal{M}.$$

If the basic method Φ_h is of order p , i.e., if $y(t_n + h) - \Phi_h(y(t_n)) = \mathcal{O}(h^{p+1})$, we have

$$F(h, y(t_{n+1}), 0) = \mathcal{O}(h^{p+1}).$$

Compared to

$$F(h, y_{n+1}, \mu) = 0$$

the implicit function theorem yields

$$y_{n+1} - y(t_{n+1}) = \mathcal{O}(h^{p+1}), \quad \mu = \mathcal{O}(h^{p+1}).$$

Symmetry of the method

Exchanging $h \rightarrow -h$ and $y_n \leftrightarrow y_{n+1}$ in the SPM yields

$$\begin{aligned}\tilde{y}_n &= y_{n+1} + G(y_{n+1})^T \mu, & g(y_{n+1}) &= 0, \\ \tilde{y}_{n+1} &= \Phi_{-h}(\tilde{y}_n), \\ y_n &= \tilde{y}_{n+1} + G(y_n)^T \mu, & g(y_n) &= 0.\end{aligned}$$

Renaming *auxiliary* variables $\mu \rightarrow -\mu$ and $\tilde{y}_n \leftrightarrow \tilde{y}_{n+1}$ gives

$$\begin{aligned}\tilde{y}_{n+1} &= y_{n+1} - G(y_{n+1})^T \mu, & g(y_{n+1}) &= 0, \\ \tilde{y}_n &= \Phi_{-h}(\tilde{y}_{n+1}), \\ y_n &= \tilde{y}_n - G(y_n)^T \mu, & g(y_n) &= 0,\end{aligned}$$

which is equivalent to the formulae of the original algorithm provided Φ_h is symmetric.

ρ -Reversibility

We know that for ρ -reversibility, a method Φ_h has to be symmetric and satisfy

$$\rho \circ \Phi_h = \Phi_{-h} \circ \rho.$$

ρ -Reversibility

We know that for ρ -reversibility, a method Φ_h has to be symmetric and satisfy

$$\rho \circ \Phi_h = \Phi_{-h} \circ \rho.$$

For a symmetric projection, this leads to the condition

$$\rho G(y)^T = G(\rho y)^T \sigma \quad \sigma \text{ constant and invertible,}$$

if the integrator Φ_h of the intermediate step is ρ -reversible.

ρ -Reversibility

We know that for ρ -reversibility, a method Φ_h has to be symmetric and satisfy

$$\rho \circ \Phi_h = \Phi_{-h} \circ \rho.$$

For a symmetric projection, this leads to the condition

$$\rho G(y)^T = G(\rho y)^T \sigma \quad \sigma \text{ constant and invertible,}$$

if the integrator Φ_h of the intermediate step is ρ -reversible.

In many interesting cases,

$$g(\rho y) = \sigma^{-T} g(y)$$

holds, which implies $\rho G(y)^T = G(\rho y)^T \sigma$ if $\rho \rho^T = I$.

Modifications

The perturbation and projection steps can be modified without destroying the symmetry.

For example use a constant projection direction:

$$\tilde{y}_n = y_n + A^T \mu, \quad y_{n+1} = \tilde{y}_{n+1} + A^T \mu \quad A \text{ constant.}$$

To guarantee the existence of the numerical solution, $G(y)A^T$ has to be invertible along the solution $y(t)$.

Modifications

The perturbation and projection steps can be modified without destroying the symmetry.

For example use a constant projection direction:

$$\tilde{y}_n = y_n + A^T \mu, \quad y_{n+1} = \tilde{y}_{n+1} + A^T \mu \quad A \text{ constant.}$$

To guarantee the existence of the numerical solution, $G(y)A^T$ has to be invertible along the solution $y(t)$.

For ρ -reversibility, A has to satisfy

$$\rho A^T = A^T \sigma$$

for an invertible matrix σ .

Example: The Pendulum

The pendulum equations in Cartesian coordinates are

$$\begin{aligned}\dot{q}_1 &= p_1, & \dot{p}_1 &= -q_1 \lambda \\ \dot{q}_2 &= p_2, & \dot{p}_2 &= -1 - q_2 \lambda,\end{aligned}$$

with $\lambda = (p^2 - q_2)/q^2$.

- The solution to these equations remains on the manifold

$$\mathcal{M} = \{(q_1, q_2, p_1, p_2) \mid q^2 = 1, q \cdot p = 0\}.$$

Example: The Pendulum

The pendulum equations in Cartesian coordinates are

$$\begin{aligned}\dot{q}_1 &= p_1, & \dot{p}_1 &= -q_1 \lambda \\ \dot{q}_2 &= p_2, & \dot{p}_2 &= -1 - q_2 \lambda,\end{aligned}$$

with $\lambda = (p^2 - q_2)/q^2$.

- The solution to these equations remains on the manifold

$$\mathcal{M} = \{(q_1, q_2, p_1, p_2) \mid q^2 = 1, q \cdot p = 0\}.$$

- The problem is ρ -reversible for both

$$\rho(q_1, q_2, p_1, p_2) = (q_1, q_2, -p_1, -p_2) \quad \text{and}$$

$$\rho(q_1, q_2, p_1, p_2) = (-q_1, q_2, p_1, -p_2).$$

The Pendulum - Projections

Three different ways to project to the Manifold \mathcal{M} were used:

- Orthogonal Projection: $G^T \mu \perp \mathcal{M}$, $G = \begin{pmatrix} 2q_1 & 2q_2 & 0 & 0 \\ p_1 & p_2 & q_1 & q_2 \end{pmatrix}$

The Pendulum - Projections

Three different ways to project to the Manifold \mathcal{M} were used:


- Orthogonal Projection: $G^T \mu \perp \mathcal{M}$, $G = \begin{pmatrix} 2q_1 & 2q_2 & 0 & 0 \\ p_1 & p_2 & q_1 & q_2 \end{pmatrix}$

Coordinate Projection:

$C^T \mu$ with

$$C_{1\pm} = \begin{pmatrix} 0 & \pm 2 & 0 & 0 \\ 0 & 0 & 0 & \pm 1 \end{pmatrix}$$

$$C_{2\pm} = \begin{pmatrix} \pm 2 & 0 & 0 & 0 \\ 0 & 0 & \pm 1 & 0 \end{pmatrix}$$



The Pendulum - Projections

Three different ways to project to the Manifold \mathcal{M} were used:


- Orthogonal Projection: $G^T \mu \perp \mathcal{M}$, $G = \begin{pmatrix} 2q_1 & 2q_2 & 0 & 0 \\ p_1 & p_2 & q_1 & q_2 \end{pmatrix}$

Coordinate Projection:

$C^T \mu$ with

$$C_{1\pm} = \begin{pmatrix} 0 & \pm 2 & 0 & 0 \\ 0 & 0 & 0 & \pm 1 \end{pmatrix}$$




$$C_{2\pm} = \begin{pmatrix} \pm 2 & 0 & 0 & 0 \\ 0 & 0 & \pm 1 & 0 \end{pmatrix}$$





- Projection violating ρ -reversibility: $A = \begin{pmatrix} -\varepsilon & -2 & 0 & 0 \\ -\varepsilon & 0 & 0 & -1 \end{pmatrix}$

The Pendulum - Results

Low Starting Position



High Starting Position



Local Coordinate Methods (LCM)

Algorithm 1 (Symmetric Local Coordinates Approach) Assume:

$y_n \in \mathcal{M}$ and ψ_a a local parametrization of \mathcal{M} with $\psi_a(0) = a$ (close to y_n).

- find z_n (close to 0) such that $\psi_a(z_n) = y_n$;
- $\tilde{z}_{n+1} = \Phi_h(z_n)$ (symmetric one-step method applied to $\dot{z} = \psi'(z)^+ f(\psi(z))$).
- $y_{n+1} = \psi_a(\tilde{z}_{n+1})$;
- choose a in the parametrization such that $z_n + \tilde{z}_{n+1} = 0$.



Symmetric use of local tangent space parametrization

Reversibility of Symmetric LCM

A symmetric local coordinates method is ρ -reversible, if the parametrization is s.t.:

$$\rho\psi_a(z) = \psi_{\rho a}(\sigma z)$$

for some invertible σ .

Reversibility of Symmetric LCM

A symmetric local coordinates method is ρ -reversible, if the parametrization is s.t.:

$$\rho\psi_a(z) = \psi_{\rho a}(\sigma z)$$

for some invertible σ .

If the initial problem is ρ -reversible, this implies σ -reversibility for

$$\dot{z} = \psi'(z)^+ f(\psi(z)).$$

The basic method Φ_h must therefore be σ -reversible.

Symmetric Lie Group Methods

Now, consider

$$\dot{Y} = A(Y)Y, \quad Y(0) = Y_0,$$

where $A(Y)$ is in the Lie algebra \mathcal{G} whenever Y is in the Lie group G .

Munthe-Kaas methods are in general not symmetric (asymmetric use of the local coordinates $Y = \exp(\Omega)Y_0$).

Symmetric Lie Group Methods

Now, consider

$$\dot{Y} = A(Y)Y, \quad Y(0) = Y_0,$$

where $A(Y)$ is in the Lie algebra \mathcal{G} whenever Y is in the Lie group G .

Munthe-Kaas methods are in general not symmetric (asymmetric use of the local coordinates $Y = \exp(\Omega)Y_0$).

Accidentally, the Lie group method based on the implicit midpoint rule


$$Y_{n+1} = \exp(\Omega)Y_n, \quad \Omega = hA(\exp(\Omega/2)Y_n)$$

is symmetric (exchange $h \leftrightarrow -h$, $Y_n \leftrightarrow Y_{n+1}$ and the auxiliary variable $\Omega \leftrightarrow -\Omega$).

Symmetric Munthe-Kaas Methods

According to the symmetric LCM, we choose a local parametrization

$$\psi_U(\Omega) = \exp(\Omega)U,$$




where $U = \exp(\Theta)Y_n$ plays the role of the midpoint on the manifold.

We put $Z_n = -\Theta$ so that $\psi_U(Z_n) = Y_n$.

Symmetric Munthe-Kaas Methods

According to the symmetric LCM, we choose a local parametrization

$$\psi_U(\Omega) = \exp(\Omega)U,$$



where $U = \exp(\Theta)Y_n$ plays the role of the midpoint on the manifold.

We put $Z_n = -\Theta$ so that $\psi_U(Z_n) = Y_n$.

Apply any symmetric Runge-Kutta method to the differential equation


$$\dot{\Omega} = A(\psi_U(\Omega)) + \sum_{k=1}^q \frac{B_k}{k!} \mathbf{ad}_{\Omega}^k (A(\psi_U(\Omega))), \quad \Omega(0) = -\Theta,$$

to obtain \tilde{Z}_{n+1} from Z_n .

Symmetric Munthe-Kaas Methods

According to the symmetric LCM, we choose a local parametrization

$$\psi_U(\Omega) = \exp(\Omega)U,$$



where $U = \exp(\Theta)Y_n$ plays the role of the midpoint on the manifold.

We put $Z_n = -\Theta$ so that $\psi_U(Z_n) = Y_n$.

Apply any symmetric Runge-Kutta method to the differential equation

$$\dot{\Omega} = A(\psi_U(\Omega)) + \sum_{k=1}^q \frac{B_k}{k!} \mathbf{ad}_{\Omega}^k (A(\psi_U(\Omega))), \quad \Omega(0) = -\Theta,$$

to obtain \tilde{Z}_{n+1} from Z_n .

Θ is implicitly given by $Z_n + \tilde{Z}_{n+1} = 0$, and the numerical result is

$$Y_{n+1} = \psi_U(\tilde{Z}_{n+1}) = \exp(\tilde{Z}_{n+1}) \exp(\Theta)Y_n = \exp(2\Theta)Y_n.$$

2-stage Gauss on Lie Groups

2-stage Gauss:

$$\begin{array}{c|cc}
 \frac{1}{2} - \frac{\sqrt{3}}{6} & \frac{1}{4} & \frac{1}{4} - \frac{\sqrt{3}}{6} \\
 \frac{1}{2} + \frac{\sqrt{3}}{6} & \frac{1}{4} + \frac{\sqrt{3}}{6} & \frac{1}{4} \\
 \hline
 & \frac{1}{2} & \frac{1}{2}
 \end{array}$$

With the coefficients of the 2-stage Gauss method and with $q = 1$, $B_1 = -\frac{1}{2}$ we thus get:

$$k_1 = A \left(\exp \left(\underbrace{-\Theta + \frac{h}{4}(k_1 + k_2) - h\frac{\sqrt{3}}{6}k_2}_{\Omega_1} \right) U \right) - \frac{1}{2} \left[\Omega_1, A(\exp(\Omega_1)U) \right]$$

$$k_2 = A \left(\exp \left(\underbrace{-\Theta + \frac{h}{4}(k_1 + k_2) + h\frac{\sqrt{3}}{6}k_1}_{\Omega_2} \right) U \right) - \frac{1}{2} \left[\Omega_2, A(\exp(\Omega_2)U) \right]$$

$$\Theta = -\Theta + \frac{h}{2}(k_1 + k_2).$$

2-stage Gauss on Lie Groups

With

$$2\Theta = \frac{h}{2}(k_1 + k_2) \Rightarrow \Omega_1 = -h\frac{\sqrt{3}}{6}k_2, \Omega_2 = h\frac{\sqrt{3}}{6}k_1$$

and $A_i := A(\exp(\Omega_i)U)$ we can rewrite this as

$$\Omega_1 = -h\frac{\sqrt{3}}{6} \left(A_2 - \frac{1}{2}[\Omega_2, A_2] \right)$$

$$\Omega_2 = h\frac{\sqrt{3}}{6} \left(A_1 - \frac{1}{2}[\Omega_1, A_1] \right)$$

$$\begin{aligned} Y_{n+1} &= \exp(2\Theta)Y_n \\ &= \exp \left(\frac{h}{2}(A_1 + A_2) - \frac{h}{4}([\Omega_1, A_1] + [\Omega_2, A_2]) \right) Y_n. \end{aligned}$$

2-stage Gauss on Lie Groups

Neglecting terms of size $\mathcal{O}(h^4)$ in Y_{n+1} gives

$$\Omega_1 = -h \frac{\sqrt{3}}{6} A_2 + \frac{h^2}{24} [A_1, A_2]$$

$$\Omega_2 = h \frac{\sqrt{3}}{6} A_1 - \frac{h^2}{24} [A_1, A_2]$$

$$Y_{n+1} = \exp \left(\frac{h}{2} (A_1 + A_2) - h^2 \frac{\sqrt{3}}{12} [A_1, A_2] \right) Y_n.$$

This method is symmetric and therefore still of order 4 (orders of symmetric methods are even).

Reversibility of 2-stage Gauss

For any linear invertible transformation ρ , the parametrization $\psi_U(\Omega) = \exp(\Omega)U$ satisfies

$$\rho\psi_U(\Omega) = \rho \exp(\Omega)U = \exp(\rho\Omega\rho^{-1})\rho U = \psi_{\rho U}(\sigma\Omega)$$

with $\sigma\Omega = \rho\Omega\rho^{-1}$.

Reversibility of 2-stage Gauss

For any linear invertible transformation ρ , the parametrization $\psi_U(\Omega) = \exp(\Omega)U$ satisfies

$$\rho\psi_U(\Omega) = \rho \exp(\Omega)U = \exp(\rho\Omega\rho^{-1})\rho U = \psi_{\rho U}(\sigma\Omega)$$

with $\sigma\Omega = \rho\Omega\rho^{-1}$.

If the initial problem is ρ -reversible, i.e. $\rho A(Y) = -A(\rho Y)\rho$, then

$$\dot{\Omega} = A(\psi_U(\Omega)) + \sum_{k+1}^q \frac{B_k}{k!} \mathbf{ad}_{\Omega}^k (A(\psi_U(\Omega))), \quad \Omega(0) = -\Theta$$

is σ -reversible for all truncation indices q .

Summary

- symmetric projection methods
 - have the same order as basic integrator
 - are ρ -reversible if the basic method is ρ -reversible and $\rho G(y)^T = G(\rho y)^T \sigma$ for some invertible σ .
- symmetric local coord. methods
 - (have the same order as basic integrator)
 - are ρ -reversible if $\rho\psi_a(z) = \psi_{\rho a}(\sigma z)$ for some invertible σ and if the basic method is σ -reversible.
- symmetric Lie group methods can be obtained by symmetrizing Munte-Kaas methods. Examples:
 - 2nd order: implicit midpoint rule as basic integrator
 - 4th order: 2-stage Gauss as basic integrator