

# Numerical solution of matrix eigenvalue problems

## Part 1: Power method and friends

Daniel Kressner  
ETH Zürich

kressner@math.ethz.ch  
<http://www.math.ethz.ch/~kressner>

ZSS 2008



## Outline

- ▶ powermethod
- ▶ subspaceiteration
- ▶ inverseiteration
- ▶ Rayleigh-quotientiteration



Input: **start vector**  $v_0 \in \{\mathbb{R}, \mathbb{C}\}^n$ , **matrix**  $A \in \{\mathbb{R}, \mathbb{C}\}^{n \times n}$ .  
**for**  $i = 0, 1, 2, \dots$  **do**  
 $v_{i+1} = Av_i$   
**end for**

Let  $\lambda_1, \dots, \lambda_n$  be eigenvalues of  $A$  s.t.

$$|\lambda_1| > |\lambda_2| \geq \dots \geq |\lambda_n|$$

with corresponding eigenvectors  $x_1, x_2, \dots, x_n$ .

Power method **converges to**  
 $x_1$ :

$$\theta(x_1, v_i) = O\left(\left|\frac{\lambda_2}{\lambda_1}\right|^i\right)$$

Angle between vectors  $v, w \in \mathbb{C}^n$ :

$$\cos \theta(v, w) = \frac{|w^H v|}{\|v\|_2 \|w\|_2}$$



## The power method

Input: **start vector**  $v_0 \in \{\mathbb{R}, \mathbb{C}\}^n$ , **matrix**  $A \in \{\mathbb{R}, \mathbb{C}\}^{n \times n}$ .  
Output: Approximation  $v_i$  to **dominant eigenvector**.  
**for**  $i = 0, 1, 2, \dots$  **do**  
 $\tilde{v}_{i+1} = Av_i$   
 $v_{i+1} = \tilde{v}_{i+1} / \|\tilde{v}_{i+1}\|_2$   
**end for**

To get an approximation to  $\lambda_1$ , choose  $\mu_i$  s.t.

$$\|Av_i - \mu_i v_i\|_2 \rightarrow \min$$

$\rightsquigarrow$  **Rayleigh quotient**  $\mu_i := v_i^H Av_i = v_i^H \tilde{v}_{i+1}$ .

$$|\mu_i - \lambda_1| = O\left(\left|\frac{\lambda_2}{\lambda_1}\right|^i\right).$$

For **symmetric**  $A$ :

$$|\mu_i - \lambda_1| \leq |\lambda_n - \lambda_1| \sin^2 \theta(x_1, v_0) \left|\frac{\lambda_2}{\lambda_1}\right|^{2k}.$$



Termination? Is  $|\mu_{i+1} - \mu_i| \approx 0$  a suitable criterion?

- ▶ For symmetric/normal matrices, **yes**:

$$|\mu_{i+1} - \mu_i| \leq \text{tol} \quad \Rightarrow \quad |\mu_i - \lambda| \leq \frac{1}{1 - |\lambda_2/\lambda_1|} \text{tol}.$$

- ▶ For nonnormal matrices, **no**. Transient growth! Show transient.m

Only reasonable choice:  $r := Av_i - \mu_i v_i \approx 0$ .

**Backward error** interpretation:  $(\mu_i, v_i)$  *exact* eigenvalue/eigenvector pair of *perturbed* matrix

$$A + \Delta A, \quad \Delta A := -rv_i^H.$$

With this choice,  $\|\Delta A\|_2 = \|r\|_2$ .



## Snippets of perturbation analysis

First-order influence of  $\Delta A$  on accuracy of eigenvalues/eigenvectors:

$$Ax = \lambda x, \quad \|x\|_2 = 1.$$

↓

$$(A + \epsilon \Delta A)x(\epsilon) = \lambda(\epsilon)x(\epsilon), \quad \|x(\epsilon)\|_2 = 1.$$

↓

$$\dot{\lambda}(0) = \frac{y^H \Delta A x}{y^H x},$$

where  $y^H A = \lambda y^H$ .



For nonsymmetric matrices:

$$|\mu_i - \lambda_1| \leq \frac{1}{|y_1^H x_1|} \|\Delta A\|_2 + O(\|\Delta A\|_2^2)$$

and

$$\theta(v_i, \theta_1) \leq \|[(I - x_1 x_1^H)(A - \lambda_1 I)(I - x_1 x_1^H)]^\dagger\|_2 \|\Delta A\|_2 + O(\|\Delta A\|_2^2).$$

If  $A$  is symmetric:

$$|\mu_i - \lambda_1| \leq \|\Delta A\|_2.$$

and

$$\sin \theta(x_1, v_i) = \frac{1}{|\mu_i - \lambda_2|} \|\Delta A\|_2.$$



## A posteriori estimates

- ▶ If  $A$  is nonsymmetric, apply power method to  $A$  and  $A^H$ . Obtain approximations  $v_i$  and  $w_i$  to right/left eigenvectors  $x_1, y_1$ . Then

$$|\mu_i - \lambda_1| \approx \frac{1}{|w_i^H v_i|} \max\{\|Av_i - \mu_i v_i\|_2, \|w_i^H A - \mu_i w_i^H\|_2\}.$$

- ▶ If  $A$  is symmetric,

$$|\mu_i - \lambda_i| \leq \|r\|_2, \quad \sin \theta(x_1, v_i) = \frac{1}{|\mu_i - \lambda_2|} \|r\|_2.$$

- ▶ Alternative for symmetric  $A$ : **Kato-Temple inequality**. Assume  $\exists$  interval  $(a, b)$  containing  $\mu_i, \lambda_1$  and **no other eigenvalue** of  $A$ . Then

$$-\frac{\|r\|_2^2}{\mu_i - a} \leq \mu_i - \lambda_i \leq \frac{\|r\|_2^2}{b - \mu_i}.$$





Let  $A$  be **symmetric** and order eigenvalues

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n.$$

**Minimax theorems:**

$$\lambda_j = \max_{\dim(\mathcal{V})=j} \min_{\substack{v \in \mathcal{V} \\ \|v\|_2=1}} v^T A v,$$
$$\lambda_{n-j+1} = \min_{\dim(\mathcal{V})=j} \max_{\substack{v \in \mathcal{V} \\ \|v\|_2=1}} v^T A v.$$

As a consequence, Ritz values **interlace** eigenvalues of  $A$ :

$$\lambda_j \geq \mu^{(j)} \geq \lambda_{n-j+1},$$

↔ Ritz values approach eigenvalues from below.



## Inverse iteration

Eigenvalue closest to  $\tau \in \mathbb{C}$  is dominant eigenvalue of

$$(A - \tau I)^{-1}.$$

Input: **start vector**  $v_0$ , **matrix**  $A$ , **target**  $\tau \in \mathbb{C}$ .

Output: Approximation  $(\mu_i, v_i)$  to eigenpair closest to  $\tau$ .

**for**  $i = 0, 1, 2, \dots$  **do**

    Solve  $(A - \tau I)\tilde{v}_{i+1} = v_i$

$v_{i+1} = \tilde{v}_{i+1} / \|\tilde{v}_{i+1}\|_2$

$\mu_{i+1} = (\tilde{v}_{i+1}^H v_{i+1})^{-1} + \tau$

**end for**

**Not suited for inexact application of  $(A - \tau I)^{-1}$ !**



## Order eigenvalues

$$|\lambda_1 - \tau| \geq \dots \geq |\lambda_{n-1} - \tau| > |\lambda_n - \tau|.$$

Convergence to  $(\lambda_n, \mathbf{x}_n)$ :

$$|\mu_i - \lambda_n| = O\left(\left|\frac{\lambda_n - \tau}{\lambda_{n-1} - \tau}\right|^i\right)$$

$$\theta(\mathbf{x}_n, \mathbf{v}_i) = O\left(\left|\frac{\lambda_n - \tau}{\lambda_{n-1} - \tau}\right|^i\right)$$



## Rayleigh-quotient iteration

**Tempting** idea: Improve convergence by adjusting  $\tau$ .

Input: **start vector**  $\mathbf{v}_0$  with  $\|\mathbf{v}_0\|_2 = 1$ , **matrix**  $A$ .

Output: Approximation  $(\mu_i, \mathbf{v}_i)$  to **some** eigenpair.

**for**  $i = 0, 1, 2, \dots$  **do**

    Set  $\tau = \mathbf{v}_i^H A \mathbf{v}_i$ .

    Solve  $(A - \tau I) \tilde{\mathbf{v}}_{i+1} = \mathbf{v}_i$

$\mathbf{v}_{i+1} = \tilde{\mathbf{v}}_{i+1} / \|\tilde{\mathbf{v}}_{i+1}\|_2$

$\mu_{i+1} = \mathbf{v}_{i+1}^H A \mathbf{v}_{i+1}$

**end for**

**Local convergence quadratic** or even cubic (when  $A$  normal).

**Global convergence critical!**

Start rqiglobal.m



Natural generalization to subspaces.

Input: start ONB  $V_0$ , symmetric matrix  $A$ .

Output: Approximate ONB  $V_i$  to some invariant subspace.

**for**  $i = 0, 1, 2, \dots$  **do**

Set  $T = V_i^H A V_i$ .

Solve Sylvester equation  $A \tilde{V}_{i+1} - \tilde{V}_{i+1} T = V_i$ .

Compute ONB  $V_{i+1}$  of  $\tilde{V}_{i+1}$ .

**end for**

**Local convergence cubic.** Advantageous over RQI, when several eigenvalues are needed:

- ▶ Avoid repeated convergence to same eigenvalues.
- ▶ In the limit, effective conditioning of  $X \mapsto AX - XT$  only depends on external gap.
- ▶ Domain of attraction primarily depends on external gap.



## Literature

- ▶ Virtually every book on numerical linear algebra covers the power method. The subspace iteration is covered, e.g., in [G. H. Golub and C. F. Van Loan. Matrix Computations. 1996], [G. W. Stewart. Matrix Algorithms. Vol. II. 2001].
- ▶ An implementation of the subspace iteration can be found in [Z. Bai and G. W. Stewart. Algorithm 776. SRRIT – A FORTRAN subroutine to calculate the dominant invariant subspaces of a nonsymmetric matrix. ACM TOMS, 23:494–513, 1998].
- ▶ The standard reference on the perturbation of *nonsymmetric* eigenvalue problems is [G. W. Stewart and J.-G. Sun. Matrix Perturbation Theory. Academic Press, New York, 1990].
- ▶ An overview of available relative perturbation bounds for *symmetric* eigenvalue problems is given in [I. C. F. Ipsen. Relative Perturbation Bounds for Matrix Eigenvalues and Singular Values. Acta Numerica, pp 151–201, (1998)].
- ▶ The Grassmann RQI and related methods are described in the recent monograph [P.-A. Absil, R. Mahony, R. Sepulchre. Optimization Algorithms on Matrix Manifolds. Princeton University Press, 2008].

