

Sparse Two-Scale FEM for Homogenization Problems

A.-M. Matache

Research Report No. 2001-09
October 2001

Seminar für Angewandte Mathematik
Eidgenössische Technische Hochschule
CH-8092 Zürich
Switzerland

Sparse Two-Scale FEM for Homogenization Problems

A.-M. Matache

Seminar für Angewandte Mathematik
Eidgenössische Technische Hochschule
CH-8092 Zürich
Switzerland

Research Report No. 2001-09

October 2001

Abstract

We analyze two-scale Finite Element Methods for the numerical solution of elliptic homogenization problems with coefficients oscillating at a small length scale $\varepsilon \ll 1$. Based on a refined two-scale regularity on the solutions, two-scale tensor product FE spaces are introduced and error estimates which are robust (i.e. independent of ε) are given. We show that under additional two-scale regularity assumptions on the solution, resolution of the fine scale is possible with substantially fewer degrees of freedom and the two-scale full tensor product spaces can be “thinned out” by means of *sparse* interpolation preserving at the same time the error estimates.

Keywords: Homogenization; two-scale FEM; sparse two-scale FEM

1 Introduction

The accurate and efficient numerical solution of partial differential equations with coefficients or geometries that oscillate periodically at a small length scale $\varepsilon \ll 1$ has received increasing attention recently. Solutions to such problems contain several length scales that differ by many orders of magnitude.

Such homogenization problems have been thoroughly analyzed by asymptotic analysis as $\varepsilon \rightarrow 0$, see [3, 7] and the references there. In this analytical approach the limiting problem as $\varepsilon \rightarrow 0$ is identified first and then solved numerically by standard methods. In particular, no scale resolution is required, since the fine scales are averaged out (homogenized). However, this analytic homogenization process does not preserve the fine scale information of the solution and the recovery of fine scale features by use of correctors proves as costly as the original problem.

The direct numerical treatment of multiple scale problems by the standard Finite Element Method (FEM) [1, 9] on the other hand faces the difficulty of representing the microstructure. The standard FEM yields reliable results only under the assumption of *scale resolution*, i.e. if the FE mesh is refined to the smallest length scale. Such an approach is however infeasible if the difference in scales is sufficiently large. However, a main feature of such problems is that the spatial variation of the solutions is concentrated at length scales which are a-priori known or can be estimated. Moreover, the regular patterns contained in the fine scale data can be used to substantially lower the number of degrees of freedom requested by the resolution of this scale.

1.1 The Homogenization Problem

We consider here the following elliptic problem in divergence form

$$L^\varepsilon \left(\frac{x}{\varepsilon}, \partial_x \right) u^\varepsilon := -\nabla \cdot \left(A \left(\frac{x}{\varepsilon} \right) \nabla u^\varepsilon \right) + a_0 \left(\frac{x}{\varepsilon} \right) u^\varepsilon = f(x), \quad (1)$$

where ε is a small parameter and we assume that $A(y)$, $a_0(y)$ are 1-periodic in each variable and that $A(\cdot) \in L^\infty_{\text{per}}(\widehat{Q})^{n \times n}_{\text{symm}}$, $a_0(\cdot) \in L^\infty_{\text{per}}(\widehat{Q})$ satisfy, for some $\gamma > 0$, $\xi^\top A(y) \xi \geq \gamma |\xi|^2$, $a_0(y) \geq \gamma$ for all $\xi \in \mathbb{R}^n$ and a.e. $y \in \widehat{Q} = [0, 1]^n$. The domain $\widehat{Q} = [0, 1]^n$ will be referred to as unit-cell domain. We consider (1) in a bounded Lipschitz domain Ω and we complete (1) by Dirichlet boundary conditions on $\partial\Omega$, i.e.,

$$u^\varepsilon = 0 \quad \text{on } \partial\Omega. \quad (2)$$

1.2 Finite Element Approximation

The FEM is based on the variational form of (1), (2)

$$\text{Find } u^\varepsilon \in H_0^1(\Omega) : B^\varepsilon(u^\varepsilon, v) = (f, v) \quad \forall v \in H_0^1(\Omega), \quad (3)$$

where the bilinear form $B^\varepsilon : H_0^1(\Omega) \times H_0^1(\Omega) \rightarrow \mathbb{R}$ is given by

$$B^\varepsilon(u, v) = \int_{\Omega} \left(A \left(\frac{x}{\varepsilon} \right) \nabla u(x) \right) \cdot \nabla v(x) + a_0 \left(\frac{x}{\varepsilon} \right) u(x)v(x) dx.$$

The variational problem (3) admits for every $\varepsilon > 0$ and every $f \in L^2(\Omega)$ a unique solution $u^\varepsilon \in H_0^1(\Omega)$.

Let $V_N^\varepsilon \subset H_0^1(\Omega)$ be any subspace of dimension $N = \dim(V_N^\varepsilon) < \infty$. Then

$$u_N^\varepsilon \in V_N^\varepsilon : B^\varepsilon(u_N^\varepsilon, v) = (f, v) \quad \forall v \in V_N^\varepsilon \quad (4)$$

defines a unique FE solution and there exists $C > 0$ independent of ε such that

$$\|u^\varepsilon - u_N^\varepsilon\|_{H^1(\Omega)} \leq C \min_{v \in V_N^\varepsilon} \|u^\varepsilon - v\|_{H^1(\Omega)}. \quad (5)$$

Even if the right hand side f , the domain Ω and the coefficients A and a_0 are smooth (i.e., C^∞), if $\varepsilon/\text{diam}(\Omega) \ll 1$ the solution u^ε exhibits oscillations on the ε -scale obstructing FE convergence. More specifically, there exist positive constants $C = C(\Omega)$ and $C(\alpha) = C(\alpha, \Omega)$, $\alpha \in \mathbb{N}^n$, such that

$$\|u^\varepsilon\|_{L^2(\Omega)} \leq C, \quad \|D^\alpha u^\varepsilon\|_{L^2(\Omega)} \leq C(\alpha)\varepsilon^{1-|\alpha|}, \quad \forall \alpha \in \mathbb{N}^n, |\alpha| > 0. \quad (6)$$

Let us denote by $V_N = S^{p,1}(\Omega, \mathcal{T}_H)$ the usual FE space of continuous, piecewise polynomials of degree $p \geq 1$ on a quasiuniform mesh \mathcal{T}_H of meshwidth H . Then, the FE error with respect to V_N satisfies the following *a-priori* estimate

$$\|u^\varepsilon - u_N^\varepsilon\|_{H^1(\Omega)} \leq C \min(1, (H/\varepsilon)^p),$$

with $C > 0$ being a constant independent of ε and H . Standard FEM, as e.g., piecewise linears on a quasiuniform mesh \mathcal{T}_H of size H , thus converge only if $H < \varepsilon$, i.e., if $N = \dim V_N^\varepsilon = O(\varepsilon^{-n})$. This *scale resolution* requirement is, especially if $n = 3$, computationally very expensive.

In view of (5), the key to a robust discretization of (1) is the design of V_N^ε . Rather than incorporating e.g., the asymptotics of u^ε (which are not always defined, see [7] and the references there) into V_N^ε , we design V_N^ε based on a refined two-scale regularity theory on u^ε . In contrast, in [5, 4, 8] a generalized FEM based on non-polynomial FE spaces was proposed and analyzed. For analytic f , this method was shown to give robust exponential rates of convergence.

1.3 Two-Scale Regularity

On the unbounded domain \mathbb{R}^n (i.e., in the absence of boundary layers) the solution $u^\varepsilon(x)$ can be viewed as a map from the ‘slow’ variable x into the ‘fast’ variable x/ε : $u^\varepsilon(x) = U^\varepsilon(x, x/\varepsilon)$, where $U^\varepsilon(x, y)$ depends smoothly on ε . In [6] we derived new, two-scale regularity results on $u^\varepsilon(x)$ by analyzing $U^\varepsilon(x, y)$. More precisely, the following two-scale shift theorem holds:

Theorem 1.1 *Assume that $A(\cdot)$, $a_0(\cdot)$ are smooth and 1-periodic in $y = x/\varepsilon \in \widehat{Q}$. Then, for $f \in H^k(\mathbb{R}^n)$ ($k \geq 0$), the solution $u^\varepsilon(x)$ of (1) on \mathbb{R}^n can be written as $u^\varepsilon(x) = U^\varepsilon(x, y)|_{y=x/\varepsilon}$, $x \in \mathbb{R}^n$, where $U^\varepsilon(x, y)$ satisfies in $\Omega = \mathbb{R}^n$ the two-scale regularity estimate*

$$\|U^\varepsilon\|_{H^r(\Omega, H_{\text{per}}^s(\widehat{Q}))} \leq C(k) \|f\|_{H^{r+s-1}(\Omega)} \quad (7)$$

provided $r + s \leq k + 1$, $r, s \geq 0$, and

$$\|\varepsilon^{-1} \nabla_y U^\varepsilon\|_{H^r(\Omega, H_{\text{per}}^{s-1}(\widehat{Q}))} \leq C(k) \|f\|_{H^{r+s-1}(\Omega)} \quad (8)$$

provided $r + s \leq k + 1$, $r, s - 1 \geq 0$. Here, $C(k)$ is independent of ε , but depends on $r + s$.

This two-scale point of view of regularity gives rise to a ‘natural’ FE discretization of (1) by means of a non-standard two-scale FE-space V_N^ε in Ω . With the two-scale FE spaces V_N^ε robust convergence rates as $h, H \rightarrow 0$ can be achieved for u_N^ε as we shall show in Section 2. These two-scale approximation results are quite general and applicable whenever the solution has the two-scale regularity. Section 2 is devoted to the definition and error analysis of the two-scale FEM and its sparse version. In Section 3 we present numerical results which support our error estimates.

2 Two-Scale Finite Element Method

2.1 FE-Spaces

We assume that the domain Ω is axis-parallel and we take \mathcal{T}_H to be a quasiuniform mesh in Ω of affine quadrilateral elements of size H . We take as macro FE space in $\Omega = (0, 1)^n$ the standard FE space $S^p(\Omega, \mathcal{T}_H) = \{u \in H^1(\Omega) \mid u|_K \circ F_K^{-1} \in S^p(\widehat{K}) \forall K \in \mathcal{T}_H\}$, where $F_K : \widehat{K} \rightarrow K$ is the affine element map associated to the element K and $S^p(\widehat{K})$ denotes the space of polynomials of degree p on the reference element $\widehat{K} = (0, 1)^n$.

Next, we resolve the fine scale by a FEM in \widehat{Q} based on a periodic mesh $\widehat{\mathcal{T}}_h$, for simplicity also affine, quasiuniform of width h . Then, we denote by $S_{\text{per}}^\mu(\widehat{Q}, \widehat{\mathcal{T}}_h)$ the finite element space of all continuous, periodic piecewise polynomials of degree μ with respect to $\widehat{\mathcal{T}}_h$.

We take as two-scale FE space V_N^ε the space of traces of the two-scale space $S^p(\Omega, \mathcal{T}_H; S_{\text{per}}^\mu(\widehat{Q}, \widehat{\mathcal{T}}_h)) = \{U(x, y) \mid \forall K \in \mathcal{T} : U(F_K(\widehat{x}), y) \text{ is polynomial of degree } p \text{ w.r.t. } \widehat{x} \text{ in } \widehat{K} \text{ and continuous, periodic p.w. polynomial w.r.t. } \widehat{\mathcal{T}}_h \text{ in } y \in \widehat{Q}\}$. More specifically,

$$V_N^\varepsilon = \mathcal{R}^\varepsilon S^p(\Omega, \mathcal{T}_H; S_{\text{per}}^\mu(\widehat{Q}, \widehat{\mathcal{T}}_h)), \quad (9)$$

where the trace operator \mathcal{R}^ε is given by $(\mathcal{R}^\varepsilon U)(x) = U(x, y)|_{y=\frac{x}{\varepsilon}}$. Since $\{1\} \subset S_{\text{per}}^\mu(\widehat{Q}, \widehat{\mathcal{T}}_h)$, V_N^ε is a generalized FE-space. Note that the elements of the FE space V_N^ε have the form $u_{FE}^\varepsilon(x) = \sum_{i,I} c_{iI} N_i(x) \phi_I(x/\varepsilon)$, $x \in \Omega$, with $c_{iI} \in \mathbb{R}$ and shape functions $N_i(\cdot) \in S^p(\Omega, \mathcal{T}_H)$, $\phi_I(\cdot) \in S_{\text{per}}^\mu(\widehat{Q}, \widehat{\mathcal{T}}_h)$.

2.2 Two-Scale Finite Element Convergence

The goal of this section is to estimate the approximation error (5) for the two-scale FE space V_N^ε (9) and to obtain robust estimates with respect to ε for $H/\varepsilon \geq 1$.

We have seen that the solution u^ε may be interpreted as $u^\varepsilon(\cdot) = \mathcal{R}U^\varepsilon(\cdot, \cdot/\varepsilon)$, where $U^\varepsilon(\cdot, \cdot)$ is defined on $\Omega \times \widehat{Q}$ and \mathcal{R} is the trace operator given by $(\mathcal{R}f)(x) = f(x, x)$. This suggests to use FE-interpolants in Ω and \widehat{Q} to approximate U^ε in $\Omega \times \widehat{Q}$ and take traces.

If H denotes the mesh size of the quasiuniform ‘macroscopic’ triangulation on Ω and h is the mesh size of the quasiuniform ‘micro’ triangulation on the unit cell \widehat{Q} , we obtain (see [6]):

Proposition 2.1 *For $p, \mu, k, s \geq 1$ and $H/\varepsilon \in \mathbb{N}$ the error in the two-scale FEM based on the space (9) can be estimated as follows:*

$$\begin{aligned} \|u^\varepsilon - u_{FE}^\varepsilon\|_{H^1(\Omega)} &\leq \\ &\leq CH^{\min(p,k)} \Phi_n(p, k) (\|\varepsilon^{-1} \nabla_y U^\varepsilon\|_{H^k(\Omega; L_{\text{per}}^2(\widehat{Q}))} + \|U^\varepsilon\|_{H^{k+1}(\Omega; L_{\text{per}}^2(\widehat{Q}))}) \\ &+ Ch^{\min(\mu,s)} \Phi_n(\mu, s) (\|\varepsilon^{-1} \nabla_y U^\varepsilon\|_{H^n(\Omega; H_{\text{per}}^s(\widehat{Q}))} + \|U^\varepsilon\|_{H^n(\Omega; H_{\text{per}}^{s+1}(\widehat{Q}))}). \end{aligned} \quad (10)$$

Here $C > 0$ is a positive constant independent of p, μ, k, s and ε and the functional $\Phi_n(p, k)$ depends only on n, p and k . For $p \rightarrow \infty$, $\Phi_n(p, k) \leq Cp^{-(k-n+1)}$.

The following result is a consequence of Proposition 2.1 and Theorem 1.1.

Theorem 2.2 *Assume for the solution u^ε of (3) the two-scale regularity (7)–(8) in Ω . Then, for $H/\varepsilon \in \mathbb{N}$, it holds:*

$$\begin{aligned} \|u^\varepsilon - u_{FE}^\varepsilon\|_{H^1(\Omega)} &\leq C_1(k) H^{\min(p,k)} \Phi_n(p, k) \|f\|_{H^k(\Omega)} \\ &+ C_2(s) h^{\min(\mu,s)} \Phi_n(\mu, s) \|f\|_{H^{n+s}(\Omega)}. \end{aligned}$$

So far we have only discussed the preasymptotic case when $H \geq \varepsilon$ and we obtained the robust (in ε) error estimate $\|e_N^\varepsilon\|_{H^1(\Omega)} \leq C(H^p + h^\mu)$. Let us choose $p = \mu$. Then, for $h \cong H$ we obtain $\|e_N^\varepsilon\|_{H^1(\Omega)} \leq Ch^p$ and $N = \dim(V_N^\varepsilon) = O(H^{-n}h^{-n}) = O(h^{-2n})$. Hence, in terms of number of degrees of freedom, the two-scale FE error estimate is qualitatively of the form

$$\|e_N^\varepsilon\|_{H^1(\Omega)} \leq CN^{-\frac{p}{2n}} \quad \text{for } N \preceq \varepsilon^{-2n}, \quad (11)$$

since the total number of degrees of freedom at the critical value $H \cong \varepsilon$, $h \cong \varepsilon$ is $N = O(\varepsilon^{-2n})$. At this transition point the fine scale is resolved and we switch from the two scale FE space to full discretization with mesh size $H = \varepsilon h$, $h \leq \varepsilon$. This is achieved by breaking the periodicity to get the full space with mesh width $H = \varepsilon h$, $h \leq \varepsilon$. The dimension of the FE space in this asymptotic regime is $N = O(H^{-n}) = O(\varepsilon^{-n}h^{-n})$. Using standard error estimates, i.e., $\|e_N^\varepsilon\|_{H^1(\Omega)} \leq CH^p \|u^\varepsilon\|_{H^{p+1}(\Omega)}$, and the *a-priori* estimate (6) for u^ε we obtain

$$\|e_N^\varepsilon\|_{H^1(\Omega)} \leq C \left(\frac{H}{\varepsilon} \right)^p = Ch^p \leq C\varepsilon^{-p} N^{-\frac{p}{n}} \leq CN^{-\frac{p}{2n}} \quad \text{for } N \succeq \varepsilon^{-2n}.$$

We see that we obtain a robust convergence rate of $O(N^{-\frac{p}{2n}})$, as compared to the (non-robust) rate of $O(N^{-\frac{p}{n}})$ of standard FEM. The robustness of the two-scale FEM was achieved by an increase in dimension and the use of tensor product approximations in $\Omega \times \widehat{Q}$.

With two-scale FEM resolution, ε independent convergence is achieved by inflating the dimension of the approximation: we resolve the fine scales by simultaneously approximating in $(x, y) \in \Omega \times \widehat{Q} \subset \mathbb{R}^n$. The tensor product two-scale FE space represents full interactions between scales. The product structure of $\Omega \times \widehat{Q}$ and the anisotropic regularity in Theorem 1.1 allow, however, to obtain the convergence in Theorem 2.2 with substantially fewer degrees of freedom: the scale interaction is ‘thinned out’ by means of sparse tensor products.

2.3 Sparse Two-Scale Interpolation

Let us consider $\Omega^\kappa \subset \mathbb{R}^n$, $\kappa = 1, 2$ two Lipschitz bounded domains, keeping in mind that for us they will be $\Omega^1 = \Omega$ and $\Omega^2 = \widehat{Q}$. We give first a rather abstract construction of a sparse grid in $\Omega^1 \times \Omega^2$ and it is in this most general setting that we prefer to work out the error estimates. Assume that $\{\mathbf{S}_L^\kappa\}_{L=0}^\infty$, $\kappa = 1, 2$, are two dense hierarchic sequences of finite dimensional subspaces of $H^1(\Omega^\kappa)$

$$\mathbf{S}_0^\kappa \subset \mathbf{S}_1^\kappa \subset \dots \subset \mathbf{S}_L^\kappa \subset H^1(\Omega^\kappa). \quad (12)$$

At level L , N_L^κ will denote the dimension of \mathbf{S}_L^κ . In order to employ this hierarchy in the context of FE methods the following *approximation property* is needed:

$$\min_{v \in \mathbf{S}_L^\kappa} \|u - v\|_{H^1(\Omega^\kappa)} \leq \Psi^\kappa(N_L^\kappa, s) \|u\|_{H^{s+1}(\Omega^\kappa)}, \quad \forall u \in H^{s+1}(\Omega^\kappa), \quad (13)$$

where $\Psi^\kappa(N, s) \rightarrow 0$ for $s > 0$ as $N \rightarrow \infty$. For regular solutions the usual FE-spaces based on quasiuniform, shape regular meshes are suitable.

Let $\{\mathcal{T}_{H_L}\}_{L \in \mathbb{N}}$ be a nested sequence of regular affine triangulations of a Lipschitz domain Ω of meshwidth $H_L = H_{L-1}/2$, $\forall L \geq 1$ and let $p \geq 1$ be a polynomial degree. Then

$$\mathbf{S}_L := S^p(\Omega, \mathcal{T}_{H_L}) := \{u \in C^0(\bar{\Omega}) : u|_K \in S^p(K) \quad \forall K \in \mathcal{T}_{H_L}\} \quad (14)$$

satisfies (12), while the approximation property reads

$$\min_{v \in S^p(\Omega, \mathcal{T}_{H_L})} \|u - v\|_{H^1(\Omega)} \leq \Psi(N_L, s) \|u\|_{H^{s+1}(\Omega)}, \quad (15)$$

where $\Psi(N, s) = O(N^{-\min(p,s)/n})$. Estimates similar to (15) also hold for p -version or spectral element methods, i.e. on fixed \mathcal{T} as $p = p_L \rightarrow \infty$.

Within this abstract setting we follow [10] and introduce the so called *hierarchical excess* of the scale (12), by

$$W_L^\kappa := \mathbf{S}_L^\kappa \ominus \mathbf{S}_{L-1}^\kappa \quad L \geq 0. \quad (16)$$

We also set $\mathbf{S}_{-1}^\kappa := \{0\}$ and we note that complements are uniquely defined by some given Hilbert structure which produces the usual topology in $H^1(\Omega^\kappa)$. Further, P_L^κ will be the orthogonal projection on \mathbf{S}_L^κ with respect to this Hilbert structure. We remark that (13) still holds for $L = -1$, by choosing $\Psi^\kappa(N_{-1}^\kappa, s)$ to be the embedding constant of $H^1(\Omega^\kappa)$ in $H^{s+1}(\Omega^\kappa)$. With respect to the Hilbert structure mentioned above, \mathbf{S}_L^κ is an orthogonal sum of the form $\mathbf{S}_L^\kappa = \bigoplus_{0 \leq i \leq L} W_i^\kappa$. Using this decomposition, we define the *full tensor product FE-space* in $\Omega^1 \times \Omega^2$ at level $L \in \mathbb{N}$ by

$$\mathbf{S}_{L,L} := \mathbf{S}_L^1 \otimes \mathbf{S}_L^2 = \bigoplus_{0 \leq i,j \leq L} (W_i^1 \otimes W_j^2) \subset H^{1,1}(\Omega^1 \times \Omega^2), \quad (17)$$

and the *sparse tensor product FE-space* at level L by

$$\widehat{\mathbf{S}}_{L,L} := \bigoplus_{0 \leq i+j \leq L} (W_i^1 \otimes W_j^2) \subset H^{1,1}(\Omega^1 \times \Omega^2). \quad (18)$$

This correlation shows the main difference between standard (full) tensor product spaces and the sparse interpolation spaces: if for the full tensor product space the resolution is defined in each direction separately, the *overall* resolution for the sparse tensor product spaces is limited by L ; the sparse two-scale spaces result therefore from skipping certain subspaces (degrees of freedom).

For a given $U \in H^1(\Omega^1 \times \Omega^2)$ we define the *sparse interpolant* of U in $\widehat{\mathbf{S}}_{L,L}$, as given by

$$\widehat{U}^L := \sum_{0 \leq i+j \leq L} (P_i^1 - P_{i-1}^1) \otimes (P_j^2 - P_{j-1}^2) U. \quad (19)$$

With these notations, the following estimate will enable us to deduce from (13) an approximation property of the sparse scale $(\widehat{\mathbf{S}}_{L,L})_{L \in \mathbb{N}}$, too.

Proposition 2.3 *Assume that the sequence (12) of FE-spaces $\{\mathbf{S}_L^\kappa\}_L$ has the approximation property (13) and $s, t > 0$. Then there exists $C > 0$ such that for all $U \in H^{s+1, t+1}(\Omega^1 \times \Omega^2)$ it holds:*

$$\begin{aligned} \min_{V \in \widehat{\mathbf{S}}_{L,L}} \|U - V\|_{H^{1,1}(\Omega^1 \times \Omega^2)} &\leq \|U - \widehat{U}^L\|_{H^{1,1}(\Omega^1 \times \Omega^2)} \\ &\leq C \left\{ \left[\sum_{i=0}^{L+1} \Psi^1(N_{i-1}^1, s) \Psi^2(N_{L-i}^2, t) \right] \|U\|_{H^{s+1, t+1}(\Omega^1 \times \Omega^2)} \right. \\ &\quad \left. + \left[\sum_{i=L+1}^{\infty} \Psi^1(N_i^1, s) \right] \|U\|_{H^{s+1, 1}(\Omega^1 \times \Omega^2)} \right\}. \end{aligned} \quad (20)$$

Concerning the approximation quality, the accuracy of the sparse grid interpolant is only slightly deteriorated with a logarithmic factor but the number of dof is reduced to that of the discretization of one subdomain only (here again up to some logarithmic factor):

Proposition 2.4 *Assume that $U \in H^{s+1,s+1}(\Omega^1 \times \Omega^2)$. Then, with the hierarchic sequences $\mathbf{S}_L^1 = S^p(\Omega^1, \mathcal{T}_{H_L})$, $\mathbf{S}_L^2 = S^\mu(\Omega^2, \mathcal{T}_{h_L})$, $H_L \cong h_L$, $p \cong \mu$, the sparse interpolant converges, as $L \rightarrow \infty$, with the rate*

$$\min_{V \in \widehat{\mathbf{S}}_{L,L}} \|U - V\|_{H^{1,1}(\Omega^1 \times \Omega^2)} \leq C(\log \widehat{N}_L)^{1+\delta} \widehat{N}_L^{-\delta} \|U\|_{H^{s+1,s+1}(\Omega^1 \times \Omega^2)}, \quad (21)$$

where $\delta = \min(p, s)/n$.

Proof. Taking into account that $N_j = O(2^{nj})$ and using (15) in (20), with $s = t$, we obtain

$$\min_{V \in \widehat{\mathbf{S}}_{L,L}} \|U - V\|_{H^{1,1}(\Omega^1 \times \Omega^2)} \leq C \cdot \log N_L \cdot N_L^{-\delta} \|U\|_{H^{s+1,s+1}(\Omega^1 \times \Omega^2)}. \quad (22)$$

In (22), $N_L = \dim(\mathbf{S}_L) = O(2^{nL})$. Also, due to (18), $\widehat{N}_L := \dim(\widehat{\mathbf{S}}_{L,L}) = O(L \cdot 2^{nL})$. Therefore, rephrasing (22) in terms of the number of degrees of freedom \widehat{N}_L we obtain (21). \square

We return now to the two-scale homogenization problem and recall that for sufficiently smooth data the solution u^ε can be viewed as $u^\varepsilon(x) = U^\varepsilon(x, x/\varepsilon)$, where $U^\varepsilon(x, y)$ depends smoothly on ε .

We have then constructed two-scale FE spaces as traces of full tensor product spaces in $\Omega \times \widehat{Q}$ and we have showed that the FE method based on these spaces leads to robust rates of convergence at the price of discretizing both in Ω and \widehat{Q} , i.e. in \mathbb{R}^{2n} . The anisotropic regularity of U^ε and the tensor product structure of the two-scale FE-spaces allow to apply the ideas of sparse interpolation.

Assume that we have the following two sequences of hierarchic FE spaces $\{\mathbf{S}_i^1\}_{i=0}^L \subset H^1(\Omega)$ and $\{\mathbf{S}_j^2\}_{j=0}^L \subset H_{\text{per}}^1(\widehat{Q})$. As before, we have in mind sequences of FE spaces of the form $\mathbf{S}_i^1 = S^p(\Omega, \mathcal{T}_{H_i})$ and $\mathbf{S}_j^2 = S_{\text{per}}^\mu(\widehat{Q}, \widehat{\mathcal{T}}_{h_j})$ where $\{\mathcal{T}_{H_i}\}$ and $\{\widehat{\mathcal{T}}_{h_j}\}$ are sequences of affine nested FE meshes in Ω and \widehat{Q} , respectively, and $\{\widehat{\mathcal{T}}_{h_j}\}$ are periodic. We assume further that $p \cong \mu$ and $H_i \cong h_i$ for all $i = 0, \dots, L$. We denote by $\widehat{\mathbf{S}}_{L,L}$ the sparse FE space associated to these sequences.

For the sparse two-scale FEM convergence result we need a slightly modified approximation property in both Ω and \widehat{Q} . More precisely, the approximation property reads: for all $\alpha \in \mathbb{N}_0^n$ with $0 \leq \alpha_j \leq 1$, for all $s \geq n-1$, $t \geq 0$ and for $i, j = 0, \dots, L$ it holds:

$$\begin{aligned} \left(\sum_{K \in \mathcal{T}_{H_i}} \|D^\alpha(u - P_i^1 u)\|_{L^2(K)}^2 \right)^{1/2} &\leq \psi_\alpha^1(N_i^1, s) \|u\|_{H^{s+1}(\Omega)} \\ \|v - P_j^2 v\|_{H_{\text{per}}^1(\widehat{Q})} &\leq \psi^2(N_j^2, t) \|v\|_{H^{t+1}(\widehat{Q})} \\ \|\nabla(v - P_j^2 v)\|_{L_{\text{per}}^2(\widehat{Q})} &\leq \psi^2(N_j^2, t) \|\nabla v\|_{H^t(\widehat{Q})} \end{aligned} \quad (23)$$

with $\Psi_\alpha^1(N, s), \Psi^2(N, t) \rightarrow 0$ as $N \rightarrow \infty$. Using then the same arguments as for the proof of Proposition 2.1, Theorem 2.2 and Proposition 2.3 we obtain the following result on convergence of the sparse two-scale FEM:

Proposition 2.5 Assume for the solution u^ε of (3) the two-scale regularity

$$\|U^\varepsilon\|_{H^{s+1}(\Omega, H_{\text{per}}^{s+1}(\hat{Q}))} + \|\varepsilon^{-1}\nabla_y U^\varepsilon\|_{H^{s+1}(\Omega, H_{\text{per}}^s(\hat{Q}))} \leq C \|f\|_{H^{2s+1}(\Omega)}$$

for some $s \geq n-1$, where $C = C(s, n) > 0$ is a constant independent of ε . Let \hat{V}_N^ε be the sparse two-scale FE space $\hat{V}_N^\varepsilon := \mathcal{R}^\varepsilon \hat{\mathbf{S}}_{L,L}$, where \mathcal{R}^ε is the trace operator given by $(\mathcal{R}^\varepsilon U)(x) = U(x, x/\varepsilon)$. Denote by \hat{u}_{FE}^ε the FE solution with respect to \hat{V}_N^ε . Then, it holds

$$\|u^\varepsilon - \hat{u}_{FE}^\varepsilon\|_{H^1(\Omega)} \leq C(\log \hat{N}_L)^{1+\delta} \hat{N}_L^{-\delta} \|f\|_{H^{2s+1}(\Omega)}, \quad (24)$$

where $\hat{N}_L = \dim \hat{V}_N^\varepsilon$, $\delta = \min(p, s)/n$ and $C > 0$ is a constant independent of \hat{N}_L , ε and f .

Proof. Let us denote by $\hat{u}^{\varepsilon,L}$ the sparse two-scale interpolant of u^ε in the sparse two-scale FE space \hat{V}_N^ε given by $\hat{u}^{\varepsilon,L} = \mathcal{R}^\varepsilon \hat{U}^{\varepsilon,L}$, where $\hat{U}^{\varepsilon,L}$ is the sparse interpolant of U^ε in $\hat{\mathbf{S}}_{L,L}$ as given by (19). We split the sparse two-scale FE-error in elemental contributions on the macro mesh $\mathcal{T}_H = \mathcal{T}_{H_L}$ and use the trace Lemma 3.1 in [6] to obtain:

$$\begin{aligned} \|u^\varepsilon - \hat{u}^{\varepsilon,L}\|_{H^1(\Omega)}^2 &\leq C \sum_{K \in \mathcal{T}_{H_L}} \left\{ \sum_{\substack{0 \leq \alpha_j \leq 1 \\ |\alpha| \geq 1}} \varepsilon^{2(|\alpha|-1)} \|D_x^\alpha (U^\varepsilon - \hat{U}^{\varepsilon,L})\|_{L^2(K \times \hat{Q})}^2 \right. \\ &\quad \left. + \sum_{0 \leq \alpha_j \leq 1} \varepsilon^{2|\alpha|} \|D_x^\alpha (\varepsilon^{-1} \nabla_y) (U^\varepsilon - \hat{U}^{\varepsilon,L})\|_{L^2(K \times \hat{Q})}^2 \right\}. \end{aligned} \quad (25)$$

Due to the approximation property (23) we have that

$$\begin{aligned} &\sum_{K \in \mathcal{T}_{H_L}} \|D_x^\alpha (U^\varepsilon - \hat{U}^{\varepsilon,L})\|_{L^2(K \times \hat{Q})}^2 \\ &\leq C \left\{ \sum_{i=0}^L \left[\sum_{K \in \mathcal{T}_{H_i}} \|D_x^\alpha (P_i^1 - P_{i-1}^1) \otimes (\text{Id}^2 - P_{L-i}^2) U^\varepsilon\|_{L^2(K \times \hat{Q})}^2 \right]^{1/2} \right. \\ &\quad \left. + \sum_{i=L+1}^\infty \left[\sum_{K \in \mathcal{T}_{H_i}} \|D_x^\alpha (P_i^1 - P_{i-1}^1) \otimes \text{Id}^2 U^\varepsilon\|_{L^2(K \times \hat{Q})}^2 \right]^{1/2} \right\}^2 \\ &\leq C \left\{ \sum_{i=0}^{L+1} \Psi_\alpha^1(N_{i-1}^1, s) \Psi^2(N_{L-i}^2, t) \|U^\varepsilon\|_{H^{s+1}(\Omega, H^{t+1}(\hat{Q}))} \right. \\ &\quad \left. + \sum_{i=L+1}^\infty \Psi_\alpha^1(N_i^1, s) \|U^\varepsilon\|_{H^{s+1}(\Omega, H_{\text{per}}^1(\hat{Q}))} \right\}^2 \end{aligned}$$

and a similar estimate can be derived for $\sum_{K \in \mathcal{T}_{H_L}} \|D_x^\alpha (\varepsilon^{-1} \nabla_y) (U^\varepsilon - \hat{U}^{\varepsilon,L})\|_{L^2(K \times \hat{Q})}^2$. For the spaces considered here there holds

$$\Psi_\alpha^1(N, s) = H^{-|\alpha|} O(N^{-\min(p, s+1)/n}) \quad \text{and} \quad \Psi^2(N, t) = O(N^{-\min(\mu, t)/n}).$$

Taking $s = t \geq n-1$, $\mu \cong p$, $H_i \cong h_i$ and using that $\varepsilon/H_L \leq 1$ we conclude the proof. \square




Figure 1: Energy error for the full version of the two-scale FEM




Figure 2: Two-scale FEM and sparse two-scale FEM with $p = \mu = 1$: relative error in the energy versus the mesh size $H = h \rightarrow 0$ (left) and versus $\#$ dof (right), respectively

2.4 Numerical Results

We illustrate our error estimates for the two-scale FEM for problem (1), (2) in the one dimensional case $\Omega = (0, 1)$ with data $f(x) = e^x$, $A(y) = 2 + \cos(2\pi y)$ and $a_0(y) = 0$. The shift Theorem 1.1 applies on Ω and the solution does not exhibit boundary layers, since $u^\varepsilon(x) = U^\varepsilon(x, x/\varepsilon)$, with $U^\varepsilon(x, y)$ smooth on $\Omega \times \hat{Q}$ and 1-periodic in y .

In Figure 1 we plot for different $p = \mu \in \{1, 2, 3, 4\}$ the energy error versus $H = h$. Computations were performed for two different ε -scales, 10^{-2} and 10^{-4} , respectively. We see that the rate of convergence of $\|u^\varepsilon - u_{FE}^\varepsilon\|_{H^1(\Omega)}^2$ is proportional to H^{2p} as expected from the error estimates in Theorem 2.2. Moreover, we observe robustness of the convergence rates with respect to the parameter ε .

In Figure 2 we compare the performances of the two-scale FEM and its sparse version for $\varepsilon = 10^{-3}$. The left plot shows the relative error in the energy versus $H = h \rightarrow 0$ for both methods in the case $p = \mu = 1$. We see that the error curves are practically at the top of each other. In the right figure we plot the same relative error in the energy versus $\#$ dof for the two-scale FEM and for the sparse two-scale FEM. We clearly see the $O(N^{-1})$ rate of convergence for the two-scale FEM and the $O(N^{-2}(\log N)^4)$ -convergence rate for the sparse two-scale FEM, as predicted by Proposition 2.5.

Acknowledgement. This research was supported under the project “Homogenization and multiple scales” HMS2000 of the EC (HPRN-CT-1999-00109) and by the Swiss National Science Foundation under Project “Hierarchic FE-Models for periodic lattice and honeycomb materials” with Number BBW 21-58754.99.

References

- [1] I. Babuška, A.K. Aziz, “Survey lectures on the mathematical foundation of the finite element method”, in *The Mathematical Foundations of the Finite Element Method with Applications to Partial Differential Equations*, A. K. Aziz, ed., Academic Press, New York (1973) 5–359.
- [2] A. Bensoussan, J.L. Lions, G. Papanicolau, “Asymptotic Analysis for Periodic Structures”, North Holland, Amsterdam, 1978.
- [3] D. Cioranescu, P. Donato, “An Introduction to Homogenization”, Oxford University Press 1999.
- [4] A.M. Matache, “Spectral- and p -Finite Elements for problems with microstructure”, Ph.D. Thesis, ETH Zürich 2000.
- [5] A.M. Matache, I. Babuška, C. Schwab, “Generalized p -FEM in Homogenization”, *Numerische Mathematik*, **86** Issue 2 (2000), 319–375.
- [6] A.M. Matache, C. Schwab, “Two-Scale FEM for Homogenization Problems” (submitted), Report 2001-06, Seminar for Applied Mathematics, ETH Zürich
<http://www.sam.math.ethz.ch/reports/details/2001/2001-06.shtml>.
- [7] O.A. Oleinik, A.S. Shamaev, G.A. Yosifian, “Mathematical Problems in Elasticity and Homogenization”, North-Holland, 1992.
- [8] C. Schwab, A.M. Matache, “High order generalized FEM for lattice materials”, *Proceedings of the 3rd European Conference on Numerical Mathematics and Advanced Applications*, Finland, 1999, ed. by P. Neittaanmäki, T. Tiihonen and P. Tarvainen, World Scientific, Singapore, 2000.
- [9] B. Szabó, I. Babuška, “Finite Element Analysis”, John Wiley & Sons, Inc. 1991.
- [10] C. Zenger, “Sparse Grids”, in “Parallel Algorithms for PDE’s ”, *Proceedings of the 6th GAMM-Seminar*, Kiel, 1990.

Research Reports

No.	Authors	Title
01-09	A.-M. Matache	Sparse Two-Scale FEM for Homogenization Problems
01-08	C. Lasser, A. Toselli	Convergence of some two-level overlapping domain decomposition preconditioners with smoothed aggregation coarse space
01-07	T. von Petersdorff, C. Schwab	Wavelet-discretizations of parabolic integro-differential equations
01-06	A.-M. Matache, C. Schwab	Two-Scale FEM for Homogenization Problems
01-05	A. Buffa, R. Hiptmair, T. von Petersdorff, C. Schwab	Boundary Element Methods for Maxwell Equations in Lipschitz Domains
01-04	R. Hiptmair, C. Schwab	Natural BEM for the Electric Field Integral Equation on polyhedra
01-03	C. Schwab, A.-M. Matache	Generalized FEM for Homogenization Problems
01-02	A. Toselli, C. Schwab	Mixed <i>hp</i> -finite element approximations on geometric edge and boundary layer meshes in three dimensions
01-01	A. Buffa, M. Costabel, C. Schwab	Boundary element methods for Maxwell's equations on non-smooth domains
00-18	B.Q. Guo, C. Schwab	Analytic regularity of Stokes flow in polygonal domains
00-17	M.H. Gutknecht, S. Röllin	Variations of Zhang's Lanczos-Type Product Method
00-16	M.H. Gutknecht, S. Röllin	The Chebyshev iteration revisited
00-15	G. Schmidlin, C. Schwab	Wavelet Galerkin BEM on unstructured meshes by aggregation
00-14	B. Cockburn, G. Kanschat, D. Schötzau, C. Schwab	Local Discontinuous Galerkin methods for the Stokes system
00-13	O. Brüker, M.J. Grote, C. Mayer, A. Reusken	Robust Parallel Smoothing for Multigrid Via Sparse Approximate Inverses
00-12	C. Lasser, A. Toselli	An overlapping domain decomposition preconditioner for a class of discontinuous Galerkin approximations of advection-diffusion problems
00-11	J. Liesen, M. Rozložník, Z. Strakoš	On Convergence and Implementation of Minimal Residual Krylov Subspace Methods for Unsymmetric Linear Systems