

# Sparse approximation of singularity functions

P.-A. Nitsche

Research Report No. 2002-18  
October 2002

Seminar für Angewandte Mathematik  
Eidgenössische Technische Hochschule  
CH-8092 Zürich  
Switzerland

# Sparse approximation of singularity functions

P.-A. Nitsche

Seminar für Angewandte Mathematik  
Eidgenössische Technische Hochschule  
CH-8092 Zürich  
Switzerland

Research Report No. 2002-18

October 2002

## Abstract

We are concerned with the sparse approximation of functions of type

$$u(x) = \operatorname{Re} (|x|^\lambda (\log |x|)^\mu \Phi(\varphi)) \eta(x) + g(x)$$

on the  $d$ -dimensional unit cube  $[0, 1]^d$  with parameters  $\lambda \in \mathbb{C}$ ,  $\mu \in \mathbb{N}_0$ , smooth  $g$ , a smooth cut-off function  $\eta$ , and a function  $\Phi(\varphi)$  of the remaining coordinates but  $|x|$ , possibly singular as well (e. g. containing edge singularities). These functions arise e. g. from corners of domains in solutions to elliptic PDEs. Usually, they deteriorate the rate of convergence of numerical algorithms to approximate these solutions.

We show, that functions of this type – for a range of  $\lambda$  covering elliptic singularities – can be approximated with respect to the  $H^1$  norm by sparse grid wavelet spaces  $V_L$ ,  $\dim(V_L) = N_L$ , of biorthogonal spline wavelets of degree  $p$  essentially at the rate  $p$ :

$$\|u - P_L u\|_{H^1([0,1]^d)} \leq C N_L^{-p} (\log_2 N_L)^s \|u\|, \quad s = s(p, d),$$

where  $\|\cdot\|$  is a weighted Sobolev norm and  $P_L u \in V_L$ .

# 1 Introduction

The efficient numerical approximation of functions of several variables is required in numerous applications: in the numerical solution of partial differential equations in dimension 3 or higher, in data mining, in numerical integration over high dimensional domains, to name but a few.

Tensor products of piecewise polynomial functions of degree  $p$  on a uniform mesh of width  $h > 0$  require  $\mathcal{O}(h^{-d})$  degrees of freedom and achieve an asymptotic convergence rate in the  $H^1$  norm of  $\mathcal{O}(h^p)$  if the  $(p + 1)$ -st derivatives of the function to be approximated are square-integrable.

The growth as  $h \rightarrow 0$  of the number of degrees of freedom in tensor product interpolants is excessive if  $d$  is large. So-called *sparse interpolants* – which require only  $\mathcal{O}(h^{-1} |\log h|^{d-1})$  degrees of freedom rather than  $\mathcal{O}(h^{-d})$  – have been proposed and successfully used in applications (see, e. g., [BG], [GOS], [Z]). There, it was shown that despite their reduced number of degrees of freedom, sparse tensor product interpolants can achieve an  $H^1$  convergence rate of  $\mathcal{O}(h^p |\log h|^{d-1})$ , i.e. up to logarithmic terms the rate of the full tensor product approximation is preserved. This is possible, however, only under very strong assumptions on the regularity of the function to be approximated: certain derivatives up to order  $d(p + 1)$  must be square-integrable in order to realize the convergence rate  $\mathcal{O}(h^p |\log h|^{d-1})$  in  $H^1$ .

In particular, in the numerical treatment of partial differential equations the solutions are *not* that regular, for example due to corner singularities in elliptic problems. The potential gain in using sparse tensor product approximation is strongly limited in this case – the reduction in the convergence rate essentially offsets the reduction in degrees of freedom in the interpolant.

Singularities in solutions of elliptic problems contain as essential parts powers of distance functions to the singular support of the domain (corner, edges, etc.). For instance for polygonal domains  $\Omega$  in 2D, there holds a decomposition of the weak solution  $u$  to a uniformly elliptic boundary value problem into

$$u = u_{\text{reg}} + u_{\text{sing}},$$

with smooth  $u_{\text{reg}}$  and a finite sum

$$(1) \quad u_{\text{sing}} = \sum^{<\infty} \text{Re} \left( r_i^{\lambda_i} (\log r_i)^{\mu_i} \Phi_i(\varphi_i) \eta_i(x) \right),$$

(in local polar coordinates  $(r_i, \varphi_i)$  associated to the corners (here not indexed)) with smooth cut-off functions  $\eta_i$ , smooth functions  $\Phi_i$  of the angle coordinates  $\varphi_i$ , and parameters  $\lambda_i \in \mathbb{C}$ ,  $\mu_i \in \mathbb{N}_0$ , see e. g. [BDLN 1-3].

For Dirichlet-Dirichlet corners, one knows  $\text{Re } \lambda_i > 1/2$ , and for Dirichlet-Neumann corners,  $\text{Re } \lambda_i > 1/4$ .

In higher dimensions, decomposition theorems are not that easy to write down, but estimates on the growth (of the function and its derivatives) towards the singular support of the domain are known, see e. g. [MR].

The rate of approximation of singular functions determines the convergence rate of Finite Element Methods for elliptic problems. The dimension  $N$  of FE spaces of continuous, piecewise polynomials of degree  $p$  on a quasiuniform mesh of meshwidth  $h$  is  $N = \mathcal{O}(h^{-d})$ . For solutions  $u \in H^{s+1}(\Omega)$ , the  $H^1$  convergence rate of the best approximation is  $\mathcal{O}(N^{-\min\{p,s\}/d})$ .

In dimension  $d = 2$ , singular terms of the above type belong to  $H^{s+1}(\Omega)$  with  $s < \text{Re}(\lambda)$ .

To overcome the reduced convergence rate, the nonuniform distribution of degrees of freedom by local mesh refinement has been proposed. This can be done *a-priori*, i. e. by grading the mesh towards the singular support of (1), or *a-posteriori*, by adaptive mesh refinement and nonlinear approximation.

In dimension 2, these techniques allow to recover the maximal asymptotic rate  $\mathcal{O}(N^{-p/2})$ , see, e. g., [BDD], [MNS] for recent results and, in particular, [CDD], where it has been proven, that the rate  $\mathcal{O}(N^{-b})$  is achievable by adaptive (non linear) wavelet approximation whenever such a rate is possible by so called best  $N$ -term approximation (for related numerical results see also [BBCCDDU]). The result in [CDD] is in particular not restricted to functions of the special type (1), but applies to all functions which belong to a scale of suitable Besov spaces (see [V] for details).

The result [CDD] applies to certain FEM bases with ‘isotropic’ support, or with ‘shape regular’ elements in FE-terminology. In polyhedra  $\Omega$  in dimension  $d = 3$ , the appearance of edge singularities with *anisotropic* regularity does not allow to achieve the rate  $\mathcal{O}(N^{-p/d})$  with local mesh refinements, as long as the basis is isotropic. However, this rate can be achieved with suitable *anisotropic* mesh refinements, (see e. g. [A]).

In the present paper, we therefore analyze the approximation of singular functions built from powers of distance functions to corners, edges, etc., in anisotropically refined *sparse* tensor product spaces.

We prove that anisotropic sparse tensor product spaces (of piecewise polynomials of degree  $p$ ) allow to approximate corner and edge singularities in 3d (defined on  $[0, 1]^d$ ) at a rate

$$(2) \quad \mathcal{O}(N^{-p} (\log_2 N)^s), \quad s = 2p + 3/2$$

in the  $H^1$  norm, where  $N$  is the number of degrees of freedom in the sparse, anisotropic tensor product. In particular, (2) shows that the reduction in the convergence rate due to the higher dimension and low Sobolev regularity can be eliminated (up to logarithmic terms).

Our proof is not restricted to dimension 3, but applies to any dimension  $d > 1$ . In particular, and at first sight somewhat surprisingly, even in dimension 2, the use of sparse tensor products of wavelet bases with local refinement for the approximation of ‘isotropic’ corner singularities (1) gives an improvement over the ‘optimal’ rate  $\mathcal{O}(N^{-p/2})$  in the  $H^1$  norm: we obtain that for each polynomial degree  $p \geq 1$ , there is a sequence of sparse spline wavelet approximants of degree  $p$  which converges at the rate  $\mathcal{O}(N^{-p} (\log_2 N)^{p+1})$ ,

as opposed to the rate  $\mathcal{O}(N^{-p/2})$  obtained from nonlinear, adaptive approximation in the 'isotropic' setting (i. e. support of wavelet bases and Besov spaces used to measure the regularity of (1) are isotropic).

Let us point out that our result yields immediately upper bounds on nonlinear and adaptive approximation schemes in anisotropic wavelet bases for the elliptic singularities, which are superior to those of [CDD] and [BDD] in the isotropic setting. These connections will be elaborated elsewhere.

The remainder of this paper is organized as follows: In section 2, we introduce the wavelets we use and prove the one-dimensional results; the key result is lemma 3. In section 3, we consider the two-dimensional case and give a detailed proof of the two-dimensional consistency estimate using an abstract tensor product argument. In section 4, we extend the two-dimensional results to  $d$  dimensions by iterating the arguments of section 3. The main theorems are theorems 4 and 5.

Concerning the application to singular functions arising from corners of polygonal domains or from changes in the type of boundary conditions, theorem 3 (for the 2d case) and theorem 6 (for the 3d case) specialize the results to these cases.

In the sequel,  $\sim$  in expressions like  $A \sim B$  means that both quantities can be uniformly bounded by constant multiples of each other. Likewise  $\lesssim$  indicates inequalities up to constant factors.

*Acknowledgement:* The author would like thank Prof. Christoph Schwab for inducting him into this subject and for a lot of helpful discussions.

## 2 Wavelet approximation with local refinement in one dimension

In this chapter we develop the one-dimensional means in order to approximate singular functions that are powers of distance functions to points. Instead of using explicit representations like (1), we derive an estimate in terms of weighted Sobolev norms. For this, we define a hierarchy of finite dimensional spaces  $V_L^\beta \subset V_{L+1}^\beta \subset H^1$  (where  $\beta$  is a parameter indicating the local refinement towards the singularity) and corresponding projectors  $P_L^\beta$ , and prove that

$$\dim(V_L^\beta) \leq C(p) 2^L \quad \text{and} \quad \|u - P_L^\beta u\|_{H^1([0,1])} \leq C(p) L^{1/2} 2^{-pL} \|u\|_*,$$

for functions  $u$  with  $\|u\|_* < \infty$  ( $p$  is the local polynomial degree of the ansatz functions). The norm  $\|\cdot\|_*$  is a weighted Sobolev norm (and will be defined precisely in the sequel). The singularity functions of interest are in these weighted spaces.

The one-dimensional framework will later be used to infer  $d$ -dimensional results by sparse 'tensor products' of the univariate results. In this regard, the use of hierarchic spaces is essential to our approach.

We begin by describing the wavelets we use. Then we define the locally (towards the singularity) refined projectors and ansatz spaces, as well as the weighted spaces. The last part of this section is devoted to the proof of the approximation property in terms of these weighted norms.

Let  $\psi, \tilde{\psi}$  be compactly supported biorthogonal spline wavelets of (a fixed) degree  $p \geq 1$  on the interval  $[0, 1]$ , such that for

$$u = \sum_{j=0}^{\infty} \sum_{k=1}^{k_j} c_{jk} \psi_{jk} = \sum_{j=0}^{\infty} \sum_{k=1}^{k_j} \langle u, \tilde{\psi}_{jk} \rangle_{L^2} \psi_{jk}$$

the following norm equivalences resp. estimates hold:

$$(3) \quad \begin{aligned} c_1 \|u\|_{H^t([0,1])}^2 &\leq \sum_{j,k} 2^{2tj} c_{jk}^2 \leq c_2 \|u\|_{H^t([0,1])}^2, & \text{for } 0 \leq t < p + 1/2, \\ \sum_{\substack{j,k \\ j \leq L}} 2^{2tj} c_{jk}^2 &\leq c_2 \|u\|_{H^t([0,1])}^2, & \text{for } p + 1/2 \leq t < p, \\ \sum_{\substack{j,k \\ j \leq L}} 2^{2tj} c_{jk}^2 &\leq c_2 L \|u\|_{H^t([0,1])}^2, & \text{for } t = p, \end{aligned}$$

where the constants  $c_1$  and  $c_2$  depend on  $t$ . Such wavelets are constructed, for instance, in [DS]; see also [DKU]. For the one-sided norm estimates, see [PSS].

On each level  $j$ , the number of wavelets  $\psi_{jk}$  is bounded by  $p(2^j + N)$  with some global constant  $N \in \mathbb{N}$ . To simplify notation, we take  $N = 0$ . We ignore the dependence of the indices  $k$  on  $p$ , thus regarding an index  $(j, k)$  standing for  $p$  local wavelets basis functions. The support of a wavelet  $\psi_{jk}$  is of order  $2^{-j}$ . Note, that the norm equivalences are a consequence of vanishing moment properties of the wavelets respectively its duals.

We define now one-dimensional wavelet projectors  $P_L^\beta$ , which are designed to approximate singular functions of type  $x \mapsto x^\lambda$ . For this, let  $I = \{(j, k) \in \mathbb{N} \times \mathbb{N} : j \in \mathbb{N}, 1 \leq k \leq 2^j\}$ , and define for  $L \in \mathbb{N}$  and fixed  $\beta \in [0, 1)$

$$(4) \quad \Lambda_L^\beta \equiv \{(j, k) \in I : (k2^{-j})^\beta \leq 2^{L-j}\}.$$

The index set  $\Lambda_L^\beta$  contains all index pairs  $(j, k)$  with  $j \leq L$ , but considerably more towards the left endpoint 0. But still it holds

$$(5) \quad \#\Lambda_L^\beta \lesssim p 2^L,$$


with a constant depending only on  $\beta$ . In the vicinity of 0, the minimal support of wavelets indexed by  $\Lambda_L^\beta$  is of order  $2^{-L/(1-\beta)}$ .

**Definition 1** (projector  $P_L^\beta$ ). Let  $u$  be a function with wavelet decomposition  $u = \sum_{j,k} c_{jk} \psi_{jk}$ . Then the projector  $P_L^\beta$  is defined by

$$(6) \quad P_L^\beta u \equiv \sum_{(j,k) \in \Lambda_L^\beta} c_{jk} \psi_{jk}.$$

We denote by  $V_L^\beta$  the image of  $P_L^\beta$ , that is  $V_L^\beta = \text{span}\{\psi_{jk} : (j,k) \in \Lambda_L^\beta\} \subset H^p([0,1])$ .

The following figure depicts a typical space  $V_L^\beta$ : the basis functions spanning  $V_L^\beta$  are marked by black dots. In this example,  $L = 2$  and  $\beta = 0.6$ ; the maximal occurring level in the vicinity of the origin is  $L/(1-\beta) = 5$ .



The projectors  $P_L^\beta$  are simultaneously stable with respect to the  $H^t$  norms,  $0 \leq t < p+1/2$ :

**Lemma 1** (stability of  $P_L^\beta$ ). Let  $t \in [0, p+1/2)$ . Then  $P_L^\beta : H^t([0,1]) \rightarrow H^t([0,1])$  is bounded independently of  $L$ .

*Proof.* Using (3), we have

$$\begin{aligned} \|P_L^\beta u\|_{H^t([0,1])}^2 &= \left\| \sum_{(j,k) \in \Lambda_L^\beta} c_{jk} \psi_{jk} \right\|_{H^t([0,1])}^2 \sim \sum_{(j,k) \in \Lambda_L^\beta} 2^{2tj} c_{jk}^2 \\ &\leq \sum_{j,k} 2^{2tj} c_{jk}^2 \sim \|u\|_{H^t([0,1])}^2. \end{aligned}$$

□

Next, we will prove an approximation result for  $P_L^\beta$  involving weighted Sobolev norms. For this, we first introduce these weighted spaces:

**Definition 2** (weighted spaces,  $d=1$ ). For  $m \in \mathbb{N}$  let  $H_\gamma^m([0, 1])$  be the space of all measurable functions  $u$  such that

$$\|u\|_{H_\gamma^m([0,1])} := \left( \sum_{i=0}^m \|(D^i u)x^\gamma\|_{L^2([0,1])}^2 \right)^{1/2}$$

is finite. The expression  $\|\cdot\|_{H_\gamma^m([0,1])}$  is a norm on this space.

We have the following embedding result:

**Lemma 2** (embedding,  $d=1$ ). If  $\gamma > p + 1/2$ , then the following embedding is continuous:

$$H_\gamma^{p+1}([0, 1]) \hookrightarrow L^q([0, 1]) \quad \text{for } q < \frac{1}{\gamma - (p + 1/2)}.$$

*Proof.* We first note, that there exists a continuous extension operator  $E : H_\gamma^{p+1}([0, 1]) \rightarrow H_\gamma^{p+1}([0, 2])$  with the property  $\text{supp } Eu \subset [0, 2]$ . This is due to the fact that close to the right boundary  $\{1\}$  the space  $H_\gamma^{p+1}$  coincides (in a local sense) with the space  $H^{p+1}$ .

Let now  $\gamma > p + 1/2$ . We write

$$\begin{aligned} |u(x)| &= |Eu(x)| \leq \int_x^2 |(Eu)'(s)| ds \leq \int_x^2 \int_{s_1}^2 \cdots \int_{s_p}^2 |D^{p+1}(Eu)(t)| dt ds_p \cdots ds_1 \\ &= \int_x^2 \int_{s_1}^2 \cdots \int_{s_p}^2 |D^{p+1}(Eu)(t) t^\gamma| t^{-\gamma} dt ds_p \cdots ds_1 \\ &\leq \int_x^2 \int_{s_1}^2 \cdots \int_{s_{p-1}}^2 \underbrace{\left( \int_{s_p}^2 |D^{p+1}(Eu)(t) t^\gamma|^2 dt \right)^{1/2}}_{\leq \|D^{p+1}(Eu) t^\gamma\|_{L^2([0,2])}} \left( \int_{s_p}^2 t^{-2\gamma} dt \right)^{1/2} ds_p \cdots ds_1 \\ &\leq c \|D^{p+1}u t^\gamma\|_{L^2([0,1])} \\ &\lesssim \|u\|_{H_\gamma^{p+1}([0,1])} \cdot \int_x^2 \int_{s_1}^2 \cdots \int_{s_{p-1}}^2 \left( \int_{s_p}^2 t^{-2\gamma} dt \right)^{1/2} ds_p \cdots ds_1 \\ &\lesssim \|u\|_{H_\gamma^{p+1}([0,1])} \cdot \int_x^2 \int_{s_1}^2 \cdots \int_{s_{p-1}}^2 s_p^{1/2-\gamma} ds_p \cdots ds_1 \\ &\lesssim \|u\|_{H_\gamma^{p+1}([0,1])} \cdot \int_x^2 \int_{s_1}^2 \cdots \int_{s_{p-2}}^2 s_{p-1}^{3/2-\gamma} ds_{p-1} \cdots ds_1 \\ &\lesssim \cdots \lesssim x^{p+1/2-\gamma} \|u\|_{H_\gamma^{p+1}([0,1])}. \end{aligned}$$



Taking this to the power  $q$ , integrating over  $[0, 1]$  and taking to the power  $1/q$  yields

$$\|u\|_{L^q([0,1])} \lesssim \|u\|_{H_\gamma^{p+1}([0,1])} \underbrace{\left( \int_0^1 x^{q(p+1/2-\gamma)} dx \right)^{1/q}}_{=: C}.$$

$C$  is finite, if  $q < 1/(\gamma - (p + 1/2))$ .  $\square$

Now we are able to prove the one-dimensional approximation result with respect to a weighted norm of the above type:

**Lemma 3** (weighted norm consistency of  $P_L^\beta$ ,  $d=1$ ). *For*

$$1 > \beta > \gamma > \frac{p + 1/2}{p + 1}$$

and  $u \in H_{\gamma(p+1)}^{p+1}([0, 1])$  it holds

$$(7) \quad \|u - P_L^\beta u\|_{L^2([0,1])} \lesssim L^{1/2} 2^{-(p+1)L} \|u\|_{H_{\gamma(p+1)}^{p+1}([0,1])}.$$

For

$$\beta > \gamma > \frac{p - 1/2}{p}$$

and  $u \in H_{\gamma p}^{p+1}([0, 1])$  it holds

$$(8) \quad \|u - P_L^\beta u\|_{H^1([0,1])} \lesssim L^{1/2} 2^{-pL} \|u\|_{H_{\gamma p}^{p+1}([0,1])}.$$

*Proof.* It suffices to prove the statements above for  $u \in H^{p+1}([0, 1])$ . The lemma then follows from a density argument.

We split the interval  $[0, 1]$  into subintervals of length  $2^{-L/(1-\beta)}$ , for simplicity assuming  $L/(1-\beta)$  to be an integer (otherwise rounded off). For  $s = 1, \dots, 2^{L/(1-\beta)}$ , set

$$I(s) = [(s-1)2^{-L/(1-\beta)}, s2^{-L/(1-\beta)}].$$

Let  $j_0(s)$  be a bound for the minimal local level of wavelets  $\psi_{jk}$  with  $(j, k) \notin \Lambda_L^\beta$  and  $\text{supp } \psi_{jk} \cap I(s) \neq \emptyset$ . Examining the condition  $(s2^{-L/(1-\beta)})^\beta \leq 2^{L-j}$ , we see that  $j_0(s)$  can be chosen to be

$$j_0(s) = \frac{L}{1-\beta} - \beta \log_2 s.$$

Applying (3) and (3), we estimate

$$\begin{aligned} \|u - P_L^\beta u\|_{L^2(I(s))}^2 &\lesssim \sum_{\substack{(j,k) \notin \Lambda_L^\beta \\ \text{supp } \psi_{jk} \cap I(s) \neq \emptyset}} c_{jk}^2 = \sum_{j \geq j_0(s)} 2^{-2(p+1)j} \sum_{\substack{k: (j,k) \notin \Lambda_L^\beta \\ \text{supp } \psi_{jk} \cap I(s) \neq \emptyset}} 2^{2(p+1)j} c_{jk}^2 \\ &\lesssim \sum_{j \geq j_0(s)} j 2^{-2(p+1)j} \|u\|_{H^{p+1}(I(s))}^2 \lesssim j_0(s) 2^{-2(p+1)j_0(s)} \|u\|_{H^{p+1}(I(s))}^2. \end{aligned}$$

Using  $2^{-2(p+1)j_0(s)} = 2^{-2(p+1)L}(s2^{-L/(1-\beta)})^{2(p+1)\beta}$  and  $j_0(s) \lesssim L$ , we infer

$$\|u - P_L^\beta u\|_{L^2(I(s))}^2 \lesssim L 2^{-2(p+1)L} \|u(s2^{-L/(1-\beta)})^{(p+1)\beta}\|_{H^{p+1}(I(s))}^2.$$

Away from 0, i. e. in case of  $s > 1$ , we have  $s2^{-L/(1-\beta)} \sim x$  for  $x \in I(s)$ . This yields

$$(9) \quad \|u - P_L^\beta u\|_{L^2([2^{-L/(1-\beta)}, 1])}^2 \lesssim L 2^{-2(p+1)L} \|ux^{(p+1)\beta}\|_{H^{p+1}([2^{-L/(1-\beta)}, 1])}^2.$$

In case of  $s = 1$ , we employ  $L^2$  stability of the projectors  $P_L^\beta$  (lemma 1) and the embedding result, lemma 2.

If  $\gamma > (p + 1/2)/(p + 1)$ , i. e.  $\gamma(p + 1) > p + 1/2$ , we estimate

$$\|u - P_L^\beta u\|_{L^2(I(1))}^2 \lesssim \|u\|_{L^2(I(1))}^2 \leq |I(1)|^{1-2/q} \|u\|_{L^q(I(1))}^2 \lesssim |I(1)|^{1-2/q} \|u\|_{H_{\gamma(p+1)}^{p+1}([0,1])}^2.$$

Here,  $q < 1/(\gamma(p + 1) - (p + 1/2))$  corresponding to lemma 2. To guarantee  $|I(1)|^{1-2/q} \leq 2^{-2(p+1)L}$ , we get the constraint  $q \geq 1/(\beta(p + 1) - (p + 1/2))$ . Hence, if  $\beta > \gamma$ , we can choose such a  $q$ . Together with (9), the  $L^2$  estimate follows.

The  $H^1$  estimate is done in an analogous fashion: Using (3) and (3), we get

$$\begin{aligned} \|u - P_L^\beta u\|_{H^1(I(s))}^2 &\lesssim \sum_{\substack{(j,k) \notin \Lambda_L^\beta \\ \text{supp } \psi_{jk} \cap I(s) \neq \emptyset}} 2^{2j} c_{jk}^2 = \sum_{j \geq j_0(s)} 2^{-2pj} \sum_{\substack{k: (j,k) \notin \Lambda_L^\beta \\ \text{supp } \psi_{jk} \cap I(s) \neq \emptyset}} 2^{2(p+1)j} c_{jk}^2 \\ &\lesssim \sum_{j \geq j_0(s)} j 2^{-2pj} \|u\|_{H^{p+1}(I(s))}^2 \lesssim j_0(s) 2^{-2pj_0(s)} \|u\|_{H^{p+1}(I(s))}^2. \end{aligned}$$

Using  $2^{-2pj_0(s)} = 2^{-2pL}(s2^{-L/(1-\beta)})^{2p\beta}$  and  $j_0(s) \lesssim L$ , we infer

$$\|u - P_L^\beta u\|_{H^1(I(s))}^2 \lesssim L 2^{-2pL} \|u(s2^{-L/(1-\beta)})^{p\beta}\|_{H^{p+1}(I(s))}^2.$$

Away from 0, i. e. in case of  $s > 1$ , we have  $s2^{-L/(1-\beta)} \sim x$  for  $x \in I(s)$ . This yields

$$(10) \quad \|u - P_L^\beta u\|_{H^1([2^{-L/(1-\beta)}, 1])}^2 \lesssim L 2^{-2pL} \|ux^{p\beta}\|_{H^{p+1}([2^{-L/(1-\beta)}, 1])}^2.$$

In case  $s = 1$ , we estimate, using lemma 2,

$$\|D(u - P_L^\beta u)\|_{L^2(I(1))}^2 \lesssim \|Du\|_{L^2(I(1))}^2 \leq |I(1)|^{1-2/q} \|Du\|_{L^q(I(1))}^2 \lesssim |I(1)|^{1-2/q} \|u\|_{H_{\gamma p}^{p+1}(I(1))}^2,$$

if  $q < 1/(\gamma p - (p - 1/2))$  and  $\gamma p > p - 1/2$ . To guarantee  $|I(1)|^{1-2/q} \leq 2^{-2pL}$ , we get the constraint  $q \geq 1/(\beta p - (p - 1/2))$ . Hence, if  $\beta > \gamma$ , we can choose such a  $q$ . Together with (10), the  $H^1$  estimate (8) follows.  $\square$

**Remark 1.** The function  $u : x \mapsto x^\lambda$  belongs to  $H_{\gamma p}^{p+1}([0, 1])$ , if

$$\gamma > 1 - \frac{2\lambda - 1}{2p}.$$

Thus, if  $\lambda > 1/2$ , there is such a  $\gamma < 1$ .

**Remark 2.** In [BN] a similar one-dimensional result for Chui prewavelet bases of degree 2 and 3 was proven, using a slightly different refinement towards the origin and a different method of proof.

A similar approach was already undertaken in [PS].

### 3 Wavelet approximation with local refinement in two dimensions

We come now to the two-dimensional case. Consider the unit square  $[0, 1]^2 \subset \mathbb{R}^2$ . We define a wavelet basis by taking tensor products of the univariate wavelets introduced above, i. e. the set  $\{\psi_{jk} \otimes \psi_{j'k'} : (j, k) \in I, (j', k') \in I\}$ . Note the anisotropy of the supports of these wavelets.

We will employ a sparse tensor product rather than a full tensor product of the projectors  $P_L^\beta$ . The sparse tensor product projector has comparable approximation properties to the full one, but uses a remarkably lower-dimensional subspace to project onto. We will give the details in the following.

Let  $\beta \in [0, 1)$  be fixed and  $L \in \mathbb{N}$ . Then we define the index sets

$$(11) \quad \hat{\Lambda}_L^\beta \equiv \bigcup_{i+j \leq L} \Lambda_i^\beta \times \Lambda_j^\beta$$

and the trial spaces

$$(12) \quad \hat{V}_L^\beta := \text{span}\{\psi_{jk} \otimes \psi_{j'k'} : (j, k, j', k') \in \hat{\Lambda}_L^\beta\} = \bigoplus_{i+j \leq L} V_i^\beta \otimes V_j^\beta.$$

From (5), we see

$$(13) \quad \#\hat{\Lambda}_L^\beta = \dim(\hat{V}_L^\beta) \lesssim p^2 L 2^L.$$

**Definition 3** (projector  $\hat{P}_L^\beta$ ). *Let  $u$  be a function with wavelet decomposition*

$$u = \sum_{j,k,j',k'} c_{jkj'k'} \psi_{jk} \otimes \psi_{j'k'}.$$

Then the projector  $\hat{P}_L^\beta$  is defined by

$$(14) \quad \hat{P}_L^\beta u \equiv \sum_{j,k,j',k' \in \hat{\Lambda}_L^\beta} c_{jkj'k'} \psi_{jk} \otimes \psi_{j'k'}.$$

**Definition 4.** Let  $(s, t) \in \mathbb{N}^2$ . Then we denote by

$$H^{s,t}([0, 1]^2)$$

the space of all measurable functions  $u : [0, 1]^2 \rightarrow \mathbb{R}$ , such that the norm

$$\|u\|_{H^{s,t}([0,1]^2)} := \left( \sum_{\substack{0 \leq i \leq s \\ 0 \leq j \leq t}} \|D_1^i D_2^j u\|_{L^2([0,1]^2)}^2 \right)^{1/2}$$

is finite. That is,  $H^{s,t}([0, 1]^2) = H^s([0, 1]) \otimes H^t([0, 1])$ .

**Lemma 4.** The projectors  $\hat{P}_L^\beta$  are stable in  $H^1 \otimes L^2$  independently of  $L$ .

*Proof.* From the one-dimensional norm equivalences, one readily derives norm equivalences for tensor product spaces in the following form:

$$\left\| \sum_{j,k,j',k'} c_{jkj'k'} \psi_{jk} \otimes \psi_{j'k'} \right\|_{H^{s,t}([0,1]^2)}^2 \sim \sum_{j,k,j',k'} 2^{2sj+2tj'} c_{jkj'k'}^2, \quad 0 \leq s, t < p + 1/2.$$

For  $s = 1$  and  $t = 0$ , one accordingly has for  $u = \sum c_{jkj'k'} \psi_{jk} \otimes \psi_{j'k'}$

$$\begin{aligned} \|\hat{P}_L^\beta u\|_{H^{1,0}([0,1]^2)}^2 &= \left\| \sum_{(j,k,j',k') \in \hat{\Lambda}_L^\beta} c_{jkj'k'} \psi_{jk} \otimes \psi_{j'k'} \right\|_{H^{1,0}([0,1]^2)}^2 \sim \sum_{(j,k,j',k') \in \hat{\Lambda}_L^\beta} 2^{2j} c_{jkj'k'}^2 \\ &\leq \sum_{j,k,j',k'} 2^{2j} c_{jkj'k'}^2 \sim \|u\|_{H^{1,0}([0,1]^2)}^2. \end{aligned}$$

□

**Definition 5.** For  $\sigma, \tau \geq 0$  and  $p, q \in \mathbb{N}$  let

$$H_{\sigma,\tau}^{p,q}([0, 1]^2) := H_\sigma^p([0, 1]) \otimes H_\tau^q([0, 1]).$$

We will derive a consistency estimate for  $\hat{P}_L^\beta$  by an abstract tensor product argument following the lines of [ST]. Since  $\|\varphi\|_{H^1([0,1]^2)}^2 \leq \|\varphi\|_{H^{1,0}([0,1]^2)}^2 + \|\varphi\|_{H^{0,1}([0,1]^2)}^2$ , it suffices (by symmetry) to estimate the  $H^{1,0}([0, 1]^2)$  norm of the error  $u - \hat{P}_L^\beta u$ . To simplify notation, we set

$$\|\cdot\| := \|\cdot\|_{H^1([0,1])} \otimes \|\cdot\|_{L^2([0,1])} = \|\cdot\|_{H^{1,0}([0,1]^2)}.$$

Let  $Q_{1,i}^\beta$  be the orthogonal projection onto  $V_i^\beta$  with respect to the  $H^1([0,1])$  norm, and let  $Q_{2,i}^\beta$  be the orthogonal projection onto  $V_i^\beta$  with respect to the  $L^2([0,1])$  norm.

We set

$$\hat{Q}_L^\beta := \sum_{i+j \leq L} (Q_{1,i}^\beta - Q_{1,i-1}^\beta) \otimes (Q_{2,j}^\beta - Q_{2,j-1}^\beta);$$

$\hat{Q}_L^\beta$  is the orthogonal projection onto  $\hat{V}_L^\beta$  with respect to  $\|\cdot\| = \|\cdot\|_{H^1,0([0,1]^2)}$ . Note, that in general  $\hat{Q}_L^\beta \neq \hat{P}_L^\beta$  (but, of course,  $\text{image } \hat{Q}_L^\beta = \text{image } \hat{P}_L^\beta = \hat{V}_L^\beta$ ).

By stability of  $\hat{P}_L^\beta$  in  $H^1 \otimes L^2$ , lemma 4, we have

$$\begin{aligned} \|u - \hat{P}_L^\beta u\| &= \|u - \hat{Q}_L^\beta u + \hat{Q}_L^\beta u - \hat{P}_L^\beta u\| = \|u - \hat{Q}_L^\beta u + \hat{P}_L^\beta \hat{Q}_L^\beta u - \hat{P}_L^\beta u\| \\ &= \|(\text{id} - \hat{P}_L^\beta)(u - \hat{Q}_L^\beta u)\| \leq c \|u - \hat{Q}_L^\beta u\|. \end{aligned}$$

It therefore suffices to estimate the error  $\|u - \hat{Q}_L^\beta u\|$  to get an approximation result for  $\hat{P}_L^\beta$ .

By orthogonality of the projectors  $\hat{Q}_i^\beta$ , we have quite generally

$$\begin{aligned} \|u - \hat{Q}_L^\beta u\|^2 &= \|u - \sum_{i+j \leq L} (Q_{1,i}^\beta - Q_{1,i-1}^\beta) \otimes (Q_{2,j}^\beta - Q_{2,j-1}^\beta) u\|^2 \\ &= \left\| \sum_{i+j \geq L+1} (Q_{1,i}^\beta - Q_{1,i-1}^\beta) \otimes (Q_{2,j}^\beta - Q_{2,j-1}^\beta) u \right\|^2 \\ &= \left\| \sum_{i=0}^{L+1} (Q_{1,i}^\beta - Q_{1,i-1}^\beta) \otimes (\text{id} - \hat{Q}_{2,L-i}^\beta) u \right\|^2 + \left\| \sum_{i=L+2}^{\infty} (Q_{1,i}^\beta - Q_{1,i-1}^\beta) \otimes \text{id} u \right\|^2 \\ &= \sum_{i=0}^{L+1} \|(Q_{1,i}^\beta - Q_{1,i-1}^\beta) \otimes (\text{id} - \hat{Q}_{2,L-i}^\beta) u\|^2 + \|(\text{id} - Q_{1,L+1}^\beta) \otimes \text{id} u\|^2. \end{aligned}$$

Applying once more orthogonality, we estimate the last line by

$$\leq \sum_{i=0}^{L+1} \|(\text{id} - Q_{1,i-1}^\beta) \otimes (\text{id} - \hat{Q}_{2,L-i}^\beta) u\|^2 + \|(\text{id} - Q_{1,L+1}^\beta) \otimes \text{id} u\|^2.$$

By stability of the orthogonal projections  $Q_{r,i}^\beta$ , we can translate this estimate back into an estimate involving the projectors  $P_L^\beta$ . To see this (using simplified notation), let  $Q, P, A$  be projectors with  $Q$  stable. Then it holds

$$\begin{aligned} \|(\text{id} - Q) \otimes A u\| &= \|(\text{id} - P) \otimes A u + P \otimes A u - Q \otimes A u\| \\ &= \|(\text{id} - P) \otimes A u + QP \otimes A u - Q \otimes A u\| \\ &= \|(\text{id} - P) \otimes A u + (Q \otimes \text{id})(P \otimes A u - \text{id} \otimes A) u\| \\ &= \|(\text{id} \otimes \text{id} - Q \otimes \text{id})(\text{id} - P) \otimes A u\| \leq c \|(\text{id} - P) \otimes A u\|. \end{aligned}$$

Hence, we finally get

$$\|u - \hat{P}_L^\beta u\|^2 \leq c \left( \sum_{i=0}^{L+1} \|(\text{id} - P_{i-1}^\beta) \otimes (\text{id} - P_{L-i}^\beta) u\|^2 + \|(\text{id} - P_{L+1}^\beta) \otimes \text{id} u\|^2 \right).$$

Now using the one-dimensional consistency result, lemma 3, we infer

$$\begin{aligned} \|u - \hat{P}_L^\beta u\|^2 &\lesssim \sum_{i=0}^{L+1} L^2 2^{-2p(i-1)} 2^{-2(p+1)(L-i)} \|u\|_{H_{\gamma p, \gamma(p+1)}^{p+1, p+1}([0,1]^2)}^2 \\ &\quad + L 2^{-2p(L+1)} \|u\|_{H_{\gamma p}^{p+1}([0,1]) \otimes L^2([0,1])}^2, \end{aligned}$$

hence,

$$\|u - \hat{P}_L^\beta u\|^2 \lesssim L^2 2^{-2pL} \|u\|_{H_{\gamma p, \gamma(p+1)}^{p+1, p+1}([0,1]^2)}^2.$$

We summarize:

**Theorem 1.** For

$$1 > \beta > \gamma > \frac{p+1/2}{p+1}$$

and  $u \in H_{\gamma p, \gamma(p+1)}^{p+1, p+1}([0,1]^2) \cap H_{\gamma(p+1), \gamma p}^{p+1, p+1}([0,1]^2)$ , it holds

$$\|u - \hat{P}_L^\beta u\|_{H^1([0,1]^2)} \lesssim L 2^{-pL} \left( \|u\|_{H_{\gamma p, \gamma(p+1)}^{p+1, p+1}([0,1]^2)} + \|u\|_{H_{\gamma(p+1), \gamma p}^{p+1, p+1}([0,1]^2)} \right).$$

**Remark 3.** Note, that trivially holds

$$H^{2(p+1)}([0,1]^2) \subset H_{\gamma p, \gamma(p+1)}^{p+1, p+1}([0,1]^2) \cap H_{\gamma(p+1), \gamma p}^{p+1, p+1}([0,1]^2) \quad \text{for } \gamma \geq 0.$$

In an analogous fashion, one proves

**Theorem 2.** For

$$1 > \beta > \gamma > \frac{p+1/2}{p+1}$$

and  $u \in H_{\gamma(p+1), \gamma(p+1)}^{p+1, p+1}([0,1]^2)$ , it holds

$$\|u - \hat{P}_L^\beta u\|_{L^2([0,1]^2)} \lesssim L^{3/2} 2^{-(p+1)L} \|u\|_{H_{\gamma(p+1), \gamma(p+1)}^{p+1, p+1}([0,1]^2)}.$$

**Remark 4.** In [O1], P. Oswald has considered best  $N$  term approximation from sparse grid spaces built on the univariate Haar system ( $p = 0$ ). He derives approximation rates in Sobolev spaces  $H^s([0,1]^2)$ ,  $-1 < s < 1/2$ . In case of approximation in  $L^2([0,1]^2)$ , i. e.  $s = 0$ , he achieves an approximation rate of  $N^{-1} (\log N)^{3/2}$  for singularity functions of type  $(1; \alpha, \alpha)$ ,  $\alpha < 1/2$ . Here, a function  $f(x_1, x_2)$  is called to be of type  $(m; \alpha, \alpha)$ , if  $|\partial_1^k \partial_2^l f| \leq C x_1^{-(\alpha+k)} x_2^{-(\alpha+l)}$  for  $0 \leq k, l \leq m$ .

A singular function of type  $(1; \alpha, \alpha)$  belongs to  $H_{\gamma, \gamma}^{1,1}([0,1]^2)$  if  $\gamma > 1/2 + \alpha$ . Hence, our result is consistent with Oswald's.

## Application to elliptic singularities in 2D

Consider an elliptic boundary value problem on a polygonal domain  $\Omega \subset \mathbb{R}^2$ :

$$Lu = f \quad \text{in } \Omega, \quad Bu = g \quad \text{on } \partial\Omega,$$

where  $L$  is a smooth, uniformly elliptic, linear second-order differential operator in divergence form,  $B$  is a smooth boundary operator of Dirichlet or Neumann type, and  $f$  and  $g$  are smooth functions.

The weak solution  $u$  is smooth (in the Sobolev scale) in the interior of  $\Omega$ , but exhibits corner singularities, which reduce global regularity dramatically. In general,  $u \notin H^2(\Omega)$ . It is well known, cf. [BDLN 1-3], that there holds a decomposition

$$u = u_{\text{reg}} + u_{\text{sing}},$$

with (arbitrarily) smooth  $u_{\text{reg}}$ , which we specialize to  $u_{\text{reg}} \in H^{2(p+1)}(\Omega)$ , and (depending on the chosen smoothness of  $u_{\text{reg}}$ ) a finite sum

$$u_{\text{sing}} = \sum^{<\infty} \text{Re} \left( r_i^{\lambda_i} (\log r_i)^{\mu_i} \Phi_i(\varphi_i) \eta_i(x) \right),$$

(in local polar coordinates  $(r_i, \varphi_i)$  associated to the corners) with smooth cut-off functions  $\eta_i$ , smooth functions  $\Phi_i$  of the angle coordinates  $\varphi_i$ , and parameters  $\lambda_i \in \mathbb{C}$ ,  $\mu_i \in \mathbb{N}_0$ . For Dirichlet-Dirichlet corners, one knows  $\text{Re } \lambda_i > 1/2$ , and for Dirichlet-Neumann corners,  $\text{Re } \lambda_i > 1/4$ .

We prove now, that this type of singularities can be approximated at a rate  $p$  in the  $H^1$  norm. Note that we can allow  $\text{Re } \lambda > 0$ :

**Theorem 3.** *Let*

$$u(x) = \text{Re} \left( |x|^\lambda (\log |x|)^\mu \Phi(\varphi) \eta(x) \right) + g(x) \in H^1([0, 1]^2)$$

*with smooth  $\eta$  and  $\Phi$ , parameters  $\lambda \in \mathbb{C}$ ,  $\text{Re}(\lambda) > 0$ ,  $\mu \in \mathbb{N}_0$ , and a function  $g \in H^{2(p+1)}([0, 1]^2)$ . Then there are  $\beta, \gamma$  (depending on  $\lambda$ ) with*

$$1 > \beta > \gamma > \frac{p + 1/2}{p + 1},$$

*such that*

$$u \in H_{\gamma p, \gamma(p+1)}^{p+1, p+1}([0, 1]^2) \cap H_{\gamma(p+1), \gamma p}^{p+1, p+1}([0, 1]^2).$$

*That means, there is a constant  $C$  depending via  $\beta$  and  $\gamma$  only on  $\lambda$ , such that*

$$\|u - \hat{P}_L^\beta u\|_{H^1([0, 1]^2)} \leq C L 2^{-pL} \left( \|u\|_{H_{\gamma p, \gamma(p+1)}^{p+1, p+1}([0, 1]^2)} + \|u\|_{H_{\gamma(p+1), \gamma p}^{p+1, p+1}([0, 1]^2)} \right).$$

Concerning approximability in  $L^2$ , we have:

If  $\operatorname{Re}(\lambda) > -1$  and the remaining assumptions as above are met, then there are  $\beta, \gamma$  with

$$1 > \beta > \gamma > \frac{p+1/2}{p+1},$$

such that

$$u \in H_{\gamma(p+1), \gamma(p+1)}^{p+1, p+1}([0, 1]^2).$$

That means, there is a constant  $C$  such that

$$\|u - \hat{P}_L^\beta u\|_{L^2([0,1]^2)} \leq C L^{3/2} 2^{-(p+1)L} \|u\|_{H_{\gamma(p+1), \gamma(p+1)}^{p+1, p+1}([0,1]^2)}^2.$$

**Remark 5.** Note, that we are only concerned with the approximation of certain singular terms, which are defined on the unit square. In this paper, we do not address the question of handling complex geometries with sparse grids within a numerical scheme.

*Proof.* The norm of  $H_{\gamma p, \gamma(p+1)}^{p+1, p+1}([0, 1]^2)$  is given by

$$\|u\|_{H_{\gamma p, \gamma(p+1)}^{p+1, p+1}([0,1]^2)} = \left( \sum_{0 \leq i, j \leq p+1} \int_{[0,1]^2} |(D_1^i D_2^j u) x_1^{\gamma p} x_2^{\gamma(p+1)}|^2 dx \right)^{1/2}.$$

Estimating  $x_1^{\gamma p} x_2^{\gamma(p+1)}$  by  $|x|^{\gamma(2p+1)}$ , we see that  $\operatorname{Re}(|x|^\lambda (\log|x|)^\mu) \in H_{\gamma p, \gamma(p+1)}^{p+1, p+1}([0, 1]^2)$  if

$$\gamma > 1 - \frac{\lambda}{2p+1}.$$

The same holds for the space  $H_{\gamma(p+1), \gamma p}^{p+1, p+1}([0, 1]^2)$ .

The smooth factors  $\eta$  and  $\Phi$  leave the regularity class unchanged, and  $g$  is trivially in the space  $H_{\gamma p, \gamma(p+1)}^{p+1, p+1}([0, 1]^2) \cap H_{\gamma(p+1), \gamma p}^{p+1, p+1}([0, 1]^2)$ , cf. remark 3.

Hence, if  $\lambda > 0$ , we can find  $1 > \beta > \gamma > 1 - \lambda/(2p+1)$  such that the assumptions of theorem 1 are met.

The proof of the  $L^2$  estimate is done analogously. □



## 4 Generalization to $d$ dimensions

The  $d$ -dimensional case with  $d > 2$  follows from the two-dimensional one by iteration. We consider the domain  $[0, 1]^d$  and the tensor product basis

$$\{\psi_{j_1 k_1 \dots j_d k_d} = \psi_{j_1 k_1} \otimes \dots \otimes \psi_{j_d k_d} : (j_i, k_i) \in I, i = 1, \dots, d\}.$$

Let  $\beta \in [0, 1)$  be fixed and  $L \in \mathbb{N}$ . Then we define the index sets

$$\hat{\Lambda}_{L,d}^\beta \equiv \bigcup_{j_1 + \dots + j_d \leq L} \Lambda_{j_1}^\beta \times \dots \times \Lambda_{j_d}^\beta$$

and the trial spaces

$$\hat{V}_{L,d}^\beta := \text{span}\{\psi_{j_1 k_1 \dots j_d k_d} : (j_1, k_1, \dots, j_d, k_d) \in \hat{\Lambda}_{L,d}^\beta\} = \bigoplus_{j_1 + \dots + j_d \leq L} V_{j_1}^\beta \otimes \dots \otimes V_{j_d}^\beta.$$

From (5), we see

$$\#\hat{\Lambda}_{L,d}^\beta = \dim(\hat{V}_{L,d}^\beta) \lesssim p^d L^{d-1} 2^L.$$

The constant depends here and in the sequel additionally on the dimension  $d$ .

**Definition 6** (projector  $\hat{P}_{L,d}^\beta$ ). *Let  $u$  be a function with wavelet decomposition*

$$u = \sum_{j_1, k_1, \dots, j_d, k_d} c_{j_1 k_1 \dots j_d k_d} \psi_{j_1 k_1} \otimes \dots \otimes \psi_{j_d k_d}.$$

*Then the projector  $\hat{P}_{L,d}^\beta$  is defined by*

$$\hat{P}_{L,d}^\beta u \equiv \sum_{j_1, k_1, \dots, j_d, k_d \in \hat{\Lambda}_{L,d}^\beta} c_{j_1 k_1 \dots j_d k_d} \psi_{j_1 k_1} \otimes \dots \otimes \psi_{j_d k_d}.$$

**Definition 7.** *Let  $(s_1, \dots, s_d) \in \mathbb{N}^d$ . Then we denote by*

$$H^{s_1, \dots, s_d}([0, 1]^d)$$

*the space of all measurable functions  $u : [0, 1]^d \rightarrow \mathbb{R}$ , such that the norm*

$$\|u\|_{H^{s_1, \dots, s_d}([0, 1]^d)} := \left( \sum_{0 \leq j_i \leq s_i} \|D_1^{j_1} \dots D_d^{j_d} u\|_{L^2([0, 1]^d)}^2 \right)^{1/2}$$

*is finite. That is,  $H^{s_1, \dots, s_d}([0, 1]^d) = \bigotimes_{i=1, \dots, d} H^{s_i}([0, 1])$ .*

**Definition 8.** *For  $\sigma_1, \dots, \sigma_d \geq 0$  and  $p_1, \dots, p_d \in \mathbb{N}$  let*

$$H_{\sigma_1, \dots, \sigma_d}^{p_1, \dots, p_d}([0, 1]^d) := \bigotimes_{i=1, \dots, d} H_{\sigma_i}^{p_i}([0, 1]).$$

We further introduce the abbreviation

$$H(\gamma, p, d) := \bigcap_{k=1, \dots, d} \left( \bigotimes_{i=1, \dots, d} H_{\gamma(p+1-\delta_{ki})}^{p+1}([0, 1]) \right)$$

and take as a norm the summed up norms.

**Remark 6.** Note, that trivially holds

$$H^{d(p+1)}([0, 1]^d) \subset H(\gamma, p, d) \quad \text{for } \gamma \geq 0.$$

With the latter notation, theorem 1 may be rephrased as follows:

$$\beta > \gamma > \frac{p+1/2}{p+1}, u \in H(\gamma, p, d) \quad \Rightarrow \quad \|u - \hat{P}_{L,2}^\beta u\|_{H^1([0,1]^d)} \lesssim L 2^{-pL} \|u\|_{H(\gamma,p,d)}.$$

Iterating the argument from section 3, we infer the following two theorems:

**Theorem 4** ( $H^1$  consistency of  $\hat{P}_L^\beta$ ). For

$$1 > \beta > \gamma > \frac{p+1/2}{p+1}$$

and  $u \in H(\gamma, p, d)$ , it holds

$$\|u - \hat{P}_{L,d}^\beta u\|_{H^1([0,1]^d)} \lesssim L^{d/2} 2^{-pL} \|u\|_{H(\gamma,p,d)}.$$

Letting  $N_L := \dim \hat{V}_L^\beta \lesssim p^d L^{d-1} 2^L$  be the number of degrees of freedom, this means

$$\|u - \hat{P}_{L,d}^\beta u\|_{H^1([0,1]^d)} \lesssim N_L^{-p} (\log_2 N_L)^s \|u\|_{H(\gamma,p,d)},$$

$$s = (d-1)p + d/2.$$

**Theorem 5** ( $L^2$  consistency of  $\hat{P}_L^\beta$ ). For

$$1 > \beta > \gamma > \frac{p+1/2}{p+1}$$

and  $u \in H_{\gamma(p+1), \dots, \gamma(p+1)}^{p+1, \dots, p+1}([0, 1]^d)$ , it holds

$$\|u - \hat{P}_{L,d}^\beta u\|_{L^2([0,1]^d)} \lesssim L^{d-1/2} 2^{-(p+1)L} \|u\|_{H_{\gamma(p+1), \dots, \gamma(p+1)}^{p+1, \dots, p+1}([0,1]^d)}.$$

In terms of number of degrees of freedom, this means

$$\|u - \hat{P}_{L,d}^\beta u\|_{L^2([0,1]^d)} \lesssim N_L^{-(p+1)} (\log_2 N_L)^s \|u\|_{H_{\gamma(p+1), \dots, \gamma(p+1)}^{p+1, \dots, p+1}([0,1]^d)},$$

$$s = (d-1)p + 2d - 3/2.$$

**Remark 7.** There holds the continuous inclusion

$$H(\gamma, p, d) \subset H_{\gamma(p+1), \dots, \gamma(p+1)}^{p+1, \dots, p+1}([0, 1]^d).$$

**Remark 8.** By interpolating the estimates of theorem 4 and 5, consistency estimates in  $H^s([0, 1]^d)$ ,  $0 < s < 1$  are readily obtained.

**Remark 9.** The function  $x \mapsto \operatorname{Re} |x|^\lambda$  on  $[0, 1]^d$  belongs to  $H(\gamma, p, d)$ , if

$$\gamma > 1 - \frac{(d-2) + 2\lambda}{2d(p+1) - 2}.$$

So, if

$$\lambda > 1 - d/2,$$

we can find such a  $\gamma < 1$ .

More generally, let  $\rho$  be the distance to a face of  $[0, 1]^d$  of dimension  $k$  containing the origin (for  $k = 0$  the corner 0, for  $k = 1$  an edge containing 0, etc.). Then the function  $x \mapsto \operatorname{Re} \rho^\lambda$  on  $[0, 1]^d$  belongs to some  $H(\gamma, p, d)$  with  $\gamma < 1$ , if

$$\lambda > 1 - (d - k)/2.$$

Note, that this condition on the exponent  $\lambda$  coincides with the condition of membership in  $H^1([0, 1]^d)$  at all, which indicates sharpness of our result with respect to powers of distance functions to lower-dimensional faces.

### Application to elliptic singularities in 3D

We apply our approximation result to elliptic corner-edge singularities in 3D. Consider on a polyhedral domain  $\Omega$  (for instance  $[0, 1]^3$ ) the boundary value problem

$$(15) \quad Lu = f \quad \text{in } \Omega, \quad Bu = g \quad \text{on } \partial\Omega,$$

where  $L$  is a smooth, uniformly elliptic, linear second-order differential operator in divergence form,  $B$  is a smooth boundary operator of face-wise Dirichlet or Neumann type, and  $f$  and  $g$  are smooth functions.

As in the 2D case, the weak solution  $u$  is smooth away from corners and edges, but exhibits singularities towards the singularities of the domain. The solution's asymptotics towards corners and edges can be described by weighted Sobolev spaces, and we quote only one result (in simplified form), which was proven by Maz'ya and Roßmann (Theorem 2 in [MR]):

There is  $\varepsilon > 0$  (depending on the domain), such that for  $\delta \in \mathbb{R}$  and  $\ell \in \mathbb{N}$  with  $|\delta - \ell - 1| < \varepsilon$  there holds

$$f \in V_\delta^\ell \quad \Rightarrow \quad u \in V_\delta^{\ell+2}.$$

Here, the space  $V_\delta^\ell$  is defined as the set of functions  $v$ , for which

$$\left( \int_\Omega \left( \prod r_i^\delta \right) \left( \prod (\rho_j/r)^\delta \right) \sum_{|\alpha| \leq \ell} \rho^{2(|\alpha| - \ell)} |D^\alpha u|^2 dx \right)^{1/2}$$

is finite. The functions  $r_i$  are the distance functions to the vertices of  $\Omega$ , the functions  $\rho_j$  are the distance functions to the edges of  $\Omega$ , and  $r = \min_i r_i$ ,  $\rho = \min_j \rho_j$ .

For the sake of simplicity of exposition, we specialize this to the case of one 'active' cube-like corner (which is supposed to be the origin). Then the norm of  $V_\delta^\ell$  reads

$$\left( \int_\Omega r^\delta \left( \prod_{j=1}^3 (\rho_j/r)^\delta \right) \sum_{|\alpha| \leq \ell} \rho^{2(|\alpha| - \ell)} |D^\alpha u|^2 dx \right)^{1/2}.$$

Using the general relations  $\rho \leq \rho_j \leq r$ , a calculation shows

$$f \in C^\infty(\bar{\Omega}) \quad \Rightarrow \quad f \in V_\delta^\ell \quad \text{for all } \delta \geq \ell.$$

Now choose  $\ell = 3p + 1$  and  $\delta = \ell + 1 - \varepsilon_0 = 3p + 2 - \varepsilon_0$  for some  $0 < \varepsilon_0 < \min\{\varepsilon, 1\}$ . From the above shift theorem, we see that  $u \in V_\delta^{3p+3}$ . But this readily implies  $u \in H(\gamma, p, 3)$  for

$$\frac{\delta}{3p+2} = \frac{3p+2-\varepsilon_0}{3p+2} < \gamma < 1.$$

We conclude:

**Theorem 6.** *The solution  $u$  of problem (15) lies locally in some space  $H(\gamma, p, 3)$  with  $\gamma < 1$  and may therefore (as a function transferred to the unit cube) be approximated at a rate  $p$  in the  $H^1$  norm.*

## References

- [A] Th. Apel: Anisotropic finite elements: Local estimates and applications; Series "Advances in Numerical Mathematics", Teubner, Stuttgart, 1999
- [BBCCDDU] A. Barinka, T. Barsch, P. Charton, A. Cohen, S. Dahlke, W. Dahmen, K. Urban: Adaptive Wavelet Schemes for Elliptic Problems – Implementations and Numerical Experiments; SIAM J. Sci. Comput. 23(3), pp. 910-939 (2001)
- [BDD] P. Binev, W. Dahmen, R. DeVore: Adaptive Finite Element Methods with Convergence Rates; preprint, 2002.
- [BDLN1] M. Bourlard, M. Dauge, M. Lubuma, S. Nicaise: Coefficients des singularités pour des problèmes aux limites elliptiques sur un domaine à points coniques. I: Résultats généraux pour le problème de Dirichlet; RAIRO Modél. Anal. Numér. 24 (1990), pp. 27-52

- [BDLN2] M. Boulard, M. Dauge, M. Lubuma, S. Nicaise: Coefficients des singularités pour des problèmes aux limites elliptiques sur un domaine à points coniques. II: Quelques opérateurs particuliers; *RAIRO Modél. Anal. Numér.* 24 (1990), pp. 343-367
- [BDLN3] M. Boulard, M. Dauge, M. Lubuma, S. Nicaise: Coefficients of the singularities for elliptic boundary value problems on domains with conical points. III: Finite Element Methods on polygonal domains; *SIAM J. Numer. Anal.* 29 (1992), Nr. 1, pp. 136-155
- [BG] H. Bungartz, M. Griebel: A Note on the Complexity of Solving Poisson's Equation for Spaces of Bounded Mixed Derivatives; *J. Compl.* 15, pp. 167-199 (1999)
- [BN] C. Bourgeois, S. Nicaise: Approximation par préondelettes augmentée de l'équation de la plaque libre polygonale; *C. R. Acad. Sci. Paris, t. 329, Série I*, pp. 727-730 (1999)
- [CDD] A. Cohen, W. Dahmen, R. DeVore: Adaptive Wavelet Methods for Elliptic Operator Equations: Convergence Rates; *Math. Comp.*, Vol. 70, Nr. 233, pp. 27-75 (2000)
- [D] M. Dauge: Elliptic boundary value problems on corner domains: smoothness and asymptotics of solutions; Springer Verlag Berlin [etc.], 1988
- [DKU] W. Dahmen, A. Kunoth, K. Urban: Biorthogonal Spline-Wavelets on the Interval – Stability and Moment Conditions; *Appl. Comput. Harm. Anal.* 6, pp. 132-196 (1999)
- [DS] W. Dahmen, R. Schneider: Wavelets with Complementary Boundary Conditions – Function Spaces on the Cube; *Result. Math.* 34, pp. 255-293 (1998)
- [G] P. Grisvard: Elliptic Problems in Nonsmooth Domains, Pitman Publishing Boston, 1985
- [GK] M. Griebel, S. Knapik: Optimized tensor-product approximation spaces; *Constr. Approx.* 16 (4), pp. 525-540 (2000)
- [GOS] M. Griebel, P. Oswald, T. Schiekofer: Sparse grids for boundary integral equations; *Numer. Math.* 83(2), pp. 279-312 (1999)
- [GT] D. Gilbarg, N. S. Trudinger: Elliptic Partial Differential Equations of Second Order; Springer-Verlag Berlin Heidelberg New York, 1998
- [KMR] V. A. Kozlov, V. G. Mazya, J. Rossmann: Elliptic boundary value problems in domains with point singularities; Rhode Island American Mathematical Society, Providence, 1997
- [MNS] P. Morin, R. Nocketto, K. Siebert: Data oscillation and convergence of adaptive FEM; *SIAM J. Numer. Anal.* 38, Nr. 2, 2000
- [MR] V. B. Maz'ya, J. Rossmann: On the Agmon-Miranda maximum principle, *Ann. Global Anal. Geom.* 9, Nr. 3, 1991, pp. 253-303
- [O1] P. Oswald: On best n-term approximation by Haar functions in  $H^s$ -norms; in *Metric Function Theory and Related Topics in Analysis*, Nicol'skij, Kashin, Izaak (eds.), AFC Moscow, 1999, pp. 137-163
- [O2] P. Oswald: Best n-term capacitance approximation on sparse grids; *Proc. 12th Intern. Conf. on Domain Decomp. Methods in Science and Engineering*, T. Chan et al. (eds.), Chiba, 1999, pp. 437-444
- [PS] T. von Petersdorff, Ch. Schwab: Boundary Element Methods with Wavelets and Mesh Refinement; Research Report No. 95-10, Seminar für Angewandte Mathematik, ETH Zürich (1995)
- [PSS] T. von Petersdorff, Ch. Schwab, R. Schneider: Multiwavelets for second-kind integral equations, *SIAM J. Numer. Anal.*, Vol. 34, No. 6, pp. 2212-2227, 1997
- [ST] Ch. Schwab, R. A. Todor: Sparse finite elements for elliptic problems with stochastic data, SAM ETHZ Report 2002-05. (online available under: <http://www.sam.math.ethz.ch/reports/>), to appear in *Numer. Math.* (2003)
- [V] R. DeVore: Nonlinear Approximation; in *Acta Numerica* 7, Cambridge University Press, Cambridge, UK, pp. 51-150 (1998)
- [Z] Ch. Zenger: Sparse Grids; in *Parallel Algorithms for PDE*, Proc. 6th GAMM Seminar, Kiel (ed. W. Hackbusch), Vieweg, Braunschweig, pp. 241-251 (1991)

# Research Reports

No.	Authors	Title
02-18	P.-A. Nitsche	Sparse approximation of singularity functions
02-17	S.H. Christiansen	Uniformly stable preconditioned mixed boundary element method for low-frequency electromagnetic scattering
02-16	S.H. Christiansen	Mixed boundary element method for eddy current problems
02-15	A. Toselli, X. Vasseur	Neumann-Neumann and FETI preconditioners for $hp$ -approximations on geometrically refined boundary layer meshes in two dimensions
02-14	Th.P. Wihler	Locking-Free DGFEM for Elasticity Problems in Polygons
02-13	S. Beuchler, R. Schneider, C. Schwab	Multiresolution weighted norm equivalences and applications
02-12	M. Kruzik, A. Prohl	Macroscopic modeling of magnetic hysteresis
02-11	A.-M. Matache, C. Schwab, T. von Petersdorff	Fast deterministic pricing of options on Lévy driven assets
02-10	D. Schötzau, C. Schwab, A. Toselli	Mixed $hp$ -DGFEM for incompressible flows
02-09	Ph. Frauenfelder, Ch. Lage	Concepts - An object-oriented software package for partial differential equations
02-08	A.-M. Matache, J.M. Melenk	Two-Scale Regularity for Homogenization Problems with Non-Smooth Fine Scale Geometry
02-07	G. Schmidlin, C. Lage, C. Schwab	Rapid solution of first kind boundary integral equations in $\mathbb{R}^3$
02-06	M. Torrilhon	Exact Solver and Uniqueness Conditions for Riemann Problems of Ideal Magnetohydrodynamics
02-05	C. Schwab, R.-A. Todor	Sparse Finite Elements for Elliptic Problems with Stochastic Data
02-04	R. Jeltsch, K. Nipp	CSE Program at ETH Zurich: Are we doing the right thing?
02-03	L. Diening, A. Prohl, M. Ruzicka	On Time-Discretizations for Generalized Newtonian Fluids
02-02	A. Toselli	$hp$ Discontinuous Galerkin Approximation for the Stokes Problem
02-01	F.M. Buchmann, W.P. Petersen	Solving Dirichlet problems numerically using the Feynman-Kac representation