

First order k -th moment finite element
analysis of nonlinear operator equations
with stochastic data

A. Chernov* and Ch. Schwab

Research Report No. 2011-51
August 2011

Seminar für Angewandte Mathematik
Eidgenössische Technische Hochschule
CH-8092 Zürich
Switzerland

*Hausdorff Center for Mathematics & Institute for Numerical Simulation, University
of Bonn, 53115 Bonn, Germany

FIRST ORDER k -TH MOMENT FINITE ELEMENT ANALYSIS OF NONLINEAR OPERATOR EQUATIONS WITH STOCHASTIC DATA

ALEXEY CHERNOV AND CHRISTOPH SCHWAB

Dedicated to W. L. Wendland on the occasion of this 75th anniversary

August 23, 2011

ABSTRACT. We develop and analyze a class of efficient algorithms for uncertainty quantification of nonlinear operator equations. The algorithms are based on sparse Galerkin discretizations of tensorized linearizations at nominal parameters. Specifically, for a class of abstract nonlinear, parametric operator equations $J(\alpha, u) = 0$ for random parameters α with realizations in a neighborhood of a nominal parameter α_0 . Under some structural assumptions on the parameter dependence, by the implicit function theorem, $J(\alpha, u) = 0$ admits locally unique solutions $u = S(\alpha)$ for all values α in some neighborhood of α_0 . Random parameters $\alpha(\omega) = \alpha_0 + r(\omega)$, are shown to imply a unique random solution $u(\omega) = S(\alpha(\omega))$. We derive a multilinear, tensorized operator equation for the deterministic computation of k -th order statistical moments of the solution fluctuations $u(\omega) - S(\alpha_0)$, provided that statistical moments of the random parameter perturbation $r(\omega)$ are known. We present a sparse tensor Galerkin discretization for the tensorized first order perturbation equation. We prove a shift theorem for the k -point correlation equation in anisotropic smoothness scales and deduce that sparse tensor Galerkin discretizations of this equation converge in accuracy vs. complexity which equals, up to logarithmic terms, that of the Galerkin discretization of a single instance of the mean field problem. We illustrate the abstract theory for nonstationary parabolic diffusion problems in random domains. We verify Fréchet differentiability by means of shape calculus, and establish the Hadamard principle that the first order, k -th moment equation is completely specified in terms of data on the boundary of the nominal space-time cylinder. We perform boundary reduction of this parabolic evolution problem and propose a novel sparse tensor space-time Galerkin discretization. In conjunction with the sparse tensor Galerkin approximation of the k -point correlation, it reduces the complexity of the Galerkin discretization to $\mathcal{O}(N(\log N)^{k-1})$ where N denotes the number of degrees of freedom for a *stationary problem* on the *boundary of the nominal domain* (rather than on the space-time cylinder), thereby generalizing [25] to the boundary reduction of parabolic problems.

2010 *Mathematics Subject Classification.* Primary 65N30, 65J15.

Key words and phrases. Nonlinear operator equations, random parameters, deterministic methods, Fréchet derivative, sparse tensor approximation, random domain.

This work was supported in part under grant ERC AdG 247277 (to CS), initiated at the Institute for Mathematics and Applications (IMA) in Minnesota, USA and completed at the Hausdorff Research Institute for Mathematics, Bonn, Germany during the trimester “High dimensional problems” May - August 2011. AC acknowledges support by the Hausdorff Center for Mathematics, Bonn and CS the excellent working conditions at the Hausdorff Research Institute.

1. INTRODUCTION

We consider uncertainty quantification for abstract, parametric, nonlinear problems: given $\alpha \in Z$, find $u \in X$ such that

$$(1.1) \quad J(\alpha, u) = 0 \quad \text{in } Y'.$$

Here X, Y, Z are Banach spaces, Y' is the dual of Y , $u \in X$ is the unknown solution and $\alpha \in Z$ represents the set of inputs (such as boundary conditions, operator coefficients, shape of the physical domain of u , etc.). We allow the functional

$$(1.2) \quad J : \begin{cases} Z \times X & \rightarrow Y' \\ (\alpha, u) & \mapsto J(\alpha, u) \end{cases}$$

to depend nonlinearly on both variables, but insist on it to satisfy structural assumptions which render it locally, in a vicinity of a *nominal parameter value* $\alpha_0 \in Z$, well-posed (these structural assumptions on (1.1) will be made precise later): there exists a unique, “nominal” solution $u_0 \in X$ such that $J(\alpha_0, u_0) = 0$ in Y' .

Well-posedness of problem (1.1) implies moreover that changes in the problem data in some (usually sufficiently small) open neighborhood $U \subset Z$ of the nominal parameter value $\alpha_0 \in U$ lead to a corresponding small change of $u_0 \in X$ such that (1.1) is satisfied. We denote the solution operator of (1.1) by $S : U \rightarrow X$

$$(1.3) \quad u = S(\alpha) \quad \text{in } X, \quad \alpha \in U$$

and assume that S is injective in U .

In many practical applications, the effective values of the problem parameters α are not known precisely, but are known (or assumed) to be close to certain *nominal values* α_0 so that the fluctuation

$$r = \alpha - \alpha_0 \neq 0.$$

Uncertainty in the parameter values can be accounted for by probabilistic modelling of the input’s fluctuation $r = r(\omega) = \alpha - \alpha_0$ as a random field with known law or statistical moments. We show that if $\alpha(\omega) = \alpha_0 + r(\omega)$ belongs to U almost surely, $u = u(\omega)$ becomes a well defined random field (cf. Lemma 4.1 below). One goal of this work to present *deterministic* strategies towards the efficient, deterministic computation of its statistics. Our approach is based on the (formal, at this stage) first order Taylor expansion of J at the nominal values (α_0, u_0) : if $J(\alpha, u)$ is twice Fréchet differentiable at the nominal values (α_0, u_0) , say, then for all (α, u) in a vicinity of (α_0, u_0) holds

$$(1.4) \quad \begin{aligned} J(\alpha, u) &= J(\alpha_0, u_0) + J'_\alpha(\alpha_0, u_0)(\alpha - \alpha_0) + J'_u(\alpha_0, u_0)(u - u_0) \\ &\quad + \mathcal{O}(\|\alpha - \alpha_0\|_Z^2) + \mathcal{O}(\|u - u_0\|_X^2). \end{aligned}$$

Here, the differentials $J'_\alpha(\alpha_0, u_0) \in \mathcal{L}(Z, Y')$ and $J'_u(\alpha_0, u_0) \in \mathcal{L}(X, Y')$ are bounded, linear operators which are assumed to be known. Using (1.1) and neglecting the second order remainders yields the *first order perturbation equation*

$$(1.5) \quad 0 = J'_\alpha(\alpha_0, u_0)(\alpha - \alpha_0) + J'_u(\alpha_0, u_0)(u - u_0) \quad \text{in } Y'.$$

Well-posedness of (1.1) implies that the differential $J'_u(\alpha, u)$ is boundedly invertible in a vicinity of the nominal pair (α_0, u_0) . Then

$$\begin{aligned} u - u_0 = S(\alpha) - S(\alpha_0) &= -[J'_u(\alpha_0, u_0)]^{-1} J'_\alpha(\alpha_0, u_0)(\alpha - \alpha_0) + \mathcal{O}(\|\alpha - \alpha_0\|_Z^2) \\ &= S'(\alpha_0)(\alpha - \alpha_0) + \mathcal{O}(\|\alpha - \alpha_0\|_Z^2). \end{aligned}$$

The aim of this paper is to develop *fast, deterministic algorithms* for the approximate computation of the k -th statistical moment $\mathcal{M}^k[u - u_0]$ (also referred to as “ k -point correlation function”) of the random solution $u(\omega)$ from known k -th moment of the perturbation $\mathcal{M}^k r$. Our algorithms will be based on trading randomness for high dimensionality: rather sampling (1.1) for numerous realizations of α , we derive an *abstract, deterministic tensor equation* for the k -point correlation function of the system’s stochastic response: in Theorem 4.8 we prove that $\mathcal{M}^k[u - u_0]$ can be approximated by $\mathcal{M}^k[S'(\alpha_0)r]$, where $S'(\alpha_0) \in \mathcal{L}(Z, X)$ is the Fréchet derivative of the solution operator (1.3) at α_0 . One main contribution of the present paper are explicit bounds for the approximation error

$$(1.6) \quad \mathcal{E}^k = \mathcal{M}^k[u - u_0] - \mathcal{M}^k[S'(\alpha_0)r].$$

The classical Implicit Function Theorem provides existence of a neighborhood U of α_0 such that S is differentiable in U . However, this classical result does not allow to control the size of U . With this we obtain only asymptotic estimates for the approximation error \mathcal{E}^k when the size of U tends to zero.

In order to derive explicit *non-asymptotic* bounds for \mathcal{E}^k we prove in the present paper variants of the classical Local Implicit Function Theorem with an explicit relation between the size of U and the size of the differentials of the nonlinear functional J . These results are Newton-Kantorovich-type theorems with quantitative control of the constants which may be of independent interest. The *local Lipschitz continuity* of J'_u , J'_α and the *local invertibility* of J'_u are the key properties allowing to control the size of U . We provide a detailed regularity analysis for $\mathcal{M}^k[u - u_0]$ and $\mathcal{M}^k[S(\alpha_0)r]$.

If the magnitude of \mathcal{E}^k is small \mathbb{P} -a.s., we propose to compute $\mathcal{M}^k[S'(\alpha_0)r]$ instead of $\mathcal{M}^k[u - u_0]$, which is a simpler task, since $S'(\alpha_0)r$ can be characterized as a solution of a *linear* operator equation with a random right-hand side. We then apply the tensorization technique [21, 24] to characterize $\mathcal{M}^k[S'(\alpha_0)r]$ as solution of a tensorized deterministic operator equation.

Ultimately, the success of this approach depends, however, on our ability to efficiently solve the tensorized, deterministic equation for the k -point correlation function. To this end, we extend and generalize the approach of [21] to the present setting by imposing further assumptions on the “sensitivity operator” $G_0 = J'_u(\alpha_0, u_0)$. We discuss this first in the particular case, when $X = Y$ are separable Hilbert spaces and when G_0 is coercive. As we showed in the linear case in [21], under additional assumptions on G_0 such as bounded invertibility on scales of smoothness spaces $\{X_s\}_{s \geq 0} \subset X$ and $\{Y_s\}_{s \geq 0} \subset Y$

In Section 4, we consider random perturbations of the parameter α in (1.1) which are formalized on a background probability space satisfying the Kolmogorov axioms of probability: throughout, \mathbb{P} shall denote a probability measure on the probability space (Ω, \mathcal{A}) . By capitals B, X, Y, Z we shall denote separable Banach spaces of functions. Solutions to parametric random boundary value problems will be sought in Bochner spaces $L^p(\Omega, \mathbb{P}; B)$ of p -summable, measurable mappings from (Ω, \mathcal{A}) to $(B, \mathcal{B}(B))$ where $\mathcal{B}(B)$ denotes the sigma algebra of Borel sets on the separable Banach space B .

The paper is organized as follows. We begin with a brief introduction to the differential calculus in Banach spaces in Section 2. In Section 3 we study properties of the implicit solution of (1.1) with a deterministic parameter α . In particular, we obtain explicit bounds on the neighborhood $U \ni \alpha_0$ where the solution operator

$u = S(\alpha)$ is well defined and is Fréchet differentiable. The size of U depends explicitly on smoothness properties of J and its Fréchet derivatives. In Section 4 we apply these results to (1.1) where α is a random field. In particular, we obtain asymptotic and non-asymptotic bounds on the approximation error \mathcal{E}^k from (1.6) in the natural tensor product norms, see Theorem 4.6, Theorem 4.8. In Section 6 we illustrate the developed abstract techniques on sparse space-time tensor BEM for instationary heat equation in random domains. In particular, we propose a sparse tensor Galerkin scheme for the approximation of the k -point correlation of the random solution at any point in the space-time cylinder with complexity of log-linear order in N , the number of degrees of freedom on the nominal boundary.

2. DIFFERENTIAL CALCULUS IN BANACH SPACES

In this section we recall definitions and results of differential calculus in Banach spaces needed in what follows (see e.g. [2, 3, 15, 5]).

Suppose X, Y and Z are Banach spaces and $U \subset Z$ is an open subset. We denote $C(U, X)$ the space of continuous (w.r.t. convergence in norm) maps $U \rightarrow X$ and by $\mathcal{L}(Z, X)$ the space of linear continuous maps $Z \rightarrow X$.

Definition 2.1. *Let U be an open subset of Z and $\alpha \in U$. A mapping F is called Fréchet differentiable at α if there exists a linear continuous mapping $A \in \mathcal{L}(Z, X)$ such that for*

$$(2.1) \quad R(r) = F(\alpha + r) - F(\alpha) - A(r) \quad \text{there holds} \quad \|R(r)\|_X = o(\|r\|_Z),$$

that is $\|R(r)\|_X/\|r\|_Z \rightarrow 0$ as $\|r\|_Z \rightarrow 0$. If it exists, A is uniquely defined and is called Fréchet derivative of F at $\alpha \in Z$ and denoted by $F'(\alpha)$. The mapping F is said to be Fréchet differentiable in $U \subset Z$ if F is differentiable at all $\alpha \in U$.

Consider a function $J(\cdot, u)$ for some fixed $u \in X$. Its differential with respect to α at u is the Fréchet derivative $J'_\alpha(\alpha, u) \in \mathcal{L}(Z, Y')$ at (α, u) with respect to α . It is uniquely defined by the requirement

$$(2.2) \quad \|J(\alpha + r, u) - J(\alpha, u) - J'_\alpha(\alpha, u)r\|_{Y'} = o(\|r\|_Z).$$

for all $r \in U$ in a neighborhood of $0 \in U$.

We will also require the following differentiation rule for composite maps.

Lemma 2.2. *(Chain rule) Let $F : U \rightarrow X$ be differentiable at $\alpha \in U$ and $G : V \rightarrow Y$ be differentiable at $u = F(\alpha) \in V$, where U and V are open subsets of Z and X respectively. Then $G \circ F : U \rightarrow Y$ is differentiable at α and there holds*

$$(2.3) \quad (G \circ F)'(\alpha)r = G'(u) [F'(\alpha)r], \quad \text{with } u = F(\alpha).$$

Theorem 2.3. *Suppose $F \in C^1(U, X)$ in an open, nonempty subset $U \subset Z$. Assume that $\alpha \in U$ and $r \in Z$ such that $[\alpha, \alpha + r] \subset U$. Then*

$$(2.4) \quad F(\alpha + r) - F(\alpha) = \int_0^1 F'(\alpha + r\theta)r \, d\theta.$$

The following classical result gives sufficient conditions for local solvability of the equation (1.1), cf. [15, X. §2], [3, Ch. 3.1], [2, Ch. 2].

Theorem 2.4 (Local Implicit Function Theorem). *Suppose X, Y, Z are Banach spaces and W is an open neighborhood of (α_0, u_0) in $Z \times X$. Suppose $J(\alpha_0, u_0) = 0$ and*

- (1) $J'_u(\alpha, u)$ exists in W and J, J'_u are continuous in (α_0, u_0)
- (2) the “sensitivity operator” $G_0 := J'_u(\alpha_0, u_0) \in \mathcal{L}(X, Y')$ admits a bounded inverse $\Gamma_0 \in \mathcal{L}(Y', X)$

Then there exists neighborhoods U of α_0 and V of u_0 and a unique mapping $S : U \rightarrow V$, such that

- a. $S(\alpha_0) = u_0$
- b. $J(\alpha, u) = 0$ in Y' is equivalent to $u = S(\alpha)$ in X for any $\alpha \in U$, i.e.

$$J(\alpha, S(\alpha)) = 0 \quad \text{holds in } Y' \quad \text{for any } \alpha \in U .$$

- c. $S \in C(U, X)$ if (2) holds and if (1) is replaced with the stronger assumption:
 - (1a) $J \in C(W, Y')$, $J'_u(\alpha, u)$ exists in W and J'_u is continuous in (α_0, u_0)
- d. if (2) holds and if (1) is replaced with the stronger assumption
 - (1b) $J'_u(\alpha, u)$, $J'_\alpha(\alpha, u)$ exist in W and J, J'_α, J'_u are continuous in (α_0, u_0) then S is differentiable in α_0 and there holds

$$(2.5) \quad S'(\alpha_0) = -\Gamma_0 J'_\alpha(\alpha_0, u_0) .$$

Slightly stronger assumptions on J are needed to ensure Lipschitz continuity of S in U . In particular, we have the following theorem.

Theorem 2.5. *Suppose the assumptions of Theorem 2.4 are satisfied and assume that $S : U \rightarrow X$ is the solution operator of (1.1). Assume in addition that*

- (1') $J \in C^1(W, Y')$
- (2') the “sensitivity operator” $G_1 := J'_u(\alpha_1, u_1) \in \mathcal{L}(X, Y')$ admits a bounded inverse $\Gamma_1 \in \mathcal{L}(Y', X)$ for any $\alpha_1 \in U$ and $u_1 = S(\alpha_1)$.

Then $S \in C^1(U, X)$ and there holds

$$(2.6) \quad S'(\alpha_1) = -\Gamma_1 J'_\alpha(\alpha_1, S(\alpha_1)) \quad \forall \alpha_1 \in U .$$

In what follows we utilize the following notations for closed and open balls in Banach spaces: for $\alpha \in Z$ and $\delta \geq 0$

$$(2.7) \quad B(\alpha_0, \delta) := \{\alpha \in Z : \|\alpha - \alpha_0\|_Z \leq \delta\}, \quad \mathring{B}(\alpha_0, \delta) := B(\alpha_0, \delta) \setminus \partial B(\alpha_0, \delta) .$$

In our notation, we do not mention the Banach space Z explicitly, since it is always clear from the context.

3. THE SIZE OF NEIGHBORHOODS

The assumptions of Theorem 2.4 and Theorem 2.5 are not sufficient to control the size of the neighborhood U of around α_0 where the solution operator $S(\alpha)$ is well defined and differentiable. Revisiting the classical proofs of this result, for example in [15, X.§2], [3, Ch. 3.1], allows to find a direct connection between the size of U and smoothness properties of J . The following result shows that the additional assumption of Lipschitz continuity of J'_α and J'_u at (α_0, u_0) is sufficient for existence of $S(\alpha)$ for $\alpha \in U$ with quantitative bounds on the size of U . This result is inspired by the Newton-Kantorovich Theorem in the form stated, for example, in [14, Ch. XVIII], [26, Ch. 5].

Theorem 3.1. *Suppose X, Y, Z are Banach spaces and W is an open neighborhood of $(\alpha_0, u_0) \in Z \times X$. Suppose $J(\alpha_0, u_0) = 0$ and that*

- (1) $J \in C(W, Y')$, J'_α, J'_u exist in W and are Lipschitz continuous in (α_0, u_0) ,

- (2) the “sensitivity operator” $G_0 := J'_u(\alpha_0, u_0) \in \mathcal{L}(X, Y')$ has a bounded inverse $\Gamma_0 \in \mathcal{L}(Y', X)$.

Define

$$(3.1) \quad \xi_0 := \|\Gamma_0 J'_\alpha(\alpha_0, u_0)\|_{Z \rightarrow X}$$

and η_0 being the smallest constant satisfying

$$(3.2) \quad \left. \begin{aligned} \|\Gamma_0 \{J'_\alpha(\alpha, u) - J'_\alpha(\alpha_0, u_0)\}\|_{Z \rightarrow X} \\ \|\Gamma_0 \{J'_u(\alpha, u) - J'_u(\alpha_0, u_0)\}\|_{X \rightarrow X} \end{aligned} \right\} \leq \eta_0 (\|\alpha - \alpha_0\|_Z + \|u - u_0\|_X)$$

for any $(\alpha, u) \in W$. Let

$$(3.3) \quad \delta_0 := \frac{1}{2\eta_0(1 + \xi_0)}, \quad \varepsilon_0(\delta) := \frac{1 - \delta\eta_0 - \sqrt{1 - 2\delta\eta_0(1 + \xi_0)}}{\eta_0}$$

and assume that $\delta < \delta_0$ is so small that $B(\alpha_0, \delta) \times B(u_0, \varepsilon_0(\delta)) \subset W$. Then the following holds:

- a. There exists a unique continuous mapping $S : B(\alpha_0, \delta) \mapsto B(u_0, \varepsilon_0(\delta))$ satisfying $J(\alpha, S(\alpha)) = 0$ for any $\alpha \in B(\alpha_0, \delta)$ and for $u_0 = S(\alpha_0)$.
- b. The mapping S satisfies

$$(3.4) \quad \|S(\alpha) - S(\alpha_0)\|_X \leq M_0 \|\alpha - \alpha_0\|_Z \quad \forall \alpha \in B(\alpha_0, \delta)$$

with Lipschitz constant $M_0 = 2\xi_0 + 1$.

Proof. Define a nonlinear map $A_\alpha : X \rightarrow X$ by

$$(3.5) \quad A_\alpha u := u - \Gamma_0 J(\alpha, u).$$

We show first that $\exists \varepsilon > 0$ such that (i) A_α is a contraction in $B(u_0, \varepsilon)$ and (ii) maps $B(u_0, \varepsilon)$ into itself.

Define $R(\alpha, u) := J(\alpha, u) - J'_u(\alpha_0, u_0)(u - u_0)$. Then from (3.5)

$$A_\alpha u = u_0 - \Gamma_0 R(\alpha, u)$$

and hence by linearity of Γ_0

$$A_\alpha u - A_\alpha u_1 = -\Gamma_0(R(\alpha, u) - R(\alpha, u_1)).$$

For the right-hand side we have by the definition of R and Theorem 2.3

$$\begin{aligned} \Gamma_0(R(\alpha, u) - R(\alpha, u_1)) &= \Gamma_0(J(\alpha, u) - J(\alpha, u_1) - J'_u(\alpha_0, u_0)(u - u_1)) \\ &= \int_0^1 \Gamma_0 \left\{ J'_u(\alpha, tu + (1-t)u_1) - J'_u(\alpha_0, u_0) \right\} (u - u_1) dt \end{aligned}$$

Then

$$(3.6) \quad \begin{aligned} \frac{\|A_\alpha u - A_\alpha u_1\|_X}{\|u - u_1\|_X} &\leq \int_0^1 \eta_0 \left\{ \|\alpha - \alpha_0\|_Z + t\|u - u_0\|_X + (1-t)\|u_1 - u_0\|_X \right\} dt \\ &= \eta_0 \left\{ \|\alpha - \alpha_0\|_Z + \frac{1}{2}\|u - u_0\|_X + \frac{1}{2}\|u_1 - u_0\|_X \right\} =: K \end{aligned}$$

We have $K < 1$, i.e. A_α is a contraction for any $\alpha \in B(\alpha_0, \delta)$ and $u, u_1 \in B(u_0, \varepsilon)$, provided that

$$(3.7) \quad \eta_0(\delta + \varepsilon) < 1 \quad \Leftrightarrow \quad \varepsilon \leq \frac{1 - \eta_0\delta}{\eta_0}.$$

For proving (ii) we need to show

$$(3.8) \quad \|A_\alpha u - u_0\|_X \leq \varepsilon \quad \forall u \in B(u_0, \varepsilon).$$

Using (3.5) and (3.6) we have for any $u \in B(u_0, \varepsilon)$

$$(3.9) \quad \begin{aligned} \|A_\alpha u - u_0\|_X &\leq \|A_\alpha u - A_\alpha u_0\|_X + \|A_\alpha u_0 - u_0\|_X \\ &\leq \eta_0 \left(\delta + \frac{\varepsilon}{2} \right) \varepsilon + \|\Gamma_0 J(\alpha, u_0)\|_X. \end{aligned}$$

Recalling $J(\alpha_0, u_0) = 0$ we estimate the last term via triangle inequality

$$\begin{aligned} \|\Gamma_0 J(\alpha, u_0)\|_X &\leq \|\Gamma_0 J'_\alpha(\alpha_0, u_0)(\alpha - \alpha_0)\|_X \\ &\quad + \|\Gamma_0 \{J(\alpha, u_0) - J(\alpha_0, u_0) - J'_\alpha(\alpha_0, u_0)(\alpha - \alpha_0)\}\|_X \end{aligned}$$

The first term in the right-hand side is bounded by $\xi_0 \|\alpha - \alpha_0\|_Z$, the expression in the norm in the last term equals to

$$\int_0^1 \Gamma_0 \left\{ J'_\alpha(s\alpha + (1-s)\alpha_0, u_0) - J'_\alpha(\alpha_0, u_0) \right\} (\alpha - \alpha_0) ds.$$

This gives

$$\begin{aligned} \|\Gamma_0 J(\alpha, u_0)\|_X &\leq \|\alpha - \alpha_0\|_Z \left(\xi_0 + \int_0^1 \eta_0 s \|\alpha - \alpha_0\|_Z ds \right) \\ &= \|\alpha - \alpha_0\|_Z \left(\xi_0 + \frac{\eta_0}{2} \|\alpha - \alpha_0\|_Z \right). \end{aligned}$$

Denote $\delta := \|\alpha - \alpha_0\|$, then (3.9) provides

$$(3.10) \quad \|A_\alpha u - u_0\|_X \leq \eta_0 \left(\delta + \frac{\varepsilon}{2} \right) \varepsilon + \left(\xi_0 + \frac{\eta_0}{2} \delta \right) \delta.$$

Therefore A_α maps $B(u_0, \varepsilon)$ into itself if right-hand side of (3.10) is bounded by ε , i.e. the radii $\delta, \varepsilon > 0$ satisfy

$$(3.11) \quad \eta_0 \left(\delta + \frac{\varepsilon}{2} \right) \varepsilon + \left(\xi_0 + \frac{\eta_0}{2} \delta \right) \delta \leq \varepsilon \quad \Leftrightarrow \quad \tilde{\varepsilon}^2 - 2(1 - \tilde{\delta})\tilde{\varepsilon} + 2\xi_0\tilde{\delta} + \tilde{\delta}^2 \leq 0.$$

Here $\tilde{\delta} := \eta_0 \delta$, $\tilde{\varepsilon} := \eta_0 \varepsilon$ are the rescaled radii. Resolving (3.11) and using (3.7) we find that $\tilde{\varepsilon} \in \mathcal{U}_{\tilde{\delta}} := [\tilde{\varepsilon}_-, \tilde{\varepsilon}_+] \cap (0, 1 - \tilde{\delta})$ where

$$\tilde{\varepsilon}_\pm = 1 - \tilde{\delta} \pm \sqrt{D} \quad \text{with} \quad D = (1 - \tilde{\delta})^2 - (2\xi_0\tilde{\delta} + \tilde{\delta}^2) = 1 - 2\tilde{\delta}(1 + \xi_0).$$

Basic calculations show that $\mathcal{U}_{\tilde{\delta}} \equiv [\tilde{\varepsilon}_-, 1 - \tilde{\delta})$ is nonempty as soon as $D > 0$, since in this case $\tilde{\varepsilon}_- > 0$ and

$$(3.12) \quad \tilde{\delta} < \frac{1}{2(1 + \xi_0)} < \frac{1}{2}.$$

Herewith we have proven that if $\tilde{\delta} := \eta_0 \delta$ satisfies (3.12), i.e. if $\delta < \delta_0$, $\alpha \in U := B(\alpha_0, \delta)$ and if $\varepsilon_0(\delta) := \tilde{\varepsilon}_- / \eta_0$ then the mapping A_α from (3.5) is a contraction on $B(u_0, \varepsilon_0(\delta))$. Furthermore, (3.5) yields that $A_\alpha(u)$ is continuous for all $(\alpha, u) \in W$, because $J \in C(W, Y')$ and Γ_0 is linear. Then the assumptions of [3, Corollary 3.1.4] are satisfied and there exists a unique $S \in C(U, X)$ satisfying $J(\alpha, S(\alpha)) = 0$ in U and $u_0 = S(\alpha_0)$.

Next, we prove the estimate (3.4). From the previous considerations, we know in particular that $\alpha \in \partial B(\alpha_0, \delta)$ implies $S(\alpha) \in B(u_0, \varepsilon_0(\delta))$, if $\delta < \delta_0$. Then

$$(3.13) \quad \|S(\alpha) - S(\alpha_0)\|_X \leq \varepsilon_0(\delta) = \frac{1 - \tilde{\delta} - \sqrt{1 - 2\tilde{\delta}(1 + \xi_0)}}{\eta_0} \leq \frac{(2\xi_0 + 1)\tilde{\delta}}{\eta_0}.$$

Then (3.4) follows, since $\delta = \|\alpha - \alpha_0\|_Z = \tilde{\delta}/\eta_0$. \square

This result does not provide computable quantitative bounds on the size of the neighborhood of α_0 in which S is differentiable. To obtain such bounds, we require additional assumptions on J .

Theorem 3.2. *Suppose that the assumptions of Theorem 3.1 are satisfied, that δ_0, ε_0 are defined as in Theorem 3.1 and that $S : B(\alpha_0, \delta) \rightarrow B(u_0, \varepsilon_0(\delta))$, $\delta < \delta_0$ is the solution operator. Assume in addition*

- (1') $J \in C^1(W, Z)$ and J'_α, J'_u are Lipschitz continuous in W .
- (2') $J'_u(\alpha_1, u_1)$ with $u_1 = S(\alpha_1)$ has a uniformly bounded inverse $\Gamma_1 : Y' \rightarrow X$ for any α_1 in the open ball $\mathring{B}(\alpha_0, \delta_0)$.

Define

$$(3.14) \quad \xi := \sup_{\alpha_1 \in \mathring{B}(\alpha_0, \delta_0)} \|\Gamma_1 J'_\alpha(\alpha_1, S(\alpha_1))\|_{Z \rightarrow X}$$

and η being the smallest constant satisfying

$$(3.15) \quad \left. \begin{aligned} \|\Gamma_1 \{J'_\alpha(\alpha, u) - J'_\alpha(\alpha_1, u_1)\}\|_{Z \rightarrow X} \\ \|\Gamma_1 \{J'_u(\alpha, u) - J'_u(\alpha_1, u_1)\}\|_{X \rightarrow X} \end{aligned} \right\} \leq \eta (\|\alpha - \alpha_1\|_Z + \|u - u_1\|_X).$$

uniformly for any $(\alpha, u), (\alpha_1, u_1) \in W$. Let

$$(3.16) \quad \delta_* := \frac{1}{2\eta(1 + \xi)}, \quad \varepsilon_*(\delta) := \frac{1 - \delta\eta - \sqrt{1 - 2\delta\eta(1 + \xi)}}{\eta}$$

and assume that $\delta < \delta_*/2$ is small enough, such that $B(\alpha_0, \delta) \times B(u_0, \varepsilon_*(\delta)) \subset W$. Then

- (a) S is Lipschitz continuous in $B(\alpha_0, \delta)$, in particular for $M := 2\xi + 1$

$$(3.17) \quad \|S(\alpha) - S(\alpha_1)\|_X \leq M\|\alpha - \alpha_1\|_Z \quad \forall \alpha, \alpha_1 \in B(\alpha_0, \delta).$$

- (b) S is differentiable in $B(\alpha_0, \delta)$ and

$$(3.18) \quad S'(\alpha_1) = -\Gamma_1 J'_\alpha(\alpha_1, S(\alpha_1)) \quad \forall \alpha_1 \in B(\alpha_0, \delta).$$

- (c) S' in Lipschitz continuous in α_0

$$(3.19) \quad \|S'(\alpha) - S'(\alpha_0)\|_{Z \rightarrow X} \leq K\|\alpha - \alpha_0\|_Z \quad \forall \alpha \in B(\alpha_0, \delta)$$

with the Lipschitz constant $K = 4\eta_0(1 + \xi_0)^2$.

Proof. (a) If we fix an arbitrary $\alpha_1 \in B(\alpha_0, \delta_0)$, then assumptions of Theorem 3.1 are satisfied with α_1 instead of α_0 . We resort to the uniform constants ξ, η and obtain the estimate

$$(3.20) \quad \|S(\alpha) - S(\alpha_1)\|_X \leq M\|\alpha - \alpha_1\|_Z \quad \forall \alpha_1 \in \mathring{B}(\alpha_0, \delta_0), \quad \forall \alpha \in \mathring{B}(\alpha_1, \delta_*).$$

with $M = 2\xi + 1$. Suppose $\alpha, \alpha_1 \in B(\alpha_0, \delta)$ with $\delta < \delta_*/2$. Then $\alpha_1 \in \mathring{B}(\alpha_0, \delta_0)$ and $\alpha \in \mathring{B}(\alpha_1, \delta_*)$, since

$$\|\alpha_1 - \alpha_0\|_Z \leq \delta_*/2 < \delta_0 \quad \text{and} \quad \|\alpha - \alpha_1\|_Z \leq \|\alpha - \alpha_0\|_Z + \|\alpha_1 - \alpha_0\|_Z \leq 2\delta < \delta_*$$

which together with (3.20) yields (3.17).

(b) We prove differentiability of S in $B(\alpha_0, \delta)$. To this end we fix an arbitrary element $\alpha_1 \in B(\alpha_0, \delta)$ and define

$$(3.21) \quad u_1 := S(\alpha_1), \quad \Gamma_1 := [J'_u(\alpha_1, u_1)]^{-1}, \quad \Lambda_1 := -\Gamma_1 J'_\alpha(\alpha_1, u_1).$$

Recall $J(\alpha_1, u_1) = 0 = J(\alpha, u)$. Then we have for $u := S(\alpha)$

$$\begin{aligned} \|S(\alpha) - S(\alpha_1) - \Lambda_1(\alpha - \alpha_1)\|_X &= \|\Gamma_1 \{J'_\alpha(\alpha_1, u_1)(\alpha - \alpha_1) - J'_u(\alpha_1, u_1)(u - u_1)\}\|_X \\ &\leq \int_0^1 \|\Gamma_1 (J'_\alpha(s\alpha + (1-s)\alpha_1, u) - J'_\alpha(\alpha_1, u_1))\|_{Z \rightarrow X} \|\alpha - \alpha_1\|_Z \, ds \\ &\quad + \int_0^1 \|\Gamma_1 (J'_u(\alpha_1, tu + (1-t)u_1) - J'_u(\alpha_1, u_1))\|_{X \rightarrow X} \|u - u_1\|_X \, dt \\ &\leq \eta \left\{ \left(\frac{1}{2} \|\alpha - \alpha_1\|_Z + \|u - u_1\|_X \right) \|\alpha - \alpha_1\|_Z + \frac{1}{2} \|u - u_1\|_X^2 \right\} \\ &\leq \frac{\eta}{2} (1 + M)^2 \|\alpha - \alpha_1\|_Z^2 \end{aligned}$$

by Lipschitz continuity of S in $B(\alpha_0, \delta)$, cf. (3.17). The right-hand side is $\mathcal{O}(\|\alpha - \alpha_1\|_Z^2)$, thus by Definition 2.1, S is differentiable in any $\alpha_1 \in B(\alpha_0, \delta)$ and $S'(\alpha_1) \equiv \Lambda_1$.

(c) Finally, we prove that S' is Lipschitz continuous in u_0 . Relation (3.18) is equivalent to

$$J'_u(\alpha, S(\alpha))S'(\alpha) + J'_\alpha(\alpha, S(\alpha)) = 0, \quad \forall \alpha \in B(\alpha_0, \delta).$$

Then for $u = S(\alpha)$, $u_0 = S(\alpha_0)$ we have the identity

$$\begin{aligned} (J'_u(\alpha, u) - J'_u(\alpha_0, u_0))S'(\alpha) + J'_u(\alpha_0, u_0)(S'(\alpha) - S'(\alpha_0)) \\ + J'_\alpha(\alpha, u) - J'_\alpha(\alpha_0, u_0) = 0. \end{aligned}$$

This yields with $\Gamma_0 = [J'_u(\alpha_0, u_0)]^{-1}$

$$\begin{aligned} \|S'(\alpha) - S'(\alpha_0)\|_{Z \rightarrow X} &\leq \|\Gamma_0 (J'_u(\alpha, u) - J'_u(\alpha_0, u_0))\|_{X \rightarrow X} \|S'(\alpha)\|_{Z \rightarrow X} \\ &\quad + \|\Gamma_0 (J'_\alpha(\alpha, u) - J'_\alpha(\alpha_0, u_0))\|_{Z \rightarrow X} \\ &\leq \eta_0 (\|\alpha - \alpha_0\|_Z + \|S(\alpha) - S(\alpha_0)\|_X) (1 + \|S'(\alpha)\|_{Z \rightarrow X}) \\ &\leq \Theta \|\alpha - \alpha_0\|_Z (1 + \|S'(\alpha_0)\|_{Z \rightarrow X} + \|S'(\alpha) - S'(\alpha_0)\|_{Z \rightarrow X}) \end{aligned}$$

with $\Theta = \eta_0(M_0 + 1) = 2\eta_0(1 + \xi_0) = 1/\delta_0$. Assume $\alpha \in B(\alpha_0, \delta)$, $\delta < \delta_*/2$. Then $\Theta \|\alpha - \alpha_0\|_Z < \delta_*/(2\delta_0) \leq 1/2$. This implies

$$\|S'(\alpha) - S'(\alpha_0)\|_{Z \rightarrow X} \leq 2\Theta(1 + \xi_0) \|\alpha - \alpha_0\|_Z.$$

The proof is complete. \square

4. RANDOM FIELDS UNDER NONLINEAR MAPS

In this section we study properties of random fields and their characteristics under nonlinear maps and apply it to solution operators of the implicit nonlinear equation (1.1). We prove the basic abstract results on measurability (Lemma 4.1) and summability (Lemma 4.2) properties of random fields under nonlinear maps. In combination with Theorem 2.4, this yields existence of the moments of the solution

in a sufficiently small neighborhood U about the nominal parameter value α_0 , cf. Theorem 4.6 and the asymptotic approximation result (4.16). Theorems 3.1 and 3.2 from the previous section allow to control the size of U . Based on this, we obtain non-asymptotic approximation bounds in Theorem 4.8 which allows for $\alpha \in U$ a.s. where U is not necessarily small. The key property here is Lipschitz continuity of the solution operator, cf. Lemma 4.7.

We begin with basic definitions, see e.g. [8, 17, 16]. Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space where Ω is a sample space, \mathcal{A} is a set of events being a σ -algebra over Ω , and \mathbb{P} is a probability measure. If (X, \mathcal{F}) and (Y, \mathcal{G}) are measurable spaces, a map $f : X \rightarrow Y$ is called \mathcal{F} - \mathcal{G} -measurable, if $f^{-1}(\mathcal{G}) \subseteq \mathcal{F}$. For a complete metric space E , we shall denote by $\mathcal{B}(E)$ the Borel σ -algebra of E generated by all open subsets of E . A mapping $\alpha : \Omega \rightarrow E$ is called a random variable if it is \mathcal{F} - $\mathcal{B}(E)$ -measurable. In this paper we are interested in random variables with values in Banach spaces. From now we assume that X, Y and Z are Banach spaces.

Under the above notations we introduce the set of all \mathcal{A} -measurable functions

$$(4.1) \quad L^0(\Omega; Z) := \{\alpha : \Omega \rightarrow Z : \alpha \text{ is } \mathcal{A}\text{-}\mathcal{B}(Z)\text{-measurable}\}$$

that is the set of all random variables $\Omega \rightarrow Z$. Suppose $\alpha \in L^0(\Omega, Z)$ and $S : Z \rightarrow X$ is a general nonlinear map (in what follows, S shall be interpreted as solution operator from Section 2). Sufficient conditions under which the composition $S \circ \alpha$ is a random variable, are given by the following classical result.

Lemma 4.1. *Suppose X, Z are Banach spaces, $\alpha \in L^0(\Omega, Z)$ and $S : Z \rightarrow X$. The mapping $u := S \circ \alpha : \Omega \rightarrow X$ is a random variable if S is Borel measurable, that is $\mathcal{B}(Z)$ - $\mathcal{B}(X)$ -measurable. A sufficient condition is $S \in C(\alpha(\Omega), X)$.*

Proof. We prove the sufficient condition. As S is continuous, $S^{-1}(V)$ is open in $\alpha(\Omega)$ for any $V \in \mathcal{B}(X)$, i.e. $\exists U \in \mathcal{B}(Z) : S^{-1}(V) = \alpha(\Omega) \cap U$. Then

$$(S \circ \alpha)^{-1}(V) = \alpha^{-1}(S^{-1}(V)) = \alpha^{-1}(\alpha(\Omega) \cap U) = \alpha^{-1}(U) \in \mathcal{A},$$

since $\alpha \in L^0(\Omega, Z)$ by assumption. \square

A function $\alpha : \Omega \rightarrow Z$ is called \mathbb{P} -almost surely separable valued if there exists a \mathbb{P} -null set N such that the image of its complement $\alpha(\Omega \setminus N)$ is separable. In order to introduce a Bochner-Lebesgue integral we resort to a subspace of \mathbb{P} -measurable random variables, which can be equivalently defined by

$$(4.2) \quad L^0(\Omega, \mathbb{P}; Z) := \{\alpha \in L^0(\Omega; Z) : \alpha \text{ is } \mathbb{P}\text{-almost surely separable valued}\}.$$

The space $L^0(\Omega, \mathbb{P}, Z)$ is also called a Bochner space of *strongly measurable functions*. For a Banach space Z we define

$$(4.3) \quad \|\alpha\|_{L^k(\Omega, \mathbb{P}; Z)} := \begin{cases} \left(\int_{\Omega} \|\alpha(\omega)\|_Z^k d\mathbb{P}(\omega) \right)^{1/k} & 1 \leq k < \infty \\ \text{ess sup}_{\omega \in \Omega} \|\alpha(\omega)\|_Z & k = \infty \end{cases}$$

where $\mathbb{E}[\cdot] := \int_{\Omega} (\cdot) d\mathbb{P}(\omega)$ with the integral denoting the Bochner-Lebesgue integral. Then

$$(4.4) \quad L^k(\Omega, \mathbb{P}; Z) := \{\alpha \in L^0(\Omega, \mathbb{P}; Z) : \|\alpha\|_{L^k(\Omega, \mathbb{P}; Z)} < \infty\} / \mathcal{N}$$

where $\mathcal{N} = \{\alpha \in L^0(\Omega, \mathbb{P}; Z) : \alpha = 0 \text{ } \mathbb{P}\text{-a.e.}\}$. The Bochner-Lebesgue space $L^k(\Omega, \mathbb{P}; Z)$ is a Banach space with the norm (4.3).

Throughout our analysis of random perturbations, we assume w.l.o.g. that the nominal parameter value $\alpha_0 \in U \subset Z$ in our linearization of the solution map $S(\alpha)$ is the mean field of $\alpha(\omega)$ or, equivalently, that the parameter fluctuation's mean field $r(\omega) = \alpha(\omega) - \alpha_0$ vanishes.

Lemma 4.2. *Suppose X, Z are Banach spaces, $\alpha \in L^k(\Omega, \mathbb{P}; Z)$ and that $S : Z \rightarrow X$ is Lipschitz continuous at $\alpha_0 \in Z$: there exists an open neighborhood $U \subseteq Z$ of α_0 and a constant $M > 0$, such that*

$$(4.5) \quad \|S(\tilde{\alpha}) - S(\alpha_0)\|_X \leq M \|\tilde{\alpha} - \alpha_0\|_Z \quad \forall \tilde{\alpha} \in U.$$

Assume in addition that

$$(4.6) \quad \mathbb{P}\{\alpha \in U\} = 1.$$

Then $u = S(\alpha) \in L^k(\Omega, \mathbb{P}; X)$.

Proof. Suppose $1 \leq k < \infty$. Then

$$(4.7) \quad \|S(\alpha) - S(\alpha_0)\|_{L^k(\Omega, \mathbb{P}; X)} \leq M \|\alpha - \alpha_0\|_{L^k(\Omega, \mathbb{P}; Z)} \quad \forall \alpha \in U \quad \mathbb{P}\text{-a.e.}$$

since

$$\int_{\Omega} \|S(\alpha) - S(\alpha_0)\|_X^k d\mathbb{P}(\omega) \leq M^k \int_{\Omega} \|\alpha - \alpha_0\|_Z^k d\mathbb{P}(\omega).$$

From (4.5) we infer that (4.7) holds also for $k = \infty$. \square

In what follows we shall introduce and study properties of k -th statistical moments. For that it is convenient to introduce the following abbreviations for tensor products.

Definition 4.3. *For any two Banach spaces X, Y we denote by $X \otimes Y$ its tensor product being a Banach space endowed with a reasonable cross-norm satisfying*

$$(4.8) \quad \|u \otimes v\|_{X \otimes Y} = \|u\|_X \|v\|_Y, \quad \forall u \in X, v \in Y.$$

The k -fold tensor product $X \otimes \cdots \otimes X$ is defined by induction and is denoted by $X^{(k)}$. Analogously for an element $u \in X$ we define its k -fold product $u^{(k)} := u \otimes \cdots \otimes u \in X^{(k)}$ and a linear map $A \in \mathcal{L}(X, Y)$ we define its k -fold tensor product $A^{(k)} = A \otimes \cdots \otimes A \in \mathcal{L}(X^{(k)}, Y^{(k)})$.

Definition 4.4. *Let $k \geq 1$ be an integer. Then for $\alpha \in L^k(\Omega, \mathbb{P}; Z)$ its k -th statistical moment is defined by*

$$(4.9) \quad \mathcal{M}^k \alpha = \mathbb{E} \left[\alpha^{(k)} \right] = \int_{\Omega} \alpha^{(k)} d\mathbb{P}(\omega) \in Z^{(k)}.$$

Note that \mathcal{M}^k is well defined as a mapping $L^k(\Omega, \mathbb{P}; Z) \rightarrow Z^{(k)}$, since

$$(4.10) \quad \|\mathcal{M}^k \alpha\|_{Z^{(k)}} \leq \|\alpha^{(k)}\|_{L^1(\Omega, \mathbb{P}; Z^{(k)})} = \|\alpha\|_{L^k(\Omega, \mathbb{P}; Z)}^k.$$

Corollary 4.5. *Suppose X, Y, Z are Banach spaces and W is a neighborhood of $(\alpha_0, u_0) \in Z \times X$. Let $\alpha \in L^0(\Omega; Z)$ be a random variable and consider a mapping J is defined in (1.2). Suppose $J(\alpha_0, u_0) = 0$ and*

- (1) $J \in C(W, Y')$ and its differential $J'_u(\alpha, u) \in \mathcal{L}(X, Y')$ exist in W and are continuous in (α_0, u_0) .
- (2) the ‘‘sensitivity operator’’ $G_0 := J'_u(\alpha_0, u_0)$ admits a bounded inverse $\Gamma_0 \in \mathcal{L}(Y', X)$.

Then there is a neighborhood U of α_0 and a unique random variable $u \in L^0(\Omega; X)$ such that $u = S(\alpha)$ satisfies (1.1) if $\mathbb{P}\{\alpha(\omega) \in U\} = 1$; and $S(\alpha_0) = u_0$.

Suppose (1) is replaced with the stronger assumption:

- (1') $J \in C(W, Y')$, $J'_\alpha \in \mathcal{L}(Z, Y')$, $J'_u \in \mathcal{L}(X, Y')$ exist in W and are Lipschitz continuous in (α_0, u_0)

Then

- (a) U can be chosen as $U = B(\alpha_0, \delta)$, $\delta < \delta_0$ with δ_0 from Theorem 3.1.
(b) If $\mathbb{P}\{\|\alpha(\omega) - \alpha_0\|_Z \leq \delta\} = 1$, then $\alpha \in L^k(\Omega, \mathbb{P}; Z)$ and $u \in L^k(\Omega, \mathbb{P}; X)$ for any $1 \leq k \leq \infty$.

Proof. Existence of a unique solution operator $S \in C(U, X)$ follows from Theorem 2.4. Lemma 4.1 yields that $u = S(\alpha)$ is $\mathcal{A}\text{-}\mathcal{B}(X)$ -measurable. Theorem 3.1 yields (a). Assertion (b) follows from

$$\|\alpha - \alpha_0\|_{L^k(\Omega, \mathbb{P}; Z)} \leq \delta_0, \quad \|u - u_0\|_{L^k(\Omega, \mathbb{P}; X)} \leq M\delta_0$$

guaranteed by (3.4) and Lemma 4.2. \square

We see from Corollary 4.5 that $u \in L^k(\Omega, \mathbb{P}; X)$ for any k , if J'_α, J'_u are Lipschitz continuous at (α_0, u_0) and the random parameter $\alpha(\omega)$ belongs \mathbb{P} -a.s. to a sufficiently small (bounded) neighborhood of $\alpha_0 \in Z$. This assumption is however of limited use in practice. In particular, α can not be modeled as a Gaussian random field taking arbitrary large values with small but *positive* probability. This difficulty cannot be overcome in general due to the nonlinear nature of J , e.g. the implicit equation (1.1) might be not uniquely solvable for certain values of α far from α_0 . Fortunately, for some classes of J unbounded values of α can be allowed, as we show in Lemma 4.7 below.

Considering models with such unbounded random parameters is possible if J is linear and continuous [21, 24, 6]. In this case the solution operator S is linear and

$$(4.11) \quad \mathcal{M}^k u = \mathbb{E}[S(\alpha) \otimes \cdots \otimes S(\alpha)] = (S \otimes \cdots \otimes S)\mathcal{M}^k \alpha.$$

In the *linear case* we obtain from (1.5)

$$(4.12) \quad J'_u u = -J'_\alpha \alpha \iff u = S(\alpha) = S' \alpha = -[J'_u]^{-1} J'_\alpha \alpha,$$

where S', J'_u, J'_α do not depend on (α, u) . If J'_u is invertible, then $\alpha \in L^k(\Omega, \mathbb{P}; Z)$ implies $u \in L^k(\Omega, \mathbb{P}; X)$ and hence the k -point correlation function to first order, $\mathcal{M}^k u$, is well-defined in $X^{(k)}$ by (4.10). In this case *sparse tensor* schemes can be constructed for approximate numerical evaluation of $S = -[J'_u]^{-1} J'_\alpha$, see [21, 24, 6].

In what follows we shall generalize this idea to a class of nonlinear problems via local implicit problems of the type (1.1). The starting point is to rewrite (4.11) in view of (4.12) in the form

$$(4.13) \quad \mathcal{M}^k u = \mathcal{M}^k S' \alpha = (S' \otimes \cdots \otimes S')\mathcal{M}^k \alpha.$$

Here both S and S' are linear operators, and S' does not depend on the evaluation point α_0 , see (2.1). If S is nonlinear, then S' depends on the evaluation point α_0 and (4.13) is not true anymore. However, $S'(\alpha_0) \in \mathcal{L}(Z, X)$ is still a bounded, linear operator and the nonlinear version of (4.13) reads

$$(4.14) \quad \mathcal{M}^k[u - u_0] \approx \mathcal{M}^k[S'(\alpha_0)(\alpha - \alpha_0)] = (S'(\alpha_0) \otimes \cdots \otimes S'(\alpha_0))\mathcal{M}^k[\alpha - \alpha_0].$$

In other words, $\mathcal{M}^k[S'(\alpha_0)(\alpha - \alpha_0)]$ can be computed pursuing the same strategy as in the linear case and requires only the knowledge of $\mathcal{M}^k[\alpha - \alpha_0]$ and of a

numerical procedure of approximate evaluation of the first variation at the nominal pair (α_0, u_0) ,

$$S'(\alpha_0) = -[J'_u(\alpha_0, u_0)]^{-1} J'_\alpha(\alpha_0, u_0) = -\Gamma_0 J'_\alpha(\alpha_0, u_0).$$

Since the exact inverse Γ_0 is usually not available, we interpret (4.14) as first order tensorized operator equation for the k -th moment $\mathcal{M}^k[u - u_0]$, i.e. we consider the tensorized equation

$$(4.15) \quad \text{find } Z^{(k)} \in X^{(k)} : \quad G_0^{(k)} Z^{(k)} = (-1)^k [J'_\alpha(\alpha_0, u_0)]^{(k)} \mathcal{M}^k[\alpha - \alpha_0]$$

where $G_0 = J'_u(\alpha_0, u_0)$. Based on (4.14), the solution $Z^{(k)} = \mathcal{M}^k[S'(\alpha_0)(\alpha - \alpha_0)]$ of (4.15) should be a good approximation of $\mathcal{M}^k[u - u_0]$. By the definition of the Fréchet derivative (2.1) we infer that the approximation must be good at least if α belongs to a sufficiently small neighborhood of α_0 as the following asymptotic result shows.

Theorem 4.6. *Suppose that the functional J and its derivatives satisfy the assumptions of Theorem 3.2 and $\mathbb{P}\{\|\alpha(\omega) - \alpha_0\|_Z \leq \delta\} = 1$. Then*

$$(4.16) \quad \|\mathcal{M}^k[u - u_0] - \mathcal{M}^k[S'(\alpha_0)(\alpha - \alpha_0)]\|_{X^{(k)}} = o(\delta^k), \quad \delta \rightarrow 0.$$

Proof. Due to Theorem 3.2, the solution operator S exist and is differentiable in $B(\alpha_0, \delta)$ for $\delta < \delta_*/2$. According to (2.1) we have for $u = S(\alpha)$, $u_0 = S(\alpha_0)$ and $r = \alpha - \alpha_0$

$$u = u_0 + S'(\alpha_0)r + R_0(r), \quad \|R_0(r)\|_X = o(\|r\|_Z).$$

Due to (4.9)

$$\mathcal{M}^k[u - u_0] - \mathcal{M}^k[S'(\alpha_0)r] = \mathbb{E}[(S'(\alpha_0)r + R_0(r))^{(k)}] - \mathbb{E}[(S'(\alpha_0)r)^{(k)}].$$

Hence, by property of the cross-norm (4.8) and the binomial theorem

$$\begin{aligned} \|\mathcal{M}^k[u - u_0] - \mathcal{M}^k[S'(\alpha_0)r]\|_{X^{(k)}} &\leq \mathbb{E}\left[\sum_{i=1}^k \binom{k}{i} \|S'(\alpha_0)r\|_X^{k-i} \|R_0(r)\|_Z^i\right] \\ &\leq \mathbb{E}\left[\sum_{i=1}^k \binom{k}{i} \|S'(\alpha_0)\|_{Z \rightarrow X}^{k-i} \|r\|_X^{k-i} \|R_0(r)\|_Z^i\right] \\ &\leq \mathbb{E}[\|r\|_Z^k] \sum_{i=1}^k \binom{k}{i} \xi_0^{k-i} \sup_{\|\tilde{r}\|_Z \leq \delta} \left(\frac{\|R_0(\tilde{r})\|_X}{\|\tilde{r}\|_Z}\right)^i. \end{aligned}$$

If $\mathbb{P}\{\|r(\omega)\|_Z \leq \delta\} = 1$ we have $\mathbb{E}[\|r\|_Z^k] \leq \delta^k$ and

$$\frac{\|\mathcal{M}^k[u - u_0] - \mathcal{M}^k[S'(\alpha_0)r]\|_{X^{(k)}}}{\delta^k} \rightarrow 0, \quad \delta \rightarrow 0.$$

The proof is complete. \square

The approximation (4.14) could be accurate even for large, possibly unbounded values for α , *provided such values occur with a small probability*. For example, such that the perturbation $\alpha - \alpha_0$ is L^m -summable for $m \geq k$ sufficiently high. The essential requirement here is local Lipschitz continuity of the Fréchet derivative of the solution operator on the possibly unbounded set containing α_0 in the sense of (4.17). We give this result first for a nonlinear sufficiently smooth mapping F in the following Lemma.

Lemma 4.7. *Suppose X, Z are Banach spaces, $\emptyset \neq U \subseteq Z$ is open and $\alpha_0 \in U$. Let $F \in C^1(U, X)$ be a nonlinear map for which F' is Lipschitz continuous at α_0 in the following sense:*

$$(4.17) \quad \exists K > 0 : \quad \|F'(\tilde{\alpha}) - F'(\alpha_0)\|_{Z \rightarrow X} \leq K \|\tilde{\alpha} - \alpha_0\|_Z \quad \forall \tilde{\alpha} \in U .$$

Assume that $B := \|F'(\alpha_0)\|_{Z \rightarrow X} < \infty$ and let $\alpha \in L^0(\Omega; Z)$ be a random field satisfying $\mathbb{P}\{\alpha(\omega) \in U\} = 1$.

(1) *If $\alpha \in L^{2k}(\Omega, \mathbb{P}; Z)$ for some $k \in \mathbb{N}$, then*

$$(4.18) \quad \begin{aligned} & \|\mathcal{M}^k[F(\alpha) - F(\alpha_0)] - \mathcal{M}^k[F'(\alpha_0)(\alpha - \alpha_0)]\|_{X^{(k)}} \\ & \leq ((B + K/2)^k - B^k) \max_{m \in \{k+1, \dots, 2k\}} \|\alpha - \alpha_0\|_{L^m(\Omega, \mathbb{P}; Z)}^m . \end{aligned}$$

(2) *If $\alpha \in L^{k+1}(\Omega, \mathbb{P}; Z)$ then*

$$(4.19) \quad \begin{aligned} & \|\mathcal{M}^k[F(\alpha) - F(\alpha_0)] - \mathcal{M}^k[F'(\alpha_0)(\alpha - \alpha_0)]\|_{X^{(k)}} \\ & \leq \frac{k}{2} K B_*^{k-1} \|\alpha - \alpha_0\|_{L^{k+1}(\Omega, \mathbb{P}; Z)}^{k+1} . \end{aligned}$$

where $B_* := \sup_{\tilde{\alpha} \in U} \|F'(\tilde{\alpha})\|_{Z \rightarrow X} < \infty$.

Proof. (1) Let $r := \alpha - \alpha_0$ be a random increment such that $r(\omega) \in U - \alpha_0$ \mathbb{P} -almost sure. In particular large realizations $r(\omega)$ are allowed, if $U \subseteq Z$ is large. By Theorem 2.3 we have in this case

$$F(\alpha) - F(\alpha_0) = \int_0^1 F'(\alpha_0 + r\theta)r \, d\theta .$$

This yields

$$\begin{aligned} \mathcal{M}^k[F(\alpha) - F(\alpha_0)] &= \mathbb{E} \left[\left(\int_0^1 F'(\alpha_0 + r(\omega)\theta)r(\omega) \, d\theta \right)^{(k)} \right] \\ &= \mathbb{E} \left[\int_{[0,1]^k} \prod_{i=1}^k F'(\alpha_0 + r(\omega)\theta_i)r(\omega) \, d\theta_1 \dots d\theta_k \right] . \end{aligned}$$

Thus

$$(4.20) \quad \begin{aligned} & \mathcal{M}^k[F(\alpha) - F(\alpha_0)] - \mathcal{M}^k[F'(\alpha_0)r] \\ &= \mathbb{E} \left[\int_{[0,1]^k} \left\{ \prod_{i=1}^k F'(\alpha_0 + r(\omega)\theta_i)r(\omega) - \prod_{i=1}^k F'(\alpha_0)r(\omega) \right\} d\theta_1 \dots d\theta_k \right] . \end{aligned}$$

For any fixed $\omega \in \Omega$ and $\theta_1, \dots, \theta_k \in [0, 1]$ the term in the inner brackets equals to

$$\sum_{j=1}^k \left(\prod_{i=1}^{j-1} F'(\alpha_0 + r\theta_i)r \right) \left(F'(\alpha_0 + r\theta_j)r - F'(\alpha_0)r \right) \left(\prod_{i=j+1}^k F'(\alpha_0)r \right) =: \Delta(\omega) .$$

Recalling definition of K, B we find $\|F'(\tilde{\alpha})\|_{Z \rightarrow X} \leq B + K\|\tilde{\alpha} - \alpha_0\|_Z$ for any $\tilde{\alpha} \in U$. Thus we obtain by triangle inequality and by the definition (4.8) of the cross-norm

$$(4.21) \quad \|\Delta(\omega)\|_{X^{(k)}} \leq \|r(\omega)\|_X^k \sum_{j=1}^k \left(\prod_{i=1}^{j-1} (B + K\theta_i\|r(\omega)\|_Z) \right) \left(K\theta_j\|r(\omega)\|_Z \right) B^{k-j} .$$

Inserting into (4.20) and integrating over $\theta_1, \dots, \theta_k$ we conclude

(4.22)

$$\begin{aligned} \|\mathcal{M}^k [F(\alpha) - F(\alpha_0)] - \mathcal{M}^k [F'(\alpha_0)r]\|_{X^{(k)}} &\leq \mathbb{E} \left[\int_{[0,1]^k} \|\Delta(\omega)\|_{X^{(k)}} d\theta_1 \dots d\theta_k \right] \\ &\leq \mathbb{E} \left[\|r(\omega)\|_Z^k \sum_{j=1}^k \left(B + \frac{K}{2} \|r(\omega)\|_Z \right)^{j-1} \frac{K}{2} \|r(\omega)\|_Z B^{k-j} \right]. \end{aligned}$$

By the binomial theorem the right-hand side equals to

$$\begin{aligned} &= \mathbb{E} \left[\frac{K}{2} \|r(\omega)\|_Z^{k+1} \sum_{j=1}^k B^{k-j} \sum_{i=0}^{j-1} \binom{j-1}{i} \left(\frac{K}{2} \|r(\omega)\|_Z \right)^i B^{j-i-1} \right] \\ &= \sum_{j=1}^k \sum_{i=0}^{j-1} \binom{j-1}{i} \left(\frac{K}{2} \right)^{i+1} B^{k-i-1} \mathbb{E} [\|r(\omega)\|_Z^{k+i+1}]. \end{aligned}$$

In this expression only the binomial coefficient depends on j . Interchanging the summation order we find for any numbers $g_i \in \mathbb{R}$

$$\sum_{j=1}^k \sum_{i=0}^{j-1} \binom{j-1}{i} g_i = \sum_{j=0}^{k-1} \sum_{i=0}^j \binom{j}{i} g_i = \sum_{i=0}^{k-1} \sum_{j=i}^{k-1} \binom{j}{i} g_i = \sum_{i=0}^{k-1} \binom{k}{i+1} g_i.$$

Changing $i+1 \mapsto i$ we obtain

(4.23)

$$\|\mathcal{M}^k [F(\alpha) - F(\alpha_0)] - \mathcal{M}^k [F'(\alpha_0)r]\|_{X^{(k)}} \leq \sum_{i=1}^k \binom{k}{i} \left(\frac{K}{2} \right)^i B^{k-i} \|r\|_{L^{k+i}(\Omega, \mathbb{P}; Z)}^{k+i}.$$

The right-hand side is finite if $\alpha \in L^{2k}(\Omega, \mathbb{P}; Z)$ and admits the bound as in (4.18) by the binomial theorem.

(2) The assertion (4.19) follows analogously. If F' is uniformly bounded, then the estimate in (4.21) can be replaced by

$$\|\Delta(\omega)\|_{X^{(k)}} \leq kK\theta_j B_*^{k-1} \|r(\omega)\|_Z^{k+1}.$$

Inserting this in (4.22) yields (4.19). Note that the right-hand side in (4.19) is finite if $\alpha \in L^{k+1}(\Omega, \mathbb{P}; Z)$, which is a weaker assumption as $\alpha \in L^{2k}(\Omega, \mathbb{P}; Z)$ in (1). \square

The following Theorem follows directly from Theorem 3.2 and Lemma 4.7.

Theorem 4.8. *Suppose that the functional J and its derivatives satisfy the assumptions of Theorem 3.2.*

(1) *If $\mathbb{P}\{\|\alpha(\omega) - \alpha_0\|_Z \leq \delta\} = 1$ then*

$$\begin{aligned} (4.24) \quad &\|\mathcal{M}^k [S(\alpha) - u_0] - \mathcal{M}^k [S'(\alpha_0)(\alpha - \alpha_0)]\|_{X^{(k)}} \\ &\leq \delta^k ((\xi_0 + \delta K/2)^k - \xi_0^k) = \mathcal{O}(\delta^{k+1}). \end{aligned}$$

(2) *Suppose S' satisfies the local Lipschitz condition*

$$(4.25) \quad \|S'(\tilde{\alpha}) - S'(\alpha_0)\|_{Z \rightarrow X} \leq K \|\tilde{\alpha} - \alpha_0\|_Z \quad \forall \tilde{\alpha} \in U,$$

where $U \subseteq Z$ is an open (possibly large or unbounded) set. Assume that $\alpha \in L^0(\Omega, \mathbb{P}; Z)$ is a random field, such that $\mathbb{P}\{\alpha(\omega) \in U\} = 1$.

(a) If $\alpha \in L^{2k}(\Omega, \mathbb{P}; Z)$, then

$$(4.26) \quad \begin{aligned} & \|\mathcal{M}^k[S(\alpha) - u_0] - \mathcal{M}^k[S'(\alpha_0)(\alpha - \alpha_0)]\|_{X^{(k)}} \\ & \leq ((\xi_0 + K/2)^k - \xi_0^k) \max_{m \in \{k+1, \dots, 2k\}} \|\alpha - \alpha_0\|_{L^m(\Omega, \mathbb{P}; Z)}^m. \end{aligned}$$

(b) If $\alpha \in L^{k+1}(\Omega, \mathbb{P}; Z)$ and $\xi_* := \sup_{\tilde{\alpha} \in U} \|S'(\tilde{\alpha})\|_{Z \rightarrow X} < \infty$, then

$$(4.27) \quad \begin{aligned} & \|\mathcal{M}^k[S(\alpha) - u_0] - \mathcal{M}^k[S'(\alpha_0)(\alpha - \alpha_0)]\|_{X^{(k)}} \\ & \leq \frac{k}{2} K \xi_*^{k-1} \|\alpha - \alpha_0\|_{L^{k+1}(\Omega, \mathbb{P}; Z)}^{k+1}. \end{aligned}$$

5. SPARSE TENSOR GALERKIN DISCRETIZATION

5.1. Formulation of k -moment equation. We address efficient sparse tensor discretizations of the abstract first order k -th moment equation (4.15), that is

$$(5.1) \quad G_0^{(k)} Z^{(k)} = [-J'_\alpha(\alpha_0, u_0)]^{(k)} \mathcal{M}^k \alpha \quad \text{in } X^{(k)}.$$

We recall that the solution $Z^{(k)} \in X^{(k)}$ of (5.1) approximates $\mathcal{M}^k u$ under smallness assumptions on the fluctuation $\alpha - \alpha_0$.

Throughout, we assume that X, Y, Z are reflexive Banach spaces and that the sensitivity operator $G_0 \in \mathcal{L}(X, Y')$ is boundedly invertible. This implies bounded invertibility of $G_0^{(k)} \in \mathcal{L}(X^{(k)}, (Y^{(k)})')$. A necessary and sufficient condition for bounded invertibility of G_0 are the *inf-sup conditions*: there exists $\gamma > 0$ such that

$$(5.2) \quad \inf_{0 \neq v \in Y} \sup_{0 \neq w \in X} \frac{\langle v, G_0 w \rangle}{\|v\|_Y \|w\|_X} \geq \gamma, \quad \inf_{0 \neq w \in X} \sup_{0 \neq v \in Y} \frac{\langle v, G_0 w \rangle}{\|v\|_Y \|w\|_X} \geq \gamma.$$

Here and in what follows, $\langle \cdot, \cdot \rangle$ denotes the $Y \times Y'$ duality. To facilitate the discussion, we associate with the sensitivity operator G_0 the bilinear form

$$(5.3) \quad g_0(w, v) := \langle v, G_0 w \rangle : X \times Y \rightarrow \mathbb{R}.$$

Based on the general assumptions made so far, the form $g_0(\cdot, \cdot) : X \times Y \mapsto \mathbb{R}$ is continuous, and satisfies the inf-sup conditions (5.2). An immediate consequence of (5.2) is the following quantification of well-posedness of the k -th moment equation (4.15): with the stability constant $\gamma > 0$ from (5.2), it holds

$$(5.4) \quad \begin{aligned} & \inf_{0 \neq v \in Y^{(k)}} \sup_{0 \neq w \in X^{(k)}} \frac{Y^{(k)} \langle v, [G_0]^{(k)} w \rangle_{X^{(k)}}}{\|v\|_{Y^{(k)}} \|w\|_{X^{(k)}}} \geq \gamma^k, \\ & \inf_{0 \neq w \in X^{(k)}} \sup_{0 \neq v \in Y^{(k)}} \frac{Y^{(k)} \langle v, [G_0]^{(k)} w \rangle_{X^{(k)}}}{\|w\|_{X^{(k)}}} \geq \gamma^k. \end{aligned}$$

This shows bounded invertibility of the tensorized operator $G_0^{(k)}$ arising in the first order k -th moment equation (4.15), however with *condition which deteriorates exponentially with respect to the moment order k* . We next address efficient discretization of (4.15). To keep technicalities and exposition simple, we shall occasionally impose the (restrictive) assumption of coercivity of G_0 :

$$(5.5) \quad X = Y \quad \text{and} \quad \exists \gamma > 0 : \forall v \in X : g_0(v, v) \geq \gamma \|v\|_X^2.$$

Clearly, (5.5) implies (5.2) and (5.4). Note that $X = Y$ does not imply $G_0 = G_0^*$.

5.2. Abstract Multilevel Galerkin Discretization. We present an abstract Galerkin discretization of (5.1). We start with the case $k = 1$, and assume the stability condition (5.2). The Galerkin discretization of (5.1) for $k = 1$ is based on two dense, nested sequences

$$(5.6) \quad \{V_\ell^X\}_{\ell=0}^\infty \subset X, \quad \{V_\ell^Y\}_{\ell=0}^\infty \subset Y$$

of equal and finite dimension, i.e. $N_\ell = \dim V_\ell^X = \dim V_\ell^Y < \infty$. In the elliptic case (5.5), we have $V_\ell^X = V_\ell^Y$ and we write $V_\ell = V_\ell^X = V_\ell^Y$.

We assume the *discrete stability condition*: there exists $\bar{\gamma} > 0$ and $\ell_0 \geq 0$ such that for all $\ell \geq \ell_0$

$$(5.7) \quad \inf_{0 \neq v \in V_\ell^Y} \sup_{0 \neq w \in V_\ell^X} \frac{\langle v, G_0 w \rangle}{\|v\|_Y \|w\|_X} \geq \bar{\gamma}, \quad \inf_{0 \neq w \in V_\ell^X} \sup_{0 \neq v \in V_\ell^Y} \frac{\langle v, G_0 w \rangle}{\|v\|_Y \|w\|_X} \geq \bar{\gamma}.$$

Evidently, in the coercive case (5.5), the discrete stability conditions (5.7) hold with $\ell_0 = 0$ and with $\bar{\gamma} = \gamma$ for *any* selection of subspaces V_ℓ .

The following result on the stability and the quasioptimality of Galerkin approximations of (5.1) follows from the Lax-Milgram Lemma.

Proposition 5.1. *Assume (5.2), (5.7). Then, for any $L \geq \ell_0$, and any $k \geq 1$, the tensorized k -linear Galerkin equations: find*

$$Z_L^{(k)} \in [V_L^X]^{(k)} \quad \forall v \in [V_L^Y]^{(k)} \quad \langle v, G_0^{(k)} Z_L^{(k)} \rangle = (-1)^k \langle v, [J'_\alpha(\alpha_0, u_0)]^{(k)} \mathcal{M}^k \alpha \rangle$$

admit a unique solution $Z_L^{(k)} \in [V_L^X]^{(k)}$ which converges quasioptimally to the unique solution $Z^{(k)} \in X^{(k)}$ of the first order, k -th moment equation (5.1). There holds

$$(5.8) \quad \|Z^{(k)} - Z_L^{(k)}\|_{X^{(k)}} \leq \left(1 + \frac{\|G_0\|^{(k)}}{\bar{\gamma}^k}\right) \inf_{w_L \in [V_L^X]^{(k)}} \|Z^{(k)} - w_L\|_{X^{(k)}}.$$

Convergence rates can then be obtained in the usual fashion, once *regularity* of $Z^{(k)}$ in (5.1) available. To state this in an abstract fashion, we embed X and Y' into two scales $\{X^s\}_{s \geq 0}$ and $\{Y'^s\}_{s \geq 0}$ of smoothness spaces of Besov type such that

$$(5.9) \quad X = X^0 \supset X^1 \supset X^2 \supset \dots \supset X^s \supset \dots, \quad Y' = Y'^0 \supset Y'^1 \supset Y'^2 \supset \dots \supset Y'^s \supset \dots.$$

Following [21, 24] we assume that the subspaces V_ℓ^X admit the *approximation property*: for $u \in X^s$ and for every $\ell \geq 0$ holds

$$(5.10) \quad \min_{w_\ell \in V_\ell^X} \|u - w_\ell\|_X \lesssim \Phi(N_\ell, s) \|u\|_{X^s}$$

where $\Phi(N, s)$ is a function tending monotonically to zero for every $s > 0$ (typically, $\Phi(N, s) = N^{-s/d}$ where $s > 0$ denotes a Sobolev resp. Besov smoothness order and $d \geq 1$ denotes the dimension of the computational domain in which the operator G_0 is defined).

Based on (5.10) and on the quasioptimality (5.8), convergence rates for the approximation of $Z^{(k)}$ (and, by Theorem 4.6), also for $\mathcal{M}^k[u - u_0]$.

To state the estimates, based on the smoothness scales (5.9), we introduce for $k \geq 2$ two families of tensorized smoothness scales: the “isotropic” scale $[X^s]_{iso}^{(k)}$ and the “mixed” scale $[X^s]_{mix}^{(k)}$ as follows:

$$(5.11) \quad \begin{aligned} [X^s]_{iso}^{(k)} &:= X^s \otimes X^{(k-1)} \cap X \otimes X^s \otimes X^{(k-2)} \cap \dots \cap X^{(k-1)} \otimes X^s, \\ [X^s]_{mix}^{(k)} &:= X^s \otimes \dots (k - \text{times}) \dots \otimes X^s. \end{aligned}$$

Then, for $\ell \geq \ell_0 \geq 0$, a tensor product argument implies the convergence rate

$$(5.12) \quad \|Z^{(k)} - Z_L^{(k)}\|_{X^{(k)}} \leq C^k \Phi(N_L, s) \|Z^{(k)}\|_{[X^s]_{iso}^{(k)}}$$

where $\dim[V_L^X]^{(k)} = N_L^k$, i.e. the full tensor product approximation of $Z^{(k)}$ converges at the same rate for all $k \geq 1$ (in particular, therefore, at the rate for the mean field problem $k = 1$). However, the number of degrees of freedom used in the full tensor approximation increases superlinearly with respect to N_L for second and higher order moments. In addition, the convergence estimates (5.12) require only the ‘‘isotropic’’ regularity of order s of $Z^{(k)}$. However, the tensor structure of the operator $G_0^{(k)}$ implies, in fact, regularity of order $s > 0$ for $Z^{(k)}$ in the mixed smoothness scales $[X^s]_{mix}^{(k)}$.

Proposition 5.2. *Assume (5.2) and that $\Gamma_0 := G_0^{-1} \in \mathcal{L}(Y^{ts}, X^s)$ for some $s > 0$. Then $\Gamma_0^{(k)} = [G_0^{(k)}]^{-1} \in \mathcal{L}([Y^{ts}]_{mix}^{(k)}, [X^s]_{mix}^{(k)})$ boundedly.*

The mixed regularity $Z^{(k)} \in [X^s]_{mix}^{(k)}$ is well known to allow for *sparse tensor approximation* at essentially the rate $\Phi(N_L, s)$, however, from sparse tensor subspaces $\widehat{V}_L^X(k) \subset [V_L^X]^{(k)}$ with substantially fewer degrees of freedom than $\dim([V_L^X]^{(k)}) = N_L^k$. Denoting for $\ell \geq 0$ by

$$W_\ell^X = V_\ell^X \setminus V_{\ell-1}^X, \quad W_\ell^Y = V_\ell^Y \setminus V_{\ell-1}^Y,$$

(with the convention that $V_{-1}^X = V_{-1}^Y = \emptyset$), the sparse tensor spaces $\widehat{V}_L^X(k)$, $\widehat{V}_L^Y(k)$ are

$$(5.13) \quad \widehat{V}_L^X(k) := \bigoplus_{\ell_1 + \dots + \ell_k \leq L} \bigotimes_{j=1}^k W_{\ell_j}^X, \quad \widehat{V}_L^Y(k) := \bigoplus_{\ell_1 + \dots + \ell_k \leq L} \bigotimes_{j=1}^k W_{\ell_j}^Y.$$

If $N_\ell = \dim(V_\ell^X) = \mathcal{O}(b^\ell)$ for some basis $b > 1$ (in the context of multilevel or wavelet methods considered below, $b = 2^d$), it is easily verified that, as $L \rightarrow \infty$,

$$(5.14) \quad \dim(\widehat{V}_L^X(k)) = \dim(\widehat{V}_L^Y(k)) = \mathcal{O}(N_L (\log(N_L))^{k-1}).$$

To harness the (generically available) additional regularity $Z^{(k)} \in [X^s]_{mix}^{(k)}$ of the solution of (5.1) and the superior approximation power of the sparse tensor product spaces (5.13) in (nonadaptive) Galerkin approximation schemes, however, the (fundamental) problem arises in the case $X \neq Y$ that the stability (5.7) in general does not appear to imply a corresponding *discrete inf-sup condition on the sparse tensor product spaces* $\widehat{V}_L^X(k)$, $\widehat{V}_L^Y(k)$. This stability problem does not appear in the coercive case (5.5): there, coercivity and conformity of the Galerkin discretizations are well-known to imply stability.

Theorem 5.3. *Assume (5.2) and that $G_0^{-1} \in \mathcal{L}(Y^{ts}, X^s)$ for some $s > 0$. Assume moreover that for $k > 1$ there exists $L_0 > 0$ such that for all $L \geq L_0$ the tensorized sensitivity operator $G_0^{(k)}$ is stable on the pairs of sparse tensor product spaces $\widehat{V}_L^X(k)$,*

$\widehat{V}_L^Y(k)$, i.e. that there exists $\bar{\gamma}(k) > 0$ such that

$$(5.15) \quad \inf_{0 \neq v \in \widehat{V}_L^Y(k)} \sup_{0 \neq w \in \widehat{V}_L^X(k)} \frac{\langle v, G_0^{(k)} w \rangle}{\|v\|_{Y^{(k)}} \|w\|_{X^{(k)}}} \geq \bar{\gamma}(k),$$

$$\inf_{0 \neq w \in \widehat{V}_L^X(k)} \sup_{0 \neq v \in \widehat{V}_L^Y(k)} \frac{\langle v, G_0^{(k)} w \rangle}{\|v\|_{Y^{(k)}} \|w\|_{X^{(k)}}} \geq \bar{\gamma}(k).$$

Then the sparse tensor Galerkin approximations: find $\widehat{Z}_L^{(k)} \in \widehat{V}_L^X(k)$ such that for all $v \in \widehat{V}_L^Y(k)$ holds

$$\langle v, G_0^{(k)} \widehat{Z}_L^{(k)} \rangle = (-1)^k \langle v, [J'_\alpha(\alpha_0, u_0)]^{(k)} \mathcal{M}^k \alpha \rangle$$

admits a unique solution $\widehat{Z}_L^{(k)} \in \widehat{V}_L^X(k)$. This solution converges quasioptimally, i.e.

$$\begin{aligned} \left\| Z^{(k)} - \widehat{Z}_L^{(k)} \right\|_{X^{(k)}} &\lesssim (\bar{\gamma}(k))^{-1} \inf_{w \in \widehat{V}_L^X(k)} \left\| Z^{(k)} - w \right\|_{X^{(k)}} \\ &\lesssim (\bar{\gamma}(k))^{-1} (\log N_L)^{\frac{k-1}{2}} \Phi(N_L, s) \left\| Z^{(k)} \right\|_{[X^s]_{miz}^{(k)}} \end{aligned}$$

with the number of degrees of freedom bounded by (5.14).

In the remainder of this paper, we illustrate the foregoing abstract theory with an example of the random solution to a nonstationary diffusion equation in a class of stationary, smooth and bounded (this could be weakened) *random domains* $D \subset \mathbb{R}^d$, $d \geq 2$.

6. NONSTATIONARY HEAT EQUATION IN RANDOM DOMAIN

A broad class of applications of the abstract theory which we developed in sections 2 - 5 consists of boundary value problems in random domains. Here, the dependence of solutions on the domain is generically nonlinear, even for linear partial differential equations. The sensitivity operator $G_0 = J'_\alpha(\alpha_0, u_0)$ is the *shape derivative of the random solution evaluated at the nominal domain*. The idea to use shape derivatives in first order second moment calculus was first proposed and developed in [12]. The rather well-developed shape calculus as presented e.g in [19] (see also [13]) immediately yields broad classes of applications. By the Hadamard formula (e.g. [19, Thm. 2.27]), *if the only source of randomness resp. uncertainty in the problem of interest is the shape of the domain D , the shape derivative (and, hence, all linearizations of the sensitivity operator G_0) takes the form of a linear boundary value problem without source terms in the nominal domain D_0* . This, and the necessity for sparse tensor spaces in the efficient Galerkin approximation of k -point correlation functions mandate *boundary integral equation reformulations and their Galerkin discretization by multiresolution schemes on the boundary ∂D_0 of the nominal domain*. Such multiresolution schemes have the additional advantage that they allow for *compression of the Galerkin discretization of G_0* which, in boundary integral formulations, is a nonlocal operator on ∂D_0 . For a stationary diffusion equation, this program was developed in [12]. Here, we extend this to time-dependent diffusion in a random domain. We consider here only *time independent random perturbations* of D_0 (and thereby avoid having to introduce stochastic

processes). We proceed along the lines of [12]: a) shape gradient and determination of G_0 , b) boundary reduction of the first order perturbation problem and c) sparse tensor Galerkin discretization of the boundary reduced problem.

6.1. Shape Calculus. As reference for the shape calculus, we follow [19, Chapter 3.4] and let $D \subset \mathbb{R}^d$ ($d \geq 2$) denote a bounded domain with smooth boundary ∂D and with exterior unit normal vector field n , and with $T > 0$ a finite time horizon, $I = (0, T)$ the time interval of interest, and denote by $Q = D \times I$ the space-time cylinder. Of particular relevance in the following will be the ‘‘mantle’’ $\Sigma = \partial D \times I$ of Q . In Q , we consider the linear, diffusion initial-boundary value problem with either Dirichlet- or Neumann boundary conditions: given $f \in L^2(Q)$ and $y_0 \in L^2(D)$, find $y = y(x, t)$ such that

$$(6.1) \quad \begin{aligned} \frac{\partial y}{\partial t} - \Delta y &= f & \text{in } Q, \\ \gamma_1 y &= \frac{\partial y}{\partial n} = 0 & \text{on } \Sigma, \\ y(x, 0) &= y_0(x) & \text{in } D. \end{aligned}$$

Here and in what follows, γ_1 denotes the (co)normal derivative on the boundary of the nominal domain, ∂D .

The weak form of (6.1) is as follows: find $y(t) \in H^1(D)$ such that for all $v \in H^1(D)$ holds

$$(6.2) \quad \begin{aligned} \int_D \left(v \frac{\partial y}{\partial t} + \nabla y \cdot \nabla v \right) dx &= \int_D f v dx & \text{a.e. } t \in I, \\ y(0) &= y_0 & \text{in } L^2(D). \end{aligned}$$

It is well-known that (6.2) admits a unique weak solution $y \in W(I; H^1(D))$ where, for any $H_0^1(D) \subseteq H \subseteq H^1(D)$, the Bochner space $W(I; H)$ is given by

$$W(I; H) = \left\{ \phi \in L^2(I; H) \left| \frac{\partial \phi}{\partial t} \in L^2(I; H') \right. \right\}.$$

Moreover, $W(I; H) \subset C^0(\bar{I}; L^2(D))$ with continuous injection so that the initial condition in (6.2) is well-defined.

Analogously, the *Dirichlet problem* reads: given $f \in L^2(Q)$ and $w_0 \in L^2(D)$, find $w = w(x, t)$ such that

$$(6.3) \quad \begin{aligned} \frac{\partial w}{\partial t} - \Delta w &= f & \text{in } Q, \\ \gamma_0 w &= w|_{\Sigma} = 0 & \text{on } \Sigma, \\ w(x, 0) &= w_0(x) & \text{in } D. \end{aligned}$$

In weak form: find $w(t) \in H_0^1(D)$ such that for all $v \in H_0^1(D)$ holds

$$(6.4) \quad \begin{aligned} \int_D \left(v \frac{\partial w}{\partial t} + \nabla w \cdot \nabla v \right) dx &= \int_D f v dx & \text{a.e. } t \in I, \\ w(0) &= w_0 & \text{in } L^2(D). \end{aligned}$$

Then (6.4) admits a unique weak solution $y \in W(I; H_0^1(D))$.

To compute the shape derivatives of y and w , we use the speed method: we transport the nominal domain D under a smooth flow and assume that the (deterministic) data f and y_0 are given in all of \mathbb{R}^d (rendering the diffusion problem meaningful on the perturbed domains). Specifically, for $\varepsilon > 0$ sufficiently small, given a velocity field $V \in C^\infty(\mathbb{R}^d; \mathbb{R}^d)$ and the one parameter family $D_s = T_s(V)(D) := \{T_s(V)(x) | x \in D\}$ of transported domains under the flow $T_s(V)$. For $\varepsilon > 0$ sufficiently small, D_s is smoothly diffeomorphic to $D = D_0$ under $T_s(V)$ for $s \in [0, \varepsilon)$.

As shown in [19, Lemma 3.7], the shape derivative y' of the solution y to the Neumann problem (6.1) with domain perturbation in direction V is the solution of an instationary Neumann problem. With γ_1 denoting the (distributional) conormal derivative on Σ , it reads: find y' such that

$$(6.5) \quad \begin{aligned} \frac{\partial y'}{\partial t} - \Delta y' &= 0 \quad \text{in } Q, \\ \gamma_1 y' &= y'_\Sigma [V \cdot n] := \gamma_0 \left(-\frac{\partial y}{\partial t} + f \right) (V \cdot n) + \operatorname{div}_{\partial D} ((V \cdot n) \nabla_{\partial D} \gamma_0 y) \quad \text{on } \Sigma, \\ y'(x, 0) &= 0 \quad \text{in } D. \end{aligned}$$

Using [19, Proposition 2.68], we obtain the alternative representation

$$(6.6) \quad y'_\Sigma [V \cdot n] = \nabla_{\partial D} (V \cdot n) \cdot \nabla_{\partial D} \gamma_0 u - (V \cdot n) \frac{\partial^2 y}{\partial n^2} \quad \text{on } \Sigma.$$

We remark that even for $\partial D \in C^\infty$ and smooth f and y_0 , this Cauchy data develops a singularity at $t = 0$, unless f and y_0 satisfy certain compatibility conditions at $\{t = 0\} \cap \partial D$.

Analogously, the shape derivative w' of the solution w of the Dirichlet problem (6.3) solves the following nonhomogeneous, instationary Dirichlet problem (see [19, Lemma 3.9]):

$$(6.7) \quad \begin{aligned} \frac{\partial w'}{\partial t} - \Delta w' &= 0 \quad \text{in } Q, \\ \gamma_0 w' &= w'_\Sigma [V \cdot n] := -(V \cdot n) \gamma_1 w \quad \text{on } \Sigma, \\ w'(x, 0) &= 0 \quad \text{in } D. \end{aligned}$$

As already observed by J. Hadamard [10], the Cauchy data y'_Σ and w'_Σ for both shape derivatives, (6.5) and (6.7), *depends linearly only on the normal component* $\alpha = V \cdot n$ *of the perturbation field*. Accordingly, we assume in what follows that random domain perturbations are defined by a speed field V of the form $V = \alpha(x, \omega)n$ where n denotes the exterior unit normal to the smooth, nominal domain $D_0 = D$, and where the scalar function α is a smooth random field on ∂D with mean zero and known two-point correlation function $\mathcal{M}^2[\alpha] = \mathbb{E}[\alpha(x, \cdot) \otimes \alpha(x', \cdot)] : \partial D \times \partial D \mapsto \mathbb{R}$. We also observe that both the Dirichlet and Neumann shape derivative problems (6.5) and (6.7) have homogeneous source terms and initial conditions.

6.2. Boundary Reduction. To develop optimal complexity Galerkin discretizations for the k -point correlation functions, we reduce (6.5) and (6.7) in the space time cylinder Q to first kind variational boundary integrodifferential equations on the

lower dimensional cylinder ‘mantle’ Σ , following [7] (see also [1] and [4]). Denote

$$(6.8) \quad E(t, x) := (4\pi t)^{-d/2} \exp(-|x|^2/4t) \vartheta(t)$$

the fundamental solution of the heat operator. Here $\vartheta(t) = \frac{1}{2}(1 + \text{sign}(t))$ denotes the Heaviside function. By κ_{t_0} we denote the time-reversal map given by $\kappa_{t_0} v(t, x) := v(t_0 - t, x)$. Then, for solutions $u(x, t)$ of the homogeneous heat equation there holds the *representation formula*

$$(6.9) \quad u = K_0(\gamma_1 u) - K_1(\gamma_0 u) \quad \text{in } Q,$$

where, for $(t_0, x_0) \in Q$ the *single layer heat potential* K_0 and the *double layer heat potential* K_1 is defined by

$$K_0(\varphi)(t_0, x_0) := \langle \varphi, \gamma_0 \kappa_{t_0} v \rangle, \quad K_1(w)(t_0, x_0) := \langle \gamma_1 \kappa_{t_0} v, w \rangle$$

with $v(t, x) = E(t, x_0 - x)$. For the formulation of the variational boundary integral equations, certain anisotropic Sobolev spaces on $\Sigma = \partial D_0 \times I$ are required: for $r \geq 0$, we denote by

$$(6.10) \quad \begin{aligned} \tilde{H}^{r, r/2}(\Sigma) &:= L^2(I; H^r(\partial D)) \cap \tilde{H}^{r/2}(I; L^2(\partial D)) \\ &\simeq L^2(I) \otimes H^r(\partial D) \cap \tilde{H}^{r/2}(I) \otimes L^2(\partial D) \end{aligned}$$

where \otimes denotes the usual tensor product of separable Hilbert spaces and where $\tilde{H}^r(I) = \{v \in H^r(I) | \tilde{v} \in H^r(-\infty, T)\}$ with \tilde{v} denoting the extension of v by zero to the negative real axis. Note that for $r < 1/2$ holds $\tilde{H}^r(I) \simeq H^r(I)$. The space $\tilde{H}^{r, r/2}(Q)$ is defined analogously. To formulate the boundary integral equations, we introduce for $(x, t) \in \Sigma$ the weakly singular and the hypersingular boundary integral operators:

$$(6.11) \quad \begin{aligned} (S\sigma)(x, t) &:= \int_0^t \int_{\partial D} \sigma(y, \tau) (\gamma_{0, y} E)(x - y, t - \tau) ds_y d\tau, \\ (H\mu)(x, t) &:= -\gamma_{1, x} \int_0^t \int_{\partial D} \mu(y, \tau) (\gamma_{1, y} E)(x - y, t - \tau) ds_y d\tau. \end{aligned}$$

Here, $\gamma_{0, y}$ denotes the trace operator taken with respect to the spatial variable y , and $\gamma_{1, x}$ denotes the (co)-normal derivative operator taken with respect to the spatial variable x . It is by now classical that the operators S and H in (6.11) are continuous and boundedly invertible in anisotropic Sobolev spaces on $\Sigma = \partial D_0 \times I$ [7, Theorem 3.7, Theorem 4.16]. Specifically, it is shown there that the following operators are bijective for every $|s| < 1/2$:

$$(6.12) \quad \begin{aligned} S &: \tilde{H}^{-1/2+s, (-1/2+s)/2}(\Sigma) \rightarrow \tilde{H}^{1/2+s, (1/2+s)/2}(\Sigma), \\ H &: \tilde{H}^{1/2+s, (1/2+s)/2}(\Sigma) \rightarrow \tilde{H}^{-1/2+s, (-1/2+s)/2}(\Sigma). \end{aligned}$$

The Sobolev range of bijectivity $|s| < 1/2$ can be enlarged to arbitrary $s > 0$ for smooth $\partial D \in C^\infty$, according to [7, Proposition 4.3]. The boundary reduction of the shape derivative problems (6.7), (6.5) is now straightforward (see [7, Corollary 3.16c), 3.17d)]): the unique variational solution w' of the Dirichlet problem (6.7) can be represented as

$$(6.13) \quad w' = K_0 \psi, \quad \text{where } S\psi = w'_\Sigma[V \cdot n],$$

and the unique solution of the Neumann problem as

$$(6.14) \quad y' = K_1 \zeta, \quad \text{where } H\zeta = -y'_\Sigma[V \cdot n]$$

with w'_Σ and y'_Σ as in (6.7), (6.5) respectively. The unique solvability of the boundary integral equations (6.13) and (6.14) is ensured by the following coercivity properties of the weakly singular boundary integral operator S and of the hypersingular boundary integral operator H (see [7, Corollary 3.13]): there exists a constant $\gamma > 0$ such that

$$(6.15) \quad \forall \psi \in H^{-1/2, -1/4}(\Sigma) : \quad \langle \psi, S\psi \rangle \geq \gamma \|\psi\|_{H^{-1/2, -1/4}(\Sigma)}^2,$$

and, with $\langle \cdot, \cdot \rangle$ denoting the corresponding duality pairings,

$$(6.16) \quad \forall \zeta \in H^{1/2, 1/4}(\Sigma) : \quad \langle \zeta, H\zeta \rangle \geq \gamma \|\zeta\|_{H^{1/2, 1/4}(\Sigma)}^2.$$

For a given, smooth velocity field $V \in C^\infty(\partial D_0; \mathbb{R}^3)$, the unique solution w' of the Dirichlet problem (6.3) is given by $w' = K_0\psi$ where the unknown surface density ψ is the solution of the *variational first kind boundary integral equation*: find

$$(6.17) \quad \psi \in H^{-1/2, -1/4}(\Sigma) : \quad \langle \bar{\psi}, S\psi \rangle = \langle \bar{\psi}, w'_\Sigma[V \cdot n] \rangle \quad \forall \bar{\psi} \in H^{-1/2, -1/4}(\Sigma).$$

By (6.15), the BIE (6.17) admits a unique solution for every $w'_\Sigma[V \cdot n] \in H^{1/2, 1/4}(\Sigma)$. Moreover, by [7] if $w'_\Sigma[V \cdot n] \in H^{1/2+s/2, 1/4+s/4}(\Sigma)$ with $s > 0$ this solution belongs to $H^{-1/2+s/2, -1/4+s/4}(\Sigma)$.

Analogously, the variational formulation of (6.14) reads: find

$$(6.18) \quad \zeta \in H^{1/2, 1/4}(\Sigma) : \quad \langle \bar{\zeta}, H\zeta \rangle = -\langle \bar{\zeta}, y'_\Sigma[V \cdot n] \rangle \quad \forall \bar{\zeta} \in H^{1/2, 1/4}(\Sigma).$$

By (6.16), for every $y'_\Sigma[V \cdot n] \in H^{-1/2, -1/4}(\Sigma)$, the BIE (6.18) admits a unique solution $\zeta \in H^{1/2, 1/4}(\Sigma)$ and if $y'_\Sigma[V \cdot n] \in H^{-1/2+s/2, -1/4+s/4}(\Sigma)$ for $s > 0$ we have $\zeta \in H^{1/2+s/2, 1/4+s/4}(\Sigma)$.

6.3. Sparse space-time Galerkin discretization. The coercivity (6.15) and (6.16) of the first kind boundary integral operators S and H implies stability and quasioptimality of Galerkin discretizations of the BIEs (6.17), (6.18) (e.g. [18]) for *any closed subspace* V_h of $V = H^{\pm 1/2, \pm 1/4}(\Sigma)$. For brevity of exposition, we develop the sparse tensor discretization only for the Galerkin BEM for the BIE (6.18) of the Neumann problem (6.5). Analogous results (and proof) hold for BIE (6.17) for the Dirichlet problem (6.7). To connect to the abstract Galerkin discretization framework in Section 5.2, we note that here $X = Y = V = H^{1/2, 1/4}(\Sigma)$ and, due to the ellipticity (6.16), for any closed subspace $V_h \subset H^{1/2, 1/4}(\Sigma)$, the Galerkin equations

$$(6.19) \quad \zeta_h \in V_h : \quad \langle \bar{\zeta}_h, H\zeta_h \rangle = -\langle \bar{\zeta}_h, y'_\Sigma[V \cdot n] \rangle \quad \forall \bar{\zeta}_h \in V_h.$$

are uniquely solvable and the Galerkin solutions $\zeta_h \in V_h$ are quasioptimal, i.e.

$$(6.20) \quad \|\zeta - \zeta_h\|_{H^{1/2, 1/4}(\Sigma)} \leq \gamma^{-1} \inf_{v_h \in V_h} \|\zeta - v_h\|_{H^{1/2, 1/4}(\Sigma)}.$$

Key to efficient Galerkin discretizations is therefore the proper choice of V_h . Owing to the (Cartesian) product structure of $\partial D \times I$, finite dimensional subspaces V_h can be constructed from *tensor products* of hierarchic multilevel subspaces of piecewise polynomials of degree $p \geq 1$ which are continuous in case of $V = H^{1/2, 1/4}(\Sigma)$ and of degree $q \geq 0$ which are possibly discontinuous in case of $V = H^{-1/2, -1/4}(\Sigma)$. Our generic notation for these spaces is as follows:

$$V_0^x \subset V_1^x \subset \dots \subset V_\ell^x \subset L^2(\partial D), \quad V_0^t \subset V_1^t \subset \dots \subset V_\ell^t \subset L^2(I)$$

Here, ℓ denotes ‘level of mesh refinement’, and we have in mind piecewise polynomial functions on uniformly refined meshes of widths $2^{-\ell_x}$ and $2^{-\ell_t}$, respectively.

Starting point for sparse space-time tensor Galerkin discretizations of the BIEs (6.17) and in (6.18) are multilevel decompositions of the hierarchies $\{V_\ell^x\}_{\ell=0}^\infty$ and $\{V_\ell^t\}_{\ell=0}^\infty$, respectively: for every level $L \geq 1$, we have

$$(6.21) \quad V_L^x = W_0^x \oplus W_1^x \oplus \dots \oplus W_L^x, \quad V_L^t = W_0^t \oplus W_1^t \oplus \dots \oplus W_L^t$$

where we *assume that we are explicitly given bases for the “detail” spaces W_ℓ^x and W_ℓ^t* (such bases are explicitly available, for example, in terms of hierarchic Finite Element bases or in terms of spline-wavelet bases as e.g. developed in [4]).

We build the *family* $\{\hat{V}_\ell\}_{\ell=0}^\infty \subset H^{1/2,1/4}(\Sigma)$ of *sparse tensor subspaces* from the hierarchies (6.21) by

$$(6.22) \quad \hat{V}_L := \bigoplus_{\ell_x + \ell_t \leq L} W_{\ell_x}^x \otimes W_{\ell_t}^t \subset V = H^{1/2,1/4}(\Sigma), \quad L = 1, 2, \dots$$

The corresponding (unique, by (6.16)) Galerkin solution of (6.19) with the subspace \hat{V}_L defined in (6.22) will be denoted by $\hat{\zeta}_L$. By the quasioptimality (6.20) and our regularity assumptions, the rate of convergence will be determined by the consistency of the family $\{\hat{V}_\ell\}_{\ell \geq 1}$ for smooth ζ , i.e. we assume

$$(6.23) \quad \zeta \in H^{1/2+s/2, 1/4+s/4}(\partial D_0 \times I) \quad \text{for all } s \geq 0.$$

In order to bound the convergence rates, we use that the anisotropic spaces (6.10) are intersection spaces and, therefore, are equipped with the sum-norms

$$\|\zeta - \hat{\zeta}_L\|_{H^{1/2,1/4}(\Sigma)} \simeq \|\zeta - \hat{\zeta}_L\|_{H^{1/2,0}(\Sigma)} + \|\zeta - \hat{\zeta}_L\|_{H^{0,1/4}(\Sigma)}$$

where the spaces $H_{mix}^{s,t}(\Sigma)$ of “square integrable, mixed highest derivative” are as in e.g. [11] defined by the tensor products: $H_{mix}^{s,t}(\Sigma) := H^s(\partial D_0) \otimes H^t(I)$. We may therefore apply [11, Theorem 7.1] (with $\sigma = 1$, $q_1 = 1/2$, $q_2 = 0$, $r_1 = r_2 = p + 1$) to obtain that for $\zeta \in H_{mix}^{s_1, s_2}(\Sigma)$ there exists a (quasi)interpolant $\hat{\zeta}_L \in \hat{V}_L$ such that for every $1/2 < s_x \leq p + 1$, $1/4 < s_t \leq p + 1$ holds

$$\|\zeta - \hat{\zeta}_L\|_{H_{mix}^{1/2,0}(\Sigma)} \lesssim 2^{-L \min\{s_x - 1/2, s_t\}} \|\zeta\|_{H_{mix}^{s_x, s_t}(\Sigma)}$$

and

$$\|\zeta - \hat{\zeta}_L\|_{H_{mix}^{0,1/4}(\Sigma)} \lesssim 2^{-L \min\{s_x, s_t - 1/4\}} \|\zeta\|_{H_{mix}^{s_x, s_t}(\Sigma)}$$

which implies that for $1/2 < s_x \leq p + 1$, $1/4 < s_t \leq p + 1$

$$(6.24) \quad \|\zeta - \hat{\zeta}_L\|_{H^{1/2,1/4}(\Sigma)} \lesssim 2^{-L \min\{s_x - 1/2, s_t - 1/4\}} \|\zeta\|_{H_{mix}^{s_x, s_t}(\Sigma)},$$

while, for $d = 3$ ([11, Theorem 4.1]), as $L \rightarrow \infty$ we have

$$(6.25) \quad \hat{N}_L := \dim(\hat{V}_L) \lesssim 2^{L(d-1)},$$

i.e. the number of degrees of freedom of the sparse tensor product space \hat{V}_L on Σ scales, as $L \rightarrow \infty$, as $\mathcal{O}(h_x^{-(d-1)})$, i.e. of a boundary element spaces on ∂D_0 with uniform mesh of meshwidth h_x (equation (6.25) remains valid also for $d = 2$, with a logarithmic factor). We sum up these results in the (slightly conservative) asymptotic bounds for the Galerkin discretization error

$$(6.26) \quad \begin{aligned} \|\zeta - \hat{\zeta}_L\|_{H^{1/2,1/4}(\Sigma)} &\lesssim 2^{-L(p+1/2)} \|\zeta\|_{H_{mix}^{p+1, p+\frac{3}{4}}(\Sigma)} \\ &\lesssim \hat{N}_L^{-(p+1/2)/(d-1)} \|\zeta\|_{H^{3p+\frac{5}{2}, \frac{3}{2}p+\frac{5}{4}}(\Sigma)} \end{aligned}$$

which is, again, the convergence rate afforded by a Galerkin boundary element discretization of a hypersingular boundary integral equation on a domain $D_0 \subset \mathbb{R}^d$

by continuous, piecewise polynomial boundary elements of degree $p \geq 1$ on ∂D_0 . Given the rather high smoothness of the heat kernel $E(t, x)$ in (6.8), by choosing wavelet bases for the detail spaces W_ℓ^x and W_ℓ^t in the decompositions (6.21) which have a sufficiently high number of vanishing moments, the stiffness matrices of the boundary integral operators H (and of S) can be compressed to $\mathcal{O}(\hat{N}_L)$ many non-vanishing entries *without* compromising the convergence rate (6.26). In addition, with the wavelets being Riesz bases in the spaces $L^2(I)$ and $L^2(\partial D_0)$, respectively, and, properly scaled, also in $H^{1/4}(I)$ and in $H^{1/2}(\partial D_0)$, the matrices of the Galerkin discretized operator S and H have condition numbers which are uniformly bounded with respect to L , allowing for optimal complexity solvers. In particular, if

$$\{\phi_\lambda^x : \lambda \in \nabla_x\} \subset H^{1/2}(\partial D_0), \quad \{\phi_\mu^t : \mu \in \nabla_t\} \subset H^{1/4}(I)$$

are collections of functions being normalized Riesz bases for $L^2(\partial D_0)$ and $L^2(I)$ respectively, then the collection

$$(6.27) \quad \left\{ (x, t) \mapsto \frac{\phi_\lambda^x(x) \phi_\mu^t(t)}{\sqrt{\|\phi_\lambda^x\|_{H^{1/2}(\partial D_0)}^2 + \|\phi_\mu^t\|_{H^{1/4}(I)}^2}} : (\lambda, \mu) \in \nabla_x \times \nabla_t \right\}$$

is a Riesz basis for $H^{1/2, 1/4}(\Sigma)$ (see, e.g. [9, 25, 23] for details).

Naturally, $\hat{\zeta}_L$ is only an approximate density on Σ . Information on the solution inside the space-time cylinder Q must be extracted by inserting $\hat{\zeta}_L$ into the representation formula (6.14); this, in fact, allows for *superconvergent approximations* of $u(x, t)$ as follows by an Aubin-Nitsche duality type argument for the Galerkin BIE (6.18). (see [22] for details on this in the elliptic setting).

6.4. Sparse space-time Galerkin discretization for the k -th moment equation. Tensorization of (6.13), (6.14) yields

$$(6.28) \quad \mathcal{M}^k[w'] = K_0^{(k)} \mathcal{M}^k[\psi], \quad \text{where } S^{(k)} \mathcal{M}^k[\psi] = (w'_\Sigma)^{(k)} \mathcal{M}^k[V \cdot n],$$

and

$$(6.29) \quad \mathcal{M}^k[y'] = K_1^{(k)} \mathcal{M}^k[\zeta], \quad \text{where } H^{(k)} \mathcal{M}^k[\zeta] = (-y'_\Sigma)^{(k)} \mathcal{M}^k[V \cdot n].$$

As discussed in Section 5, the coercivity (6.15), (6.16) of the boundary integral operators S and H implies well-posedness of (6.28) and (6.29) and stability and quasioptimality of Galerkin discretizations of the k -th moment BIE. We again work out the details for the Neumann problem (6.5) and (6.29). Analogous results also hold for (6.7) and (6.28).

Suppose \hat{V}_L is a family of sparse tensor subspaces defined in (6.22). We introduce

$$(6.30) \quad \widehat{V}_L^{(k)} := \bigoplus_{L_1 + \dots + L_k \leq L} \bigotimes_{j=1}^k \hat{W}_{L_j} \subset V^{(k)}, \quad V = H^{1/2, 1/4}(\Sigma)$$

with the ‘‘detail’’ spaces

$$(6.31) \quad \hat{W}_{L_j} := \hat{V}_{L_j} \setminus \hat{V}_{L_j-1} = \bigoplus_{\ell_x + \ell_t = L_j} W_{\ell_x}^x \otimes W_{\ell_t}^t.$$

Due to (6.16) the Galerkin equations

$$(6.32) \quad \zeta_L^{(k)} \in \widehat{V}_L^{(k)} : \langle v, H^{(k)} \zeta_L^{(k)} \rangle = (-1)^k \langle v, (y'_\Sigma)^{(k)} [V \cdot n] \rangle \quad \forall v \in \widehat{V}_L^{(k)}$$

are uniquely solvable and the Galerkin solutions $\zeta_L^{(k)} \in \widehat{V}_L^{(k)}$ are quasioptimal, i.e. for $d = 3$ (6.26) yields

$$(6.33) \quad \begin{aligned} \|\mathcal{M}^k[\zeta] - \zeta_L^{(k)}\|_{[H^{1/2,1/4}(\Sigma)]^{(k)}} &\leq \gamma^{-k} \inf_{v_L \in \widehat{V}_L^{(k)}} \|\mathcal{M}^k[\zeta] - v_L\|_{[H^{1/2,1/4}(\Sigma)]^{(k)}} \\ &\lesssim \gamma^{-k} L^{\frac{k-1}{2}} 2^{-L(p+1/2)} \|\mathcal{M}^k[\zeta]\|_{[H^{r,r/2}(\Sigma)]_{mix}^{(k)}} \end{aligned}$$

where $r = 3p + \frac{5}{2}$. As $L \rightarrow \infty$, the cardinality of $\widehat{V}_L^{(k)}$ scales as

$$\dim(\widehat{V}_L^{(k)}) \sim \log(\hat{N}_L)^{k-1} \hat{N}_L \lesssim L^{k-1} 2^{L(d-1)},$$

i.e. (up to a logarithmic factor) as the cardinality of the boundary element space on ∂D_0 with the uniform meshwidth $h_x \sim 2^{-L}$.

The approximation \mathcal{Y}'_L to $\mathcal{M}^k[y']$ is defined via the representation formula

$$(6.34) \quad \mathcal{Y}'_L := K_1^{(k)} \zeta_L^{(k)}.$$

The mapping $K_1 : H^{1/2,1/4}(\Sigma) \rightarrow \tilde{H}^{1,1/2}(Q)$ is continuous, thus

$$\|\mathcal{M}^k[y'] - \mathcal{Y}'_L\|_{[H^{1,1/2}(Q)]_{mix}^{(k)}} \leq \|K_1\|^k \|\mathcal{M}^k[\zeta] - \zeta_L^{(k)}\|_{[H^{1/2,1/4}(\Sigma)]^{(k)}}.$$

Finally, we utilize (4.24) from Theorem 4.8 and obtain by the triangle inequality the error estimate for the k -th moment:

$$(6.35) \quad \begin{aligned} \|\mathcal{M}^k[\frac{y_s - y_0}{s}] - \mathcal{Y}'_L\|_{[H^{1,1/2}(Q)]_{mix}^{(k)}} &\leq \|\mathcal{M}^k[\frac{y_s - y_0}{s}] - \mathcal{M}^k[y']\|_{[H^{1,1/2}(Q)]_{mix}^{(k)}} + \|\mathcal{M}^k[y'] - \mathcal{Y}'_L\|_{[H^{1,1/2}(Q)]_{mix}^{(k)}} \\ &\leq (\delta/s)^k ((\xi_0 + \delta K/2)^k - \xi_0^k) + C \|K_1\| L^{\frac{k-1}{2}} 2^{-L(p+1/2)} \|\mathcal{M}^k[\zeta]\|_{[H^{r,r/2}(\Sigma)]_{mix}^{(k)}} \end{aligned}$$

where y_s is the exact solution of (6.1) in the perturbed spatial domain $D_s = T_s(V)(D)$, see Subsection 6.1. The constants ξ_0 , K and δ were explicitly defined in Section 4 and depend on the size of the random boundary perturbation α .

Remark 6.1. *The bound (6.35) tells that the level L of the Finite Element discretization $\widehat{V}_L^{(k)}$ should balance the linearization error represented by the first term in the right-hand side of (6.35) $L = L^{opt}$. For $L > L^{opt}$ the linearization error dominates the total error in (6.35) and further refinement will not provide a better approximation of the k -th moment $\mathcal{M}^k[\frac{y_s - y_0}{s}]$.*

The linearization error depends on ξ_0 , K and δ . In the next subsection we exemplify the abstract linearization error bounds by giving explicit values of these constants for the parabolic model problem under consideration.

6.5. Bounds on the linearization error. We recall definitions (3.1), (3.19) and Theorem 4.8(1) of the constants ξ_0 , K and δ . For the parabolic model problem (6.1), the spaces X , Y , Z from Section 4 are given by

$$(6.36) \quad \begin{aligned} X &:= \tilde{H}^{1,\frac{1}{2}}(Q, \frac{\partial}{\partial t} - \Delta) := \left\{ u \in \tilde{H}^{1,\frac{1}{2}}(Q) \mid (\frac{\partial}{\partial t} - \Delta)u \in L^2(Q) \right\}, \\ Y &:= \tilde{H}^{1,\frac{1}{2}}(Q), \quad Z := C^k(\mathbb{R}^d) \end{aligned}$$

with $k \geq 1$ being sufficiently large. Note that $\{u \in W(I, H^1(D)) | u(x, 0) = 0\} \subset \tilde{H}^{1, \frac{1}{2}}(Q)$ by interpolation. In this setting we have

$$\xi_0 = \|\Gamma_0 J'_\alpha(\alpha_0, y_0)\|_{C^k(\mathbb{R}^d) \rightarrow H^{1, \frac{1}{2}}(Q)} = \sup_{\alpha \in C^k(\mathbb{R}^d)} \frac{\|y'_0(\alpha)\|_{H^{1, \frac{1}{2}}(Q)}}{\|\alpha\|_{C^k(\mathbb{R}^d)}}$$

and

$$\|y' - y'_0\|_{C^k(\mathbb{R}^d) \rightarrow H^{1, \frac{1}{2}}(Q)} \leq K \|\alpha - \alpha_0\|_{C^k(\mathbb{R}^d)}.$$

To avoid ambiguity we denote here by y'_0 the solution of (6.5) in the nominal domain $D = D_0$ corresponding to the zero perturbation $\alpha_0 = 0$ whereas y' the solution of (6.5) on the perturbed domain $D_s = T_s(V)(D)$ corresponding to the perturbation parameter α . First, we derive an explicit bound for ξ_0 . By continuity of K_1 and coercivity (6.16) of H we obtain

$$\|y'\|_{H^{1, 1/2}(Q)} \leq \|K_1\| \|\zeta\|_{H^{\frac{1}{2}, \frac{1}{4}}(\Sigma)} \leq \frac{\|K_1\|}{\gamma} \|y'_\Sigma[\alpha]\|_{H^{-\frac{1}{2}, -\frac{1}{4}}(\Sigma)}$$

and by (6.6)

$$\|y'_\Sigma[\alpha]\|_{H^{-\frac{1}{2}, -\frac{1}{4}}(\Sigma)} \leq \|\alpha\|_{W^{1, \infty}(\partial D)} \left(\|\nabla_{\partial D} \gamma_0 y\|_{H^{-\frac{1}{2}, -\frac{1}{4}}(\Sigma)} + \left\| \frac{\partial^2 y}{\partial n^2} \right\|_{H^{-\frac{1}{2}, -\frac{1}{4}}(\Sigma)} \right).$$

Thus

$$(6.37) \quad \xi_0 = \frac{\|K_1\|}{\gamma} \left(\|\nabla_{\partial D} \gamma_0 y\|_{H^{-\frac{1}{2}, -\frac{1}{4}}(\Sigma)} + \left\| \frac{\partial^2 y}{\partial n^2} \right\|_{H^{-\frac{1}{2}, -\frac{1}{4}}(\Sigma)} \right)$$

provided the nominal solution y is sufficiently regular.

Next, we identify the local Lipschitz constant K . If the second shape derivative of y exists, then

$$K = \sup_{\|\alpha\|_{C^k(\mathbb{R}^d)} \leq \delta} \|y''(\alpha)\|_{C^k(\mathbb{R}^d) \times C^k(\mathbb{R}^d) \rightarrow H^{1, \frac{1}{2}}(Q)}.$$

The second shape derivative of y'' can be identified as a solution of the following problem:

$$(6.38) \quad \begin{aligned} \frac{\partial y''}{\partial t} - \Delta y &= 0 && \text{in } Q_s \\ \frac{\partial y''}{\partial n} &= y''_\Sigma [V \cdot n, W \cdot n] && \text{on } \Sigma_s \\ y''(x, 0) &= 0 && \text{in } D_s \end{aligned}$$

where

$$(6.39) \quad \begin{aligned} y''_\Sigma[\alpha, \beta] &= -\alpha\beta \frac{\partial^3 y}{\partial n^3} + \nabla_{\partial D_s} \beta \cdot \nabla_{\partial D_s} y'[\alpha] - \beta \frac{\partial^2 y'[\alpha]}{\partial n_s^2} \\ &\quad + \nabla_{\partial D_s} \alpha \cdot \nabla_{\partial D_s} y'[\beta] - \alpha \frac{\partial^2 y'[\beta]}{\partial n_s^2}. \end{aligned}$$

Similarly as above, we obtain

$$(6.40) \quad \begin{aligned} K &= \frac{\|K_1\|}{\gamma} \left(\|\nabla_{\partial D_s} y'[\cdot]\|_{C^k(\mathbb{R}^d) \rightarrow H^{-\frac{1}{2}, -\frac{1}{4}}(\Sigma)} \right. \\ &\quad \left. + \left\| \frac{\partial^2 y'[\cdot]}{\partial n_s^2} \right\|_{C^k(\mathbb{R}^d) \rightarrow H^{-\frac{1}{2}, -\frac{1}{4}}(\Sigma)} + \left\| \frac{\partial^3 y}{\partial n_s^3} \right\|_{H^{-\frac{1}{2}, -\frac{1}{4}}(\Sigma)} \right). \end{aligned}$$

The value of k in (6.36) and (6.40) must be chosen so that the operator norms in the right-hand side of (6.40) remain bounded. By (6.14) this task reduces to the study of the mapping properties of non-classical boundary integral operators

$$\nabla_{\partial D} K_1, \quad \text{and} \quad \frac{\partial^2}{\partial n^2} K_1$$

which is beyond the scope of the present paper, see [7] for more details.

7. CONCLUDING REMARKS

In this paper we provide the first order k -th moment analysis for nonlinear operator equations with random data. In Section 3 we studied solvability of a general nonlinear parametric equation in Banach spaces for varying parameters and obtained *precise estimates on the size of the neighborhoods where the solution operator is well defined* (Theorem 3.1, Theorem 3.2). These results are new and provide a generalization of the classical Implicit Function Theorem.

In Section 4 this result has been applied to abstract nonlinear equations in Banach spaces with random parameters. We established precise *non-asymptotic error bounds* with *explicit constants* for the magnitude of the linearization error for the k -th moment equation (see Theorem 4.8 being a generalization of the asymptotic estimate in Theorem 4.6).

In Section 5 we introduced an abstract multilevel Galerkin discretization for the linearized k -th moment equations. As a result of Theorem 5.3, the *k -th moment equation in the k -fold product domain can be solved at essentially the same cost as a single linearized problem in the original non-tensorized domain.*

In Section 6 we applied the general methodology to model nonstationary non-homogeneous parabolic equations in randomly perturbed domains. By means of Shape Calculus for the underlying partial differential equations we obtain a formulation for the shape derivative of the solution which does not contain source terms. It can, therefore, be efficiently treated numerically via boundary reduction to strongly positive first kind boundary integral equations on the boundary of the space-time cylinder.

We introduce sparse tensor product Galerkin discretizations of these boundary integral equations. Sparsity enters here on two levels: first, the space and time discretizations are performed by sparse tensorization, thereby compressing the Galerkin approximation unknown surface density to a format with complexity equal to that of a boundary integral equation of a stationary problem.

Second, the hierarchic multilevel space-time tensor basis allows to exploit the mixed regularity of the k -point correlations functions. This leads to a priori error estimates for the resulting nonstationary k -th moment equation which equal, up to logarithmic terms, those for the mean field of a stationary diffusion problem on the boundary of the physical domain. In particular, we solve the second moment problem for a three-dimensional nonstationary heat conduction, being an *eight-dimensional problem*, with the same error versus number of unknowns as a stationary diffusion problem in a *two-dimensional* domain. Finally, we derive a bound for the total error (6.35) with explicit bounds for the linearization and the discretization errors.

We solved the problems on the boundary of the space-time cylinder using boundary reduction via first kind boundary integral equations. Their stability in parabolic trace spaces is a result of the coercivity of the underlying integral operators. An

alternative approach using *boundary integral equations of the second kind* is also possible.

REFERENCES

- [1] Douglas N. Arnold and Patrick J. Noon. Coercivity of the single layer heat potential. *J. Comput. Math.* **7** (1989) 100–104.
- [2] Antonio Ambrosetti and Giovanni Prodi. *A primer of nonlinear analysis*, volume 34 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1993.
- [3] Melvin S. Berger. *Nonlinearity and functional analysis*. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1977. Lectures on nonlinear problems in mathematical analysis, Pure and Applied Mathematics.
- [4] Ch. Bourgeois and S. Nicaise. Prewavelet Analysis of the heat equation. *Numer. Math.* **87** (2001) 407–434.
- [5] Alexey Chernov. Abstract sensitivity analysis for nonlinear equations and applications. In Karl Kunisch, Günther Of, and Olaf Steinbach, editors, *Numerical Mathematics and Advanced Applications*, Proceedings of ENUMATH 2007, Graz, Austria, pages 407–414. Springer, September 2008.
- [6] Alexey Chernov and Christoph Schwab. Sparse p -version BEM for first kind boundary integral equations with random loading. *Appl. Numer. Math.*, 59(11):2698–2712, 2009.
- [7] Martin Costabel. Boundary integral operators for the heat equation. *Int. Eqns. and Operator Theory* **13** (1990) 498–552.
- [8] Giuseppe Da Prato. *An introduction to infinite-dimensional analysis*. Universitext. Springer-Verlag, Berlin, 2006. Revised and extended from the 2001 original by Da Prato.
- [9] Michael Griebel and Peter Oswald. *Tensor product type subspace splittings and multilevel iterative methods for anisotropic problems*. *Adv. Comput. Math.* **4** (1995), no. 1-2, 171–206
- [10] Jacques Hadamard. *Lessons on the Calculus of Variations* (in french). (1910).
- [11] Michael Griebel and Helmut Harbrecht. *On the construction of sparse tensor product spaces*. INS Preprint 1104, Universität Bonn, 2011, submitted.
- [12] Helmut Harbrecht, Reinhold Schneider and Christoph Schwab. *Sparse second moment analysis for elliptic problems in stochastic domains*. *Numer. Math.* **109** (2008) 167–188.
- [13] R. Hiptmair and J. Li. *Shape derivatives in differential forms I: An intrinsic perspective*. SAM-Report 2011-42, ETH Zurich.
- [14] L. V. Kantorovich and G. P. Akilov. *Funktsionalnyi analiz (Russian) [Functional analysis]*. “Nauka”, Moscow, third edition, 1984.
- [15] A. N. Kolmogorov and S. V. Fomin. *Elementy teorii funktsii i funktsionalnogo analiza (Russian) [Elements of the theory of functions and functional analysis]*. FIZMATLIT, Moscow, seventh edition, 2004.
- [16] Michel Ledoux and Michel Talagrand. *Probability in Banach spaces*, volume 23 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1991. Isoperimetry and processes.
- [17] Claudia Prévôt and Michael Röckner. *A concise course on stochastic partial differential equations*, volume 1905 of *Lecture Notes in Mathematics*. Springer, Berlin, 2007.
- [18] Stefan Sauter and Christoph Schwab. *Boundary Element Methods*, Springer Ser. Comp. Math, Springer Verlag (2010).
- [19] Jan Sokolowski and Jean-Paul Zolesio. *Introduction to Shape Optimization*. Springer Ser. Computational Mathematics **16** Springer Verlag Berlin, Heidelberg, New York (1992).
- [20] O. Steinbach and W. L. Wendland. On C. Neumanns Method for Second-Order Elliptic Systems in Domains with Non-smooth Boundaries *J. Math. Anal. Appl.* **262** (2001), no. 2, 733–748.
- [21] Christoph Schwab and Radu Alexandru Todor. Sparse finite elements for stochastic elliptic problems— higher order moments. *Computing*, 71(1):43–63, 2003.
- [22] Christoph Schwab and Wolfgang Wendland. On the extraction technique in boundary integral equations. *Math. Comp.* **68** (1999) 91–122.
- [23] Christoph Schwab and Claude Jeffrey Gittelson. *Sparse tensor discretizations of high-dimensional parametric and stochastic PDEs* *Acta Numerica* **20** (2011), Cambridge University Press.

- [24] Christoph Schwab and Tobias von Petersdorff. Sparse Finite Element Methods for Operator Equations with Stochastic Data, *Applications of Mathematics* **51**(2) (2006) 145-180.
- [25] Christoph Schwab and Rob Stevenson, Space-Time adaptive wavelet methods for parabolic evolution equations, *Math. Comp.* **78** (2009), no. 267, 1293 - 1318.
- [26] Eberhard Zeidler. *Nonlinear functional analysis and its applications. I.* Springer-Verlag, New York, 1986. Fixed-point theorems, Translated from the German by Peter R. Wadsack.

HAUSDORFF CENTER FOR MATHEMATICS & INSTITUTE FOR NUMERICAL SIMULATION, UNIVERSITY OF BONN, 53115 BONN, GERMANY

E-mail address: `chernov@hcm.uni-bonn.de`

SEMINAR FOR APPLIED MATHEMATICS, ETH ZÜRICH, 8092 ZÜRICH, SWITZERLAND

E-mail address: `schwab@sam.math.ethz.ch`

Research Reports

No.	Authors/Title
11-51	<i>A. Chernov and Ch. Schwab</i> First order k -th moment finite element analysis of nonlinear operator equations with stochastic data
11-50	<i>E. Fonn</i> Shearlet Galerkin for transport equations: implementation and stability
11-49	<i>M.J. Castro, U.S. Fjordholm, S. Mishra and C. Parés</i> Entropy conservative and entropy stable schemes for non-conservative hyperbolic systems
11-48	<i>D. Kressner and Ch. Tobler</i> Preconditioned low-rank methods for high-dimensional elliptic PDE eigenvalue problems
11-47	<i>H. Heumann and R. Hiptmair</i> Convergence of lowest order semi-Lagrangian schemes
11-46	<i>K. Grella and Ch. Schwab</i> Sparse discrete ordinates method in radiative transfer
11-45	<i>X. Claeys and R. Hiptmair</i> Boundary integral formulation of the first kind for acoustic scattering by composite structures
11-44	<i>A. Chkifa, A. Cohen, R. DeVore and Ch. Schwab</i> Sparse adaptive Taylor approximation algorithms for parametric and stochastic elliptic PDEs
11-43	<i>S. Chen and S. Mao</i> Anisotropic error bounds of Lagrange interpolation with any order in two and three dimensions
11-42	<i>R. Hiptmair and J. Li</i> Shape derivatives in differential forms I: An intrinsic perspective
11-41	<i>Ph. Grohs and Ch. Schwab</i> Sparse twisted tensor frame discretization of parametric transport operators
11-40	<i>J. Li, H. Liu, H. Sun and J. Zou</i> Imaging acoustic obstacles by hypersingular point sources
11-39	<i>U.S. Fjordholm, S. Mishra and E. Tadmor</i> Arbitrarily high order accurate entropy stable essentially non-oscillatory schemes for systems of conservation laws