

# Energy conservative and -stable schemes for the two-layer shallow water equations

U.S. Fjordholm

Research Report No. 2010-41  
November 2010

Seminar für Angewandte Mathematik  
Eidgenössische Technische Hochschule  
CH-8092 Zürich  
Switzerland

# Energy conservative and -stable schemes for the two-layer shallow water equations.

Ulrik Skre Fjordholm  
*ETH Zurich, Rämistrasse 101, 8092 Zurich*  
*E-mail: ulrikf@sam.math.ethz.ch*

## Abstract

By extending the work in [4] we design an energy stable finite difference scheme for the two-layer shallow water equations.

## 1 Introduction

The two-layer shallow water equations model the flow of two fluids of different densities, superimposed on each other, under the influence of gravity. The main assumption in the derivation is that the horizontal length scales are much bigger than the vertical scales, and that one can therefore neglect variations in depth of density and velocity. Furthermore, it is assumed that no mixing occurs between the fluids. In one spatial dimension the equations have the form

$$\begin{aligned} (h_1)_t + (h_1 u_1)_x &= 0, & (h_2)_t + (h_2 u_2)_x &= 0, \\ (h_1 u_1)_t + \left( \frac{1}{2} g h_1^2 + h_1 u_1^2 \right)_x &= -g h_1 (b + r h_2)_x, & (1.1) \\ (h_2 u_2)_t + \left( \frac{1}{2} g h_2^2 + h_2 u_2^2 \right)_x &= -g h_2 (b + h_1)_x, \end{aligned}$$

where subscripts 1 and 2 denote the lower and upper layers, respectively,  $b = b(x)$  is the bottom topography,  $h$  is the layer height,  $u$  is layer velocity,  $r := \frac{\rho_2}{\rho_1} < 1$  and  $\rho$  is the layer density, with  $\rho_1 > \rho_2$ . In this equation, mass ( $h_1$  and  $h_2$ ) and total momentum ( $\rho_1 h_1 u_1 + \rho_2 h_2 u_2$ ) are the conserved variables. (1.1) is a *balance law*, which in general has the form

$$\mathbf{u}_t + \mathbf{f}(\mathbf{u}_x) = \mathbf{s}(\mathbf{u}, \mathbf{u}_x, x).$$

The system (1.1) is equipped with an entropy pair, the *energy* of the

solution:

$$\begin{aligned}\eta &= \frac{1}{2} (\rho_1 (h_1 u_1^2 + g h_1^2) + \rho_2 (h_2 u_2^2 + g h_2^2)) + g \rho_1 h_1 b + g \rho_2 h_2 (h_1 + b) \\ q &= \sum_{i=1}^2 \rho_i \left( \frac{1}{2} h_i u_i^3 + g h_i u_i (h_i + b) \right) + g \rho_2 h_1 h_2 (u_1 + u_2).\end{aligned}\tag{1.2}$$

The relevant entropy condition for (1.1) is therefore

$$\eta(\mathbf{u})_t + q(\mathbf{u})_x \leq 0.\tag{1.3}$$

The two-layer shallow water equations entails several difficult problems:

- The right-hand side products (e.g.  $h_1 \partial_x h_2$ ) are undefined at discontinuities, as we get the product of a distribution ( $h_1$ ) with a measure ( $\partial_x h_2$ ), a *nonconservative product*. The theoretical framework for nonconservative products was set by DalMaso, LeFloch and Murat in [3], and numerical methods (so-called path-conservative schemes) have been developed to fit into this framework. However, as has been recently reported [1], such schemes may not converge to the correct weak solution.
- As (1.1) is a 4-by-4 system, a direct calculation of its eigenvalues can be hard. However, a first-order approximation in  $u_2 - u_1$  was found in [2] to be

$$\begin{aligned}\lambda_{\text{int}}^{\pm} &= U_c \pm \sqrt{g' \frac{h_1 h_2}{h_1 + h_2} \left( 1 - \frac{(u_2 - u_1)^2}{g'(h_1 + h_2)} \right)}, \\ \lambda_{\text{ext}}^{\pm} &= U_m \pm \sqrt{g(h_1 + h_2)},\end{aligned}\tag{1.4}$$

where  $U_c := \frac{h_1 u_2 + h_2 u_1}{h_1 + h_2}$ ,  $U_m := \frac{h_1 u_1 + h_2 u_2}{h_1 + h_2}$ ,  $g' := g(1 - r)$ . From this we see that the system is only hyperbolic in the regime

$$\frac{(u_2 - u_1)^2}{g'(h_1 + h_2)} \leq 1.$$

This loss of hyperbolicity can be linked with Kelvin-Helmholtz instabilities – violent mixing that occurs when the relative difference in velocities between the two layers becomes too large.

From the above approximation of eigenvalues we also get a bound on the wave speeds of the system (1.1):

$$|\lambda| \leq |U_m| + \sqrt{g(h_1 + h_2)}.$$

This will be useful when determining CFL conditions and adding diffusion in numerical computations.

- It is easily seen that (1.1) has the steady state

$$u_1, u_2 \equiv 0, \quad b + h_1 \equiv \text{const}, \quad h_2 \equiv \text{const}. \quad (1.5)$$

This is the so-called *lake at rest* steady state. A major challenge in numerical schemes for shallow water models is the preservation of the lake at rest. Its importance is evident in the modelling of lakes and oceans, where the flows of interest are small perturbations of the lake at rest.

## 1.1 Numerical methods

We consider finite difference schemes to solve (1.1). Our spatial domain is partitioned into a uniform grid  $\{x_j\}_j$  with  $x_{j+1} - x_j \equiv \Delta x$ , and we solve for the point values  $u_j(t) \approx u(x_j, t)$ . The general form of a finite difference scheme for (1.1) is then

$$\frac{d}{dt} \mathbf{u}_j + \frac{1}{\Delta x} (\mathbf{F}_{j+1/2} - \mathbf{F}_{j-1/2}) = \mathbf{S}_j, \quad (1.6)$$

where we suppress the  $t$ -dependence of  $u_j$  for notational convenience. Temporal integration is performed with the explicit Euler method for first-order schemes and strong stability preserving Runge Kutta method for second-order schemes.

Our main goal in this paper is to design a finite difference scheme that satisfies a discrete entropy inequality

$$\frac{d}{dt} \eta(\mathbf{u}_j) + \frac{1}{\Delta x} (Q_{j+1/2} - Q_{j-1/2}) \leq 0. \quad (1.7)$$

for some numerical entropy flux  $Q_{j+1/2} = Q(\mathbf{u}_j, \mathbf{u}_{j+1})$  that is consistent with the entropy flux  $q$ . Such a scheme will be called *energy stable* after the work of Tadmor [6]. To this end, we first construct an *entropy conservative* scheme – one which satisfies

$$\frac{d}{dt} \eta(\mathbf{u}_j) + \frac{1}{\Delta x} (\widehat{Q}_{j+1/2} - \widehat{Q}_{j-1/2}) = 0, \quad (1.8)$$

and then add a numerical diffusion operator to obtain entropy stability. Define the *entropy variables* as  $\mathbf{v}(\mathbf{u}) := \nabla \eta(\mathbf{u})$  and the *entropy potential* as  $\psi(\mathbf{u}) := \mathbf{v}(\mathbf{u})^\top \mathbf{f}(\mathbf{u}) - q(\mathbf{u})$ . For the two-layer shallow water equations, the entropy variables are

$$V = \begin{bmatrix} \rho_1 \left( g(h_1 + rh_2 + b) - \frac{u_1^2}{2} \right) \\ \rho_2 \left( g(h_1 + h_2 + b) - \frac{u_2^2}{2} \right) \\ \rho_1 u_1 \\ \rho_2 u_2 \end{bmatrix}. \quad (1.9)$$

It was shown in [6] that a scheme

$$\frac{d}{dt} \mathbf{u}_j + \frac{1}{\Delta x} (\mathbf{F}_{j+1/2} - \mathbf{F}_{j-1/2}) = 0 \quad (1.10)$$

is entropy conservative if its numerical flux  $\mathbf{F}_{j+1/2}$  satisfies

$$\llbracket \mathbf{v} \rrbracket_{j+1/2}^\top \mathbf{F}_{j+1/2} = \llbracket \psi \rrbracket_{j+1/2},$$

where  $\llbracket x \rrbracket_{j+1/2} := x_{j+1} - x_j$ . Moreover, it was shown that if a numerical flux function  $\mathbf{F}$  can be written as  $\mathbf{F}_{j+1/2} = \mathbf{F}_{j+1/2}^* - D_{j+1/2} \llbracket \mathbf{v} \rrbracket_{j+1/2}$  for an entropy conservative flux  $\mathbf{F}^*$  and a positive definite diffusion matrix  $D_{j+1/2}$ , then the resulting scheme is entropy *stable*. Using these results, the authors designed energy conservative and energy stable schemes for the shallow water equations in [4]. (The entropy used is the total energy of the solution, hence the word “energy”.) A consistent discretization of the bottom topography source term in [5] led to energy conservative and energy stable, *well-balanced* methods for the shallow water equations with variable bottom topography.

## 2 The EEC scheme

It turns out that the generalization from the setting of [5] to the two-layer shallow water equations is more or less trivial. Without going into details we give the spatial discretization and refer to [4, 5] for further information:

$$\mathbf{F}_{j+1/2}^* = \begin{bmatrix} \overline{h_1 u_1} \\ \overline{h_2 u_2} \\ \frac{1}{2} g \overline{h_1^2} + \overline{h_1 u_1^2} \\ \frac{1}{2} g \overline{h_2^2} + \overline{h_2 u_2^2} \end{bmatrix}_{j+1/2}, \quad (2.1)$$

$$\mathbf{S}_j = -\frac{g}{2\Delta x} \begin{bmatrix} 0 \\ 0 \\ \overline{(h_1)_{j+1/2}} \llbracket b + rh_2 \rrbracket_{j+1/2} + \overline{(h_1)_{j-1/2}} \llbracket b + rh_2 \rrbracket_{j-1/2} \\ \overline{(h_2)_{j+1/2}} \llbracket b + h_1 \rrbracket_{j+1/2} + \overline{(h_2)_{j-1/2}} \llbracket b + h_1 \rrbracket_{j-1/2} \end{bmatrix},$$

where we use the notation  $\bar{x}_{j+1/2} := \frac{x_j + x_{j+1}}{2}$ . We denote the scheme (1.6), (2.1) the *EEC* (Explicit Energy Conservative) scheme.

**Theorem 2.1.** *The EEC scheme is consistent, second-order accurate and energy conservative – solutions  $\{\mathbf{u}_j\}_j$  satisfy the discrete entropy*

equality (1.8) with

$$\begin{aligned}\widehat{Q}_{j+1/2} &= \sum_{i=1}^2 \rho_i \left( \frac{1}{2} \overline{(h_i)_{j+1/2}} \overline{(u_i)_{j+1/2}} (u_i)_j (u_i)_{j+1} \right. \\ &\quad \left. + g \overline{(h_i)_{j+1/2}} \left( \frac{(u_i)_j (h_i + b)_{j+1} + (u_i)_{j+1} (h_i + b)_j}{2} \right) \right) \\ &\quad + \rho_2 g \left( \overline{(h_1)_{j+1/2}} \left( \frac{(u_1)_j (h_2)_{j+1} + (u_1)_{j+1} (h_2)_j}{2} \right) \right. \\ &\quad \left. + \overline{(h_2)_{j+1/2}} \left( \frac{(u_2)_j (h_1)_{j+1} + (u_2)_{j+1} (h_1)_j}{2} \right) \right)\end{aligned}$$

(compare to (1.2)). Furthermore, it is well-balanced – given initial data satisfying (1.5), the computed solution is constant in time.

*Proof.* Consistency and order of accuracy is straight-forward to check. For entropy conservation, we take the inner product of (1.6) with  $\mathbf{v}(\mathbf{u}_j)$  (see (1.9)) and obtain (1.8). Last, if the solution at some point of time satisfies the lake at rest conditions (1.5), then the two first components of the flux and source in (1.6) drop out immediately, and the third reduces to

$$\begin{aligned}&\overline{(h_1^2)_{j+1/2}} - \overline{(h_1^2)_{j-1/2}} + \overline{(h_1)_{j+1/2}} \llbracket b + rh_2 \rrbracket_{j+1/2} + \overline{(h_1)_{j-1/2}} \llbracket b + rh_2 \rrbracket_{j-1/2} \\ &= \frac{\llbracket h_1^2 \rrbracket_{j+1/2} + \llbracket h_1^2 \rrbracket_{j-1/2}}{2} + \overline{(h_1)_{j+1/2}} \llbracket b + rh_2 \rrbracket_{j+1/2} + \overline{(h_1)_{j-1/2}} \llbracket b + rh_2 \rrbracket_{j-1/2} \\ &= \overline{(h_1)_{j+1/2}} \llbracket b + h_1 + rh_2 \rrbracket_{j+1/2} + \overline{(h_1)_{j-1/2}} \llbracket b + h_1 + rh_2 \rrbracket_{j-1/2} \\ &= 0\end{aligned}$$

by (1.5). An analogous argument holds for the fourth component. Thus, we end up with  $\frac{d}{dt} \mathbf{u}_j = 0$ , whence  $\mathbf{u}_j(t) \equiv \text{constant}$  for all  $j$ .  $\square$

## 2.1 Numerical experiments

We test the EEC scheme on a problem taken from [2]. The bottom topography is set to be

$$b(x) = \begin{cases} (\cos(10\pi(x - 0.5)) + 1) / 5 & \text{if } |x - 0.5| \leq 0.1 \\ 0 & \text{otherwise.} \end{cases}$$

The initial data is the lake at rest data  $u_1, u_2 \equiv 0$ ,  $h_1 + b(x) \equiv 0.5$ ,  $h_2 \equiv 0.1$ . We discretize the domain  $[0, 1]$  into 100 grid cells and compute up to  $t = 1$ . We set  $g = 1$ ,  $\rho_1 = 1$  and  $\rho_2 = 0.9$  in this and the remaining




Figure 2.1: Lake at rest computed with the EEC scheme

experiments. As can be seen in Figure 2.1(a), there is no unphysical generation of spurious waves, and hence the initial lake at rest data is preserved.

Figure 2.1(b) shows the same experiment at  $t = 0.4$ , but with a small initial perturbation of  $+0.05$  in  $h_2$  in the range  $x \in [0.38, 0.42]$ . The two resulting shock waves are resolved, but in-between there are unphysical oscillations that are due to a lack of numerical diffusion. To fix this we will add numerical diffusion to obtain an *energy stable* scheme.

### 3 The ERus scheme

To obtain energy decay near discontinuities we add a Rusanov-type numerical diffusion operator of the form  $c_{j+1/2} \llbracket \mathbf{u} \rrbracket_{j+1/2}$  to the numerical flux, where  $c_{j+1/2}$  is some approximation of the largest eigenvalues of the system. Assume for the moment that the bottom topography is flat,  $b \equiv 0$ . As the energy  $\eta$  is strictly convex, the mapping  $\mathbf{u} \mapsto \mathbf{v}(\mathbf{u})$  is injective, and so the change-of-variables matrix  $\partial_{\mathbf{v}} \mathbf{u}(\mathbf{v})$  is well-defined and positive definite. By the mean value theorem there is a state  $\mathbf{v}_{j+1/2}$  such that

$$\llbracket \mathbf{u} \rrbracket_{j+1/2} = \partial_{\mathbf{v}} \mathbf{u}(\mathbf{v}_{j+1/2}) \llbracket \mathbf{v} \rrbracket_{j+1/2}. \quad (3.1)$$

This is the form of our numerical diffusion operator, and we define the *ERus* (Energy stable Rusanov) flux to be

$$\mathbf{F}_{j+1/2} = \mathbf{F}_{j+1/2}^* - \frac{1}{2} c_{j+1/2} \partial_{\mathbf{v}} \mathbf{u}(\mathbf{v}_{j+1/2}) \llbracket \mathbf{v} \rrbracket_{j+1/2}, \quad (3.2)$$

where  $\mathbf{F}^*$  is the EEC flux (2.1). In the more general case  $b \neq 0$  we still use the above expression, although the identity (3.1) no longer holds since  $\mathbf{v}$ , unlike  $\mathbf{u}$ , depends explicitly on  $b$ . We set the diffusion coefficient  $c_{j+1/2}$  to be

$$c_{j+1/2} = \max(c_j, c_{j+1}),$$

where  $c_j := |(U_m)_j| + \sqrt{g((h_1)_j + (h_2)_j)}$  (see (1.4)).

**Theorem 3.1.** *The ERus scheme is consistent, first-order accurate and energy stable – solutions  $\{\mathbf{u}_j\}_j$  satisfy (1.7) with  $Q_{j+1/2} = \widehat{Q}_{j+1/2} - \frac{1}{2}\bar{\mathbf{v}}_{j+1/2}^\top D_{j+1/2} \llbracket \mathbf{v} \rrbracket_{j+1/2}$  and  $D_{j+1/2} := \frac{1}{2}c_{j+1/2} \partial_{\mathbf{v}} \mathbf{u}(\mathbf{v}_{j+1/2})$ . Furthermore, it is well-balanced – given initial data satisfying (1.5), the computed solution is constant in time.*

*Proof.* First-order accuracy comes from the  $\mathcal{O}(\Delta x)$  term  $\llbracket \mathbf{v} \rrbracket_{j+1/2}$ . The proof of the discrete entropy inequality follows [6, Theorem 5.2]: Multiply the ERus scheme (1.6) by  $\mathbf{v}_j^\top$ , use the entropy conservativity of the EEC flux  $\mathbf{F}^*$  and rearrange the diffusion terms to obtain (??).

For well-balancedness, we note that when the solution satisfies (1.5), the entropy variables (1.9) are constant in space. Hence, the ERus flux reduces to the EEC flux, which we have already shown is well-balanced.  $\square$

### 3.1 Numerical experiments

We repeat the numerical experiments from the previous section with the ERus scheme. The lake at rest is preserved exactly; even after long-time simulations the relative error stays within  $10^{-16}$ . Figure 3.1(a) shows the perturbed lake at rest experiment. Clearly, the unphysical oscillations of the EEC scheme are gone.

Figure 3.2 shows computed  $h_1$  and  $u_1$  on a problem from [2]. The results compare well to those of [2].




Figure 3.1: Perturbed lake at rest computed with the ERus scheme.






Figure 3.2: Problem from [2].  $h_1$  to the left,  $u_1$  to the right. Top row with 500 grid points, bottom row with 10000.

## References

- [1] R. Abgrall and S. Karni. A comment on the computation of non-conservative products. *Journal of Computational Physics*, 229:2759–2763, April 2010.
- [2] Rémi Abgrall and Smadar Karni. Two-layer shallow water system: A relaxation approach. *SIAM Journal on Scientific Computing*, 31(3):1603–1627, 2009.
- [3] G. DalMaso, P. G. LeFloch, and F. Murat. Definition and weak stability of nonconservative products. *Journal de Mathématiques Pures et Appliquées*, 74(6):483–548, 1995.
- [4] U. S. Fjordholm, S. Mishra, and E. Tadmor. Energy preserving and energy stable schemes for the shallow water equations. In *Foundations of Computational Mathematics*, Proc. FoCM held in Hong Kong 2008, volume 363 of *London Math. Soc. Lecture Notes Series*, pages 93–139, 2009.
- [5] U. S. Fjordholm, S. Mishra, and E. Tadmor. Well-balanced, energy stable schemes for the shallow water equations with varying topography. Submitted, 2010.
- [6] E. Tadmor. The numerical viscosity of entropy stable schemes for systems of conservation laws. I. *Mathematics of Computation*, 49:91–103, 1987.

# Research Reports

No.	Authors/Title
10-41	<i>U.S. Fjordholm</i> Energy conservative and -stable schemes for the two-layer shallow water equations
10-40	<i>R. Andreev and Ch. Schwab</i> Sparse tensor approximation of parametric eigenvalue problems
10-39	<i>R. Hiptmair, A. Moiola and I. Perugia</i> Stability results for the time-harmonic Maxwell equations with impedance boundary conditions
10-38	<i>I. Hnětynková, M. Plešinger, D.M. Sima, Z. Strakoš and S. Van Huffel</i> The total least squares problem in $AX \approx B$ . A new classification with the relationship to the classical works
10-37	<i>S. Mishra</i> Robust finite volume schemes for simulating waves in the solar atmosphere
10-36	<i>C. Effenberger, D. Kressner and C. Engström</i> Linearization techniques for band structure calculations in absorbing photonic crystals
10-35	<i>R. Hiptmair and C. Jerez-Hanckes</i> Multiple traces boundary integral formulation for Helmholtz transmission problems
10-34	<i>H. Harbrecht and Ch. Schwab</i> Sparse tensor finite elements for elliptic multiple scale problems
10-33	<i>K. Grella and C. Schwab</i> Sparse tensor spherical harmonics approximation in radiative transfer
10-32	<i>P. Kauf, M. Torrilhon and M. Junk</i> Scale-induced closure for approximations of kinetic equations
10-31	<i>M. Hansen</i> On tensor products of quasi-Banach spaces
10-30	<i>P. Corti</i> Stable numerical scheme for the magnetic induction equation with Hall effect
10-29	<i>H. Kumar</i> Finite volume methods for the two-fluid MHD equations