

An A Priori Error Estimate for Interior Penalty Discretizations of the Curl-Curl Operator on Non-Conforming Meshes

R. Casagrande and R. Hiptmair

Research Report No. 2014-40
December 2014

Seminar für Angewandte Mathematik
Eidgenössische Technische Hochschule
CH-8092 Zürich
Switzerland

An A Priori Error Estimate for Interior Penalty Discretizations of the Curl-Curl Operator on Non-Conforming Meshes

Raffael Casagrande and Ralf Hiptmair

Abstract We prove an a-priori error estimate for conductivity-regularized Curl-Curl Problems which are discretized by the Interior Penalty/Nitsche's Method on meshes non-conforming across interfaces. It is shown that the total error can be bounded by the best approximation error which in turn depends on the concrete choice of the approximation space V_h . In this work we show that if V_h is the space of edge functions of the first kind of order k we can expect (suboptimal) convergence $O(h^{k-1})$ as the mesh is refined. The numerical experiments in Casagrande, Winkelmann, Hiptmair and Ostrowski, SAM Report 2014-32, ETH Zürich, indicate that this bound is sharp for $k = 1$. Moreover it is shown that the regularization term can be made arbitrarily small without affecting the error in the $|\cdot|_{\text{curl}}$ semi-norm. A numerical experiment shows that the regularization parameter can be chosen in a wide range of values such that, at the same time, the discrete problem remains solvable and the error due to regularization is negligible compared to the discretization error.

1 Introduction

In this work we study the 3D, *regularized* Curl-Curl boundary value problem,

$$\nabla \times (\mu^{-1} \nabla \times \mathbf{A}) + \varepsilon \mathbf{A} = \mathbf{j}^i, \quad \text{in } \Omega \quad (1)$$


$$\mathbf{n} \times \mathbf{A} = \mathbf{g}_D \quad \text{on } \partial\Omega, \quad (2)$$

which can be used to calculate the magnetic field that originates from a stationary current \mathbf{j}^i . Herein μ denotes the magnetic permeability, \mathbf{g}_D prescribes Dirichlet boundary data and $\varepsilon > 0$ is the regularization parameter that renders the solution unique. We seek the magnetic vector potential \mathbf{A} that fulfills (1-2). The magnetic field is then $\mathbf{B} = \nabla \times \mathbf{A}$. Note that if $\mathbf{g}_D \equiv 0$ on $\partial\Omega$ then (2) implies $(\nabla \times \mathbf{A}) \cdot \mathbf{n} = \mathbf{B} \cdot \mathbf{n} = 0$ on $\partial\Omega$ which reflects the decay of the fields away from the source.

Raffael Casagrande, Ralf Hiptmair
Seminar for Applied Mathematics, ETH Zürich, Rämistr. 101, CH-8092 Zürich,
e-mail: raffael.casagrande@sam.math.ethz.ch, hiptmair@sam.math.ethz.ch

In some applications like the simulation of electric machines or magnetic actuators, magnetic fields have to be computed in the presence of moving, rigid parts. Then one may use separate, moving sub-meshes for them in order to avoid remeshing. However, this leads to so-called “sliding interfaces”, i.e. meshes with hanging nodes (cf. Fig. 1).

Fig. 1 Initially conforming sub-meshes become non-conforming when the upper sub-mesh starts moving.



In [1] the authors applied the Interior Penalty/Nitsche’s Method [2] to Curl-Curl type problems in order to handle arbitrary, non-conforming mesh interfaces. Therein it was shown experimentally that the Interior Penalty Method solves problem (1-2) successfully if second order edge functions of the first kind are used. Moreover it was shown that first order edge functions fail to converge to the exact solution as the mesh is refined. In this work we intend to give theoretical explanations of these observations and investigate the effect of the regularization term in (1).

We start our discussion in Section 2 by introducing Discontinuous Galerkin (DG) notations that were already introduced in [1] and which are needed to state the interior penalty formulation of (1-2) in Section 3. Section 3 also proves an a-priori bound on the total error in terms of the best approximation error for piecewise-polynomial test- and trial spaces V_h . In Section 4 we analyze the particular case where V_h is the space of k -th order edge functions, R^k . Combining the results of Sections 3 and 4 we get rates of convergence for the regularized problem (1-2). Section 5 is devoted to the choice of the local length scale appearing in the Interior Penalty formulation and Section 6 discusses the role of the regularity parameter ε and how to choose it. We end our presentation with a short conclusion and outlook in Section 7.

2 Preliminaries

Before we can introduce the Symmetric Weighted Interior penalty (SWIP) formulation of (1-2) we give some definitions and notations (cf. [1]):

Subdomains and sub meshes: Let us assume that the domain Ω , on which (1-2) is posed, is a simply connected polyhedron with Lipschitz boundary. Furthermore we assume Ω to be split into two non-overlapping subdomains $\overline{\Omega}_1 \cup \overline{\Omega}_2 = \overline{\Omega}$. On each subdomain we introduce a sequence of conforming, simplicial meshes $\mathcal{T}_{\mathcal{H},1}, \mathcal{T}_{\mathcal{H},2}$ and let $\mathcal{T}_{\mathcal{H}} = \mathcal{T}_{\mathcal{H},1} \cup \mathcal{T}_{\mathcal{H},2}$. For any $h \in \mathcal{H}$ we let \mathcal{T}_h denote a particular mesh in the sequence $\mathcal{T}_{\mathcal{H}}$ and $T \in \mathcal{T}_h$ a mesh element (tetrahedron). The meshwidth is then defined as $h = \max_{T \in \mathcal{T}_h} h_T$ where h_T is the diameter of element T .

Furthermore we define \mathbb{F}_T to be the set of the four *facets* of tetrahedron T . The intersection of two facets, belonging to two neighboring elements, is called an inner *face* while the intersection of a facet with the boundary $\partial\Omega$ is called a boundary face. Note that facets are always triangular while inner faces are convex polygons with up to six nodes and boundary faces can have virtually any polygonal shape. We denote by \mathcal{F}_h^b the set of all boundary faces, \mathcal{F}_h^i the set of all inner faces

and define $\mathcal{F}_h = \mathcal{F}_h^b \cup \mathcal{F}_h^i$ to be the set of all faces. Furthermore, \mathcal{F}_T stands for the set of all faces which lie on the boundary of element T .

Mesh assumptions We assume that the elements are *shape regular* in the sense of Ciarlet: There is a constant σ_{\max} such that $\forall h \in \mathcal{H}, \forall T \in \mathcal{T}_h$ we have

$$\frac{h_T}{\rho_T} \leq \sigma_{\max} \quad (3)$$

where ρ_T is the radius of the largest ball inscribed in T . It is easy to check that this condition is satisfied if two sequences of static sub-meshes are moved against each other. We will make additional assumptions about the mesh when we discuss choices for the local length scale in Sect. 5.

Magnetic Permeability: We assume there exists a partition $P_\Omega = \{\Omega_{i,\mu}\}$ such that each $\Omega_{i,\mu}$ is a polyhedron and such that the permeability $\mu > 0$ is constant on each $\Omega_{i,\mu}$. Furthermore the mesh sequence $\mathcal{T}_\mathcal{H}$ is *compatible* with the partition P_Ω : For each $\mathcal{T}_h \in \mathcal{T}_\mathcal{H}$, each element $T \in \mathcal{T}_h$ belongs to exactly one $\Omega_{i,\mu} \in P_\Omega$. I.e. the magnetic permeability is allowed to jump over element boundaries, and in particular over the non-conforming interface $\Gamma = \overline{\Omega}_1 \cap \overline{\Omega}_2$

Polynomial approximation: Later on we will seek our discrete solution in the piecewise polynomial space (cf. [3]),

$$\mathbb{P}^k(\mathcal{T}_h) := \left\{ p \in L^2(\Omega) \mid \forall T \in \mathcal{T}_h, p|_T \in \mathbb{P}^k(T) \right\} \quad (4)$$

where $\mathcal{T}_h \in \mathcal{T}_\mathcal{H}$ and $\mathbb{P}^k(T)$ is the usual space of polynomials up to degree k on mesh element T . Note that functions of $\mathbb{P}^k(\mathcal{T}_h)$ are discontinuous across element boundaries.

Mesh Faces, Jump and Average Operators For each mesh face F and *vector valued* function $\mathbf{A}_h \in \mathbb{P}^k(\mathcal{T}_h)^3$, we define the *tangential jump* as

$$\begin{aligned} \text{if } F \in \mathcal{F}_h^i, F = \partial T_1 \cap \partial T_2 : & \quad [\mathbf{A}_h]_T = \mathbf{n}_F \times (\mathbf{A}_h|_{T_1} - \mathbf{A}_h|_{T_2}), \\ \text{if } F \in \mathcal{F}_h^b, F \subseteq \partial T_1 \cap \partial \Omega : & \quad [\mathbf{A}_h]_T = \mathbf{n}_F \times \mathbf{A}_h|_{T_1}, \end{aligned}$$

and the *weighted average* as

$$\begin{aligned} \text{if } F \in \mathcal{F}_h^i, F = \partial T_1 \cap \partial T_2 : & \quad \{\mathbf{A}_h\}_\omega = \omega_1 \mathbf{A}_h|_{T_1} + \omega_2 \mathbf{A}_h|_{T_2}, \\ \text{if } F \in \mathcal{F}_h^b, F \subseteq \partial T_1 \cap \partial \Omega : & \quad \{\mathbf{A}_h\}_\omega = \mathbf{A}_h|_{T_1}. \end{aligned}$$

Here \mathbf{n}_F always points from T_1 to T_2 and $\omega_1, \omega_2 \in [0, 1]$ such that $\omega_1 + \omega_2 = 1$. Note that the jump and average operators are well defined for all $\mathbf{p} \in \mathbb{P}^k(\mathcal{T}_h)^3$. Indeed, since $\mathbf{p}|_T \in C^\infty(T)$, the trace operator $\gamma_0(\mathbf{p}) = \mathbf{p}|_{\partial T}$ is well defined.

The following Lemma relates the trace of a polynomial function to its L^2 norm on the element:

Lemma 1 (Discrete Trace Inequality). *Let $\mathcal{T}_\mathcal{H}$ be a sequence of shape regular, possibly non-conforming, simplicial meshes. Then for all $h \in \mathcal{H}$, all $v_h \in \mathbb{P}^k(\mathcal{T}_h)$, and all $T \in \mathcal{T}_h$ we have*

$$h_T^{1/2} \|v_h|_T\|_{L^2(\partial T)} \leq C_{Tr} \|v_h\|_{L^2(T)} \quad (5)$$

where C_{Tr} is independent of T , h but depends on σ_{\max} , k .

Proof. We split the integral over ∂T into the four triangular facets $F_{T,i} \in \mathbb{F}_T$, $i = 1, \dots, 4$. Let now $\Phi_T : \hat{T} \rightarrow T$, $\hat{\mathbf{x}} \mapsto \mathbf{B}_T \hat{\mathbf{x}} + \mathbf{b}_T$ be the affine transformation that maps the reference element \hat{T} to T . Then,

$$\begin{aligned} \|v_h\|_{L^2(F_{T,i})}^2 &\leq \text{area}(F_{T,i}) \|v_h\|_{L^\infty(F_{T,i})}^2 \leq h_T^2 \|v_h\|_{L^\infty(\hat{T})}^2 \\ &\leq Ch_T^2 \|v_h\|_{L^2(\hat{T})}^2, \end{aligned}$$

where we have used the equivalence of norms in finite dimensions in the last step. Now, transforming back to the element T we get:

$$\|v_h\|_{L^2(F_{T,i})}^2 \leq C \frac{h_T^2}{|\det(\mathbf{B}_T)|} \|v_h\|_{L^2(T)}^2 \leq C' h_T^{-1} \sigma_{\max}^3 \|v_h\|_{L^2(T)}^2$$

where we have used $|\det(\mathbf{B}_T)| \geq C'' \rho_T^3$ (see [9][Lemma 5.10]) and the shape regularity (3). Summing over all facets $F_{T,i}$ yields the assertion. \square

Function Spaces We will use the following spaces

$$\begin{aligned} \mathbf{H}(\mathbf{curl}; \Omega) &:= \left\{ \mathbf{f} \in L^2(\Omega)^3 \mid \|\nabla \times \mathbf{f}\|_{L^2(\Omega)^3} < \infty \right\} \\ H^s(P_\Omega) &:= \left\{ f \in L^2(\Omega) \mid \forall \Omega_{i,\mu} \in P_\Omega : f|_{\Omega_{i,\mu}} \in H^s(\Omega_{i,\mu}) \right\}, \end{aligned}$$

Herein L^2 denotes the usual space of square integrable functions and $H^s(\Omega) = W^{s,2}(\Omega)$ is the Sobolev space of order s with Hölder coefficient $p = 2$. The associated norms and semi-norms are:

$$\begin{aligned} \|\mathbf{f}\|_{\mathbf{H}(\mathbf{curl}; \Omega)}^2 &:= \|\mathbf{f}\|_{L^2(\Omega)^3}^2 + \|\mathbf{f}\|_{\mathbf{H}(\mathbf{curl}; \Omega)}^2, & \|\mathbf{f}\|_{\mathbf{H}(\mathbf{curl}; \Omega)}^2 &:= \|\nabla \times \mathbf{f}\|_{L^2(\Omega)^3}^2, \\ \|f\|_{H^s(P_\Omega)}^2 &:= \sum_{i=1}^N \|f\|_{H^s(\Omega_{i,\mu})}^2. \end{aligned}$$

3 Symmetric Weighted Interior Penalty (SWIP) Formulation

We chose an arbitrary subspace $V_h \subseteq \mathbb{P}^k(\mathcal{T}_h)^3$ as discrete test and trial space, and use integration by parts (cf. [3, 7] for details) to arrive at the SWIP formulation of (1): Find $\mathbf{A}_h \in V_h$ such that

$$a_h^{\text{SWIP}}(\mathbf{A}_h, \mathbf{A}'_h) = \ell_h(\mathbf{A}'_h) \quad \forall \mathbf{A}'_h \in V_h \quad (6)$$

with

$$\begin{aligned}
a_h^{\text{SWIP}}(\mathbf{A}_h, \mathbf{A}'_h) &= \int_{\Omega} (\mu^{-1} \nabla \times \mathbf{A}_h) \cdot (\nabla \times \mathbf{A}'_h) - \sum_{F \in \mathcal{F}_h} \int_F \{\mu^{-1} \nabla \times \mathbf{A}_h\}_{\omega} \cdot [\mathbf{A}'_h]_T \\
&\quad - \sum_{F \in \mathcal{F}_h} \int_F \{\mu^{-1} \nabla \times \mathbf{A}'_h\}_{\omega} \cdot [\mathbf{A}_h]_T + \sum_{F \in \mathcal{F}_h} \frac{\eta \gamma_{\mu, F}}{a_F} \int_F [\mathbf{A}_h]_T \cdot [\mathbf{A}'_h]_T + \varepsilon \int_{\Omega} \mathbf{A}_h \cdot \mathbf{A}'_h, \quad (7)
\end{aligned}$$

$$\begin{aligned}
\ell_h(\mathbf{A}'_h) &= \int_{\Omega} \mathbf{j}^i \cdot \mathbf{A}'_h - \sum_{F \in \mathcal{F}_h^b} \int_F \{\mu^{-1} \nabla \times \mathbf{A}'_h\}_{\omega} \cdot (\mathbf{n} \times \mathbf{g}_D) \\
&\quad + \sum_{F \in \mathcal{F}_h^b} \frac{\eta \gamma_{\mu, F}}{a_F} \int_F [\mathbf{A}'_h]_T \cdot (\mathbf{n} \times \mathbf{g}_D). \quad (8)
\end{aligned}$$

where η is the penalty parameter. The last four terms of a_h^{SWIP} are called consistency, symmetry, penalty, regularization term, respectively. For an inner face $F \in \mathcal{F}_h^i$, $F = \partial T_1 \cap \partial T_2$, we chose the weights as

$$\gamma_{\mu, F} := \frac{1}{\mu_1 + \mu_2}, \quad \omega_1 := \frac{\mu_1}{\mu_1 + \mu_2}, \quad \omega_2 := \frac{\mu_2}{\mu_1 + \mu_2}.$$

If $F \in \mathcal{F}_h^b$ we choose $\gamma_{\mu, F} := \mu^{-1}$. The term a_F is the *local length scale* of face F and can be chosen in different ways (e.g. $a_F = \frac{1}{2}(h_{T_1} + h_{T_2})$ where h_{T_1}, h_{T_2} are the diameters of the neighboring elements). For now we assume that there exists a constant $\zeta_2 > 0$ such that for all $h \in \mathcal{H}$, all $T \in \mathcal{T}_h$, and all $F \in \mathcal{F}_T$:

$$0 < a_F \leq \zeta_2 h_T. \quad (9)$$

In Section 5 we will look at concrete choices of a_F and discuss the circumstances under which (9) is fulfilled. It will turn out that depending on the choice of a_F we have to make additional assumptions about the mesh regularity to guarantee (9).

Remark If $V_h \subseteq \mathbf{H}(\mathbf{curl}; \Omega)$, then all inner tangential jumps in (7) will drop out [7, Lemma 3.8] and only jumps at the boundary remain, i.e. we are left with a standard FEM formulation where the inhomogeneous boundary conditions (2) are enforced in a weak sense.

3.1 A Priori Error Estimate

In the following we derive an error estimate in the ‘‘energy-norm’’ for the variational problem (6).

Regularity of the exact solution We assume that the exact solution \mathbf{A} of (1-2) (in the sense of distributions) is such that

$$\mathbf{A} \in V^* := \{ \mathbf{A} \in \mathbf{H}(\mathbf{curl}; \Omega) \cap H^1(P_{\Omega})^3 \mid \nabla \times \mathbf{A} \in H^1(P_{\Omega})^3 \}$$

Furthermore we set $V_h^* := V^* + V_h$. Note that because $\mathbf{A}, \nabla \times \mathbf{A} \in H^1(P_{\Omega})^3$ the traces of \mathbf{A} and $\nabla \times \mathbf{A}$ are well defined on the faces of the mesh elements (cf. [3, Remark 1.26]). Indeed, let $T \in \mathcal{T}_h$ be a mesh element, then by the multiplicative trace inequality [4, Thm. 1.6.6]: $\|\mathbf{A}\|_{L^2(\partial T)^3} <$

$C_{\text{tr}} \|\mathbf{A}\|_{L^2(T)^3}^{1/2} \|\mathbf{A}\|_{H^1(T)^3}^{1/2}$ and the same estimate holds for $\nabla \times \mathbf{A}$. Therefore we see that $a_h^{\text{SWIP}} : V_h^* \times V_h \rightarrow \mathbb{R}$ is well defined.

In order for the right-hand side to be well-posed we assume $\mathbf{j}^i \in L^2(\Omega)^3$ and $\mathbf{g}_D = L^2(\partial\Omega)^3$.

We begin the proof of the a priori error estimate by showing that the exact solution \mathbf{A} fulfills equation (6):

Lemma 2 (Consistency). *Assume $\mathbf{A} \in V^*$ is the exact solution of (1-2). Then, for all $\mathbf{A}'_h \in V_h$,*

$$a_h^{\text{SWIP}}(\mathbf{A}, \mathbf{A}'_h) = \ell_h(\mathbf{A}'_h) \quad (10)$$

Proof. Since $\mathbf{A} \in \mathbf{H}(\mathbf{curl}, \Omega)$, \mathbf{A} is tangentially continuous across all element boundaries (cf. Lemma 3.8 in [7]). Thus all inner jump terms drop out,

$$\begin{aligned} a_h^{\text{SWIP}}(\mathbf{A}, \mathbf{A}'_h) &= \int_{\Omega} \left(\frac{1}{\mu} \nabla \times \mathbf{A} \right) \cdot (\nabla \times \mathbf{A}'_h) - \sum_{F \in \mathcal{F}_h} \int_F \{ \mu^{-1} \nabla \times \mathbf{A} \}_{\omega} \cdot [\mathbf{A}'_h]_T \\ &- \sum_{F \in \mathcal{F}_h^b} \int_F \{ \mu^{-1} \nabla \times \mathbf{A}'_h \}_{\omega} \cdot [\mathbf{A}]_T + \sum_{F \in \mathcal{F}_h^b} \frac{\eta \gamma_{\mu, F}}{a_F} \int_F [\mathbf{A}]_T \cdot [\mathbf{A}'_h]_T + \varepsilon \int_{\Omega} \mathbf{A}_h \cdot \mathbf{A}'_h \end{aligned} \quad (11)$$

Note that the two last two sums include only boundary faces. Next we make use of the following identity (which holds for any interior face $F = \partial T_1 \cap \partial T_2$):

$$\begin{aligned} [\mathbf{a} \times \mathbf{b}]_n &= (\mathbf{a}_1 \times \mathbf{b}_1 - \mathbf{a}_2 \times \mathbf{b}_2) \cdot \mathbf{n}_F, \\ &= ((\omega_1 \mathbf{a}_1 + \omega_2 \mathbf{a}_2) \times (\mathbf{b}_1 - \mathbf{b}_2) + (\mathbf{a}_1 - \mathbf{a}_2) \times (\omega_2 \mathbf{b}_1 + \omega_1 \mathbf{b}_2)) \cdot \mathbf{n}_F \\ &= -\{\mathbf{a}\}_{\omega} \cdot [\mathbf{b}]_T + [\mathbf{a}]_T \cdot \{\mathbf{b}\}_{\bar{\omega}}, \end{aligned}$$

where $\{\mathbf{b}\}_{\bar{\omega}} := (\omega_2 \mathbf{b}_1 + \omega_1 \mathbf{b}_2)$ is the skew-weighted average and $[\mathbf{b}]_n = (\mathbf{b}_1 - \mathbf{b}_2) \cdot \mathbf{n}_F$ is the normal jump. Let us apply the identity to the second term of (11):

$$\begin{aligned} - \sum_{F \in \mathcal{F}_h^b} \int_F \{ \mu^{-1} \nabla \times \mathbf{A} \}_{\omega} \cdot [\mathbf{A}'_h]_T &= \sum_{F \in \mathcal{F}_h^i} \int_F [(\mu^{-1} \nabla \times \mathbf{A}) \times \mathbf{A}'_h]_n \\ &- \sum_{F \in \mathcal{F}_h^i} \int_F \underbrace{[\mu^{-1} \nabla \times \mathbf{A}]_T}_{=0} \cdot \{\mathbf{A}'_h\}_{\bar{\omega}} - \sum_{F \in \mathcal{F}_h^b} \int_F \{ \mu^{-1} \nabla \times \mathbf{A} \}_{\omega} \cdot [\mathbf{A}'_h]_T, \end{aligned}$$

where the second term on the right-hand side vanishes because \mathbf{A} is a solution of the strong formulation (1) and thus $\mu^{-1} \nabla \times \mathbf{A} \in \mathbf{H}(\mathbf{curl}; \Omega)$ which implies that $\mu^{-1} \nabla \times \mathbf{A}$ is tangentially continuous. Note that for $F \in \mathcal{F}_h^b$: $\{ \mu^{-1} \nabla \times \mathbf{A} \}_{\omega} \cdot [\mathbf{A}'_h]_T = - [(\mu^{-1} \nabla \times \mathbf{A}) \times \mathbf{A}'_h]_n$ so we can rearrange the face contributions to the elements boundaries:

$$- \sum_{F \in \mathcal{F}_h^b} \int_F \{ \mu^{-1} \nabla \times \mathbf{A} \}_{\omega} \cdot [\mathbf{A}'_h]_T = \sum_{T \in \mathcal{T}_h} \int_{\partial T} (\mu^{-1} (\nabla \times \mathbf{A}) \times \mathbf{A}'_h) \cdot \mathbf{n}_T \quad (12)$$

Now substitute (12) into (11) and use integration by parts on each mesh element [9, Thm. 3.29]:

$$\begin{aligned}
a_h^{\text{SWIP}}(\mathbf{A}, \mathbf{A}'_h) &= \sum_{T \in \mathcal{T}_h} \int_T \nabla \times \left(\frac{1}{\mu} \nabla \times \mathbf{A} \right) \cdot \mathbf{A}'_h + \varepsilon \int_{\Omega} \mathbf{A}_h \cdot \mathbf{A}'_h \\
&\quad - \sum_{F \in \mathcal{F}_h^b} \int_F \{ \mu^{-1} \nabla \times \mathbf{A}'_h \}_\omega \cdot [\mathbf{A}]_T + \sum_{F \in \mathcal{F}_h^b} \frac{\eta \gamma_{\mu, F}}{a_F} \int_F [\mathbf{A}]_T \cdot [\mathbf{A}'_h]_T \\
&\stackrel{(1-2)}{=} \ell_h(\mathbf{A}'_h)
\end{aligned}$$

□

Let us introduce the following (semi-)norms on the space V_h^* :

$$\begin{aligned}
\|\mathbf{A}\|_{\text{SWIP}}^2 &:= \left\| \mu^{-1/2} \nabla \times \mathbf{A} \right\|_{L^2(\Omega)^3}^2 + \left\| \varepsilon^{1/2} \mathbf{A} \right\|_{L^2(\Omega)^3}^2 + |\mathbf{A}|_{j, \mu}^2 \\
|\mathbf{A}|_{j, \mu}^2 &:= \sum_{F \in \mathcal{F}_h} \frac{\gamma_{\mu, F}}{a_F} \left\| [\mathbf{A}]_T \right\|_{L^2(F)^3}^2 \\
\|\mathbf{A}\|_{\text{SWIP},*}^2 &:= \|\mathbf{A}\|_{\text{SWIP}}^2 + \sum_{T \in \mathcal{T}_h} h_T \left\| \mu^{-1/2} \nabla \times \mathbf{A} \Big|_T \right\|_{L^2(\partial T)^3}^2
\end{aligned}$$

Lemma 3 (Bound on consistency term). *For all $\mathbf{A}, \mathbf{A}' \in V_h^*$ there holds:*

$$\left| \sum_{F \in \mathcal{F}_h} \int_F \{ \mu^{-1} \nabla \times \mathbf{A} \}_\omega \cdot [\mathbf{A}']_T \right| \leq \zeta_2^{1/2} \left\{ \sum_{T \in \mathcal{T}_h} h_T \left\| \mu^{-1/2} \nabla \times \mathbf{A} \Big|_T \right\|_{L^2(\partial T)^3}^2 \right\}^{1/2} |\mathbf{A}'|_{j, \mu}$$

where $h_T = \max \{ \|x - y\| \mid x, y \in T \}$ is the diameter of mesh element $T \in \mathcal{T}_h$.

Proof. For an arbitrary inner face $F = \partial T_1 \cap \partial T_2$ we have by the Cauchy-Schwarz (CS) inequality:

$$\left| \int_F \{ \mu^{-1} \nabla \times \mathbf{A} \}_\omega \cdot [\mathbf{A}']_T \right| \leq \left| \int_F \left(\frac{\omega_1}{\mu_1} \nabla \times \mathbf{A}_1 + \frac{\omega_2}{\mu_2} \nabla \times \mathbf{A}_2 \right)^2 \right|^{1/2} \left| \int_F \left\| [\mathbf{A}']_T \right\|^2 \right|^{1/2} \quad (13)$$

Using Cauchy-Schwarz again we see that

$$\begin{aligned}
\left\| \frac{\omega_1}{\mu_1} \nabla \times \mathbf{A}_1 + \frac{\omega_2}{\mu_2} \nabla \times \mathbf{A}_2 \right\| &\leq \left(\frac{\omega_1^2}{\mu_1} + \frac{\omega_2^2}{\mu_2} \right)^{1/2} \left(\left\| \mu_1^{-1/2} \nabla \times \mathbf{A}_1 \right\|^2 + \left\| \mu_2^{-1/2} \nabla \times \mathbf{A}_2 \right\|^2 \right)^{1/2} \\
&= a_F^{1/2} \left(\frac{\gamma_{\mu, F}}{a_F} \right)^{1/2} \left(\left\| \mu_1^{-1/2} \nabla \times \mathbf{A}_1 \right\|^2 + \left\| \mu_2^{-1/2} \nabla \times \mathbf{A}_2 \right\|^2 \right)^{1/2}
\end{aligned}$$

Substitute this back into (13):

$$\begin{aligned}
\left| \int_F \{ \mu^{-1} \nabla \times \mathbf{A} \}_\omega \cdot [\mathbf{A}']_T \right| &\leq \left[a_F \int_F \left(\left\| \mu_1^{-1/2} \nabla \times \mathbf{A}_1 \right\|^2 + \left\| \mu_2^{-1/2} \nabla \times \mathbf{A}_2 \right\|^2 \right) \right]^{1/2} \\
&\quad \left(\frac{\gamma_{\mu, F}}{a_F} \right)^{1/2} \left\| [\mathbf{A}']_T \right\|_{L^2(F)^3}. \quad (14)
\end{aligned}$$

Similarly, for a boundary face $F \in \mathcal{F}_h^b$ we have:

$$\left| \int_F \{ \mu^{-1} \nabla \times \mathbf{A} \}_\omega \cdot [\mathbf{A}']_T \right| \stackrel{\text{(CS)}}{\leq} \left| a_F \int_F \left\| \mu^{-1/2} \nabla \times \mathbf{A} \right\|^2 \right|^{1/2} \frac{1}{\sqrt{\mu a_F}} \| [\mathbf{A}']_T \|_{L^2(F)} \quad (15)$$

Now use (14-15) to bound the sum over all faces:

$$\begin{aligned} & \left| \sum_{F \in \mathcal{F}_h^b} \int_F \{ \mu^{-1} \nabla \times \mathbf{A} \}_\omega \cdot [\mathbf{A}']_T + \sum_{F \in \mathcal{F}_h^i} \int_F \{ \mu^{-1} \nabla \times \mathbf{A} \}_\omega \cdot [\mathbf{A}']_T \right| \\ & \stackrel{(14-15)}{\leq} \sum_{F \in \mathcal{F}_h^b} \left\{ \left| a_F \int_F \left\| \mu^{-1/2} \nabla \times \mathbf{A} \right\|^2 \right|^{1/2} \frac{1}{\sqrt{\mu a_F}} \| [\mathbf{A}']_T \|_{L^2(F)} \right\} + \\ & \quad \sum_{F \in \mathcal{F}_h^i} \left\{ \left[a_F \int_F \left(\left\| \mu_1^{-1/2} \nabla \times \mathbf{A}_1 \right\|^2 + \left\| \mu_2^{-1/2} \nabla \times \mathbf{A}_2 \right\|^2 \right) \right]^{1/2} \left(\frac{\gamma_{\mu,F}}{a_F} \right)^{1/2} \| [\mathbf{A}']_T \|_{L^2(F)^3} \right\} \\ & \stackrel{\text{(CS)}}{\leq} \left\{ \sum_{F \in \mathcal{F}_h} \frac{\gamma_{\mu,F}}{a_F} \| [\mathbf{A}']_T \|_{L^2(F)^3}^2 \right\}^{1/2} \\ & \quad \left\{ \sum_{F \in \mathcal{F}_h^i} \left[a_F \int_F \left(\left\| \mu_1^{-1/2} \nabla \times \mathbf{A}_1 \right\|^2 + \left\| \mu_2^{-1/2} \nabla \times \mathbf{A}_2 \right\|^2 \right) \right] + \sum_{F \in \mathcal{F}_h^b} a_F \int_F \left\| \mu^{-1/2} \nabla \times \mathbf{A} \right\|^2 \right\}^{1/2} \\ & \leq \zeta_2^{1/2} \left\{ \sum_{T \in \mathcal{T}_h} h_T \int_{\partial T} \left\| \mu^{-1/2} \nabla \times \mathbf{A} \right|_T \right\}^{1/2} |\mathbf{A}'|_{j,\mu} \end{aligned}$$

where we have regrouped the face contributions in the last step and used that $a_F \leq \zeta_2 h_T$, cf. (9). \square

Using Lemma 3 we can finally prove discrete coercivity:

Lemma 4 (Discrete Coercivity). *The bilinear form a_h^{SWIP} is coercive: For all $\eta > C_{tr}^2 \zeta_2$ there exists a constant $C_{stab} > 0$ such that for all $h \in \mathcal{H}$*

$$a_h^{SWIP}(\mathbf{A}_h, \mathbf{A}_h) \geq C_{stab} \|\mathbf{A}_h\|_{SWIP}^2 \quad \forall \mathbf{A}_h \in V_h$$

with $C_{stab} = \min\left(\frac{\eta - C_{tr}^2 \zeta_2}{1 + \eta}, 1\right)$. The constant C_{tr} stems from the discrete trace inequality (5) and is independent of $h, \mu, \varepsilon, \zeta_2$.

Proof.

$$a_h^{\text{SWIP}}(\mathbf{A}_h, \mathbf{A}_h) = \left\| \mu^{-1/2} \nabla \times \mathbf{A}_h \right\|_{L^2(\Omega)^3}^2 - 2 \sum_{F \in \mathcal{F}_h} \int_F \{ \mu^{-1} \nabla \times \mathbf{A}_h \}_\omega \cdot [\mathbf{A}_h]_T + \sum_{F \in \mathcal{F}_h} \frac{\eta \gamma_{\mu, F}}{a_F} \| [\mathbf{A}_h]_T \|_{L^2(F)^3}^2 + \left\| \varepsilon^{1/2} \mathbf{A}_h \right\|_{L^2(\Omega)^3}^2$$

Now let us give a bound on the second term on the right-hand side using Lemma 3:

$$\begin{aligned} \left| \sum_{F \in \mathcal{F}_h} \int_F \{ \mu^{-1} \nabla \times \mathbf{A}_h \}_\omega \cdot [\mathbf{A}_h]_T \right| &\leq \zeta_2^{1/2} \left\{ \sum_{T \in \mathcal{T}_h} h_T \left\| \mu^{-1/2} \nabla \times \mathbf{A}_h \right\|_{L^2(\partial T)^3}^2 \right\}^{1/2} |\mathbf{A}_h|_{j, \mu} \\ &\leq C_{\text{tr}} \zeta_2^{1/2} \left\| \mu^{-1/2} \nabla \times \mathbf{A}_h \right\|_{L^2(\Omega)^3} |\mathbf{A}_h|_{j, \mu} \end{aligned}$$

where we have used the discrete trace inequality (5) componentwise in the last step. Hence,

$$\begin{aligned} a_h^{\text{SWIP}}(\mathbf{A}_h, \mathbf{A}_h) &\geq \left\| \varepsilon^{1/2} \mathbf{A}_h \right\|_{L^2(\Omega)^3}^2 \\ &\quad + \underbrace{\left\| \mu^{-1/2} \nabla \times \mathbf{A}_h \right\|_{L^2(\Omega)^3}^2}_{:=x^2} - 2 \underbrace{C_{\text{tr}} \zeta_2^{1/2}}_{:=\beta} \left\| \mu^{-1/2} \nabla \times \mathbf{A}_h \right\|_{L^2(\Omega)^3} |\mathbf{A}_h|_{j, \mu} + \eta \underbrace{|\mathbf{A}_h|_{j, \mu}^2}_{:=y^2}. \end{aligned}$$

Now use the inequality $x^2 - 2\beta xy + \eta y^2 \geq \frac{\eta - \beta^2}{1 + \eta} (x^2 + y^2)$ which holds for arbitrary β, η, x, y (it follows from $(\beta x - \eta y)^2 + (x - \beta y)^2 \geq 0$):

$$\begin{aligned} a_h^{\text{SWIP}}(\mathbf{A}_h, \mathbf{A}_h) &\geq \frac{\eta - C_{\text{tr}}^2 \zeta_2}{1 + \eta} \left(\left\| \mu^{-1/2} \nabla \times \mathbf{A}_h \right\|_{L^2(\Omega)^3}^2 + |\mathbf{A}_h|_{j, \mu}^2 \right) + \varepsilon \left\| \mathbf{A}_h \right\|_{L^2(\Omega)^3}^2, \\ &\geq C_{\text{stab}} \left(\left\| \mu^{-1/2} \nabla \times \mathbf{A}_h \right\|_{L^2(\Omega)^3}^2 + |\mathbf{A}_h|_{j, \mu}^2 + \varepsilon \left\| \mathbf{A}_h \right\|_{L^2(\Omega)^3}^2 \right) \end{aligned}$$

Finally, we note that $C_{\text{stab}} > 0$ if $\eta > C_{\text{tr}}^2 \zeta_2$ which completes the proof. \square

Lemma 5 (Boundedness). *There exists a constant $C_{\text{bnd}} > 0$ independent of h, μ , and ε such that for all $\mathbf{A} \in V_h^*$, all $\mathbf{A}'_h \in V_h$, all $h \in \mathcal{H}$:*

$$a_h^{\text{SWIP}}(\mathbf{A}, \mathbf{A}'_h) \leq C_{\text{bnd}} \|\mathbf{A}\|_{\text{SWIP},*} \|\mathbf{A}'_h\|_{\text{SWIP}}$$

Proof.

$$\begin{aligned} a_h^{\text{SWIP}}(\mathbf{A}, \mathbf{A}'_h) &= \int_{\Omega} (\mu^{-1} \nabla \times \mathbf{A}) \cdot (\nabla \times \mathbf{A}'_h) - \sum_{F \in \mathcal{F}_h} \int_F \{ \mu^{-1} \nabla \times \mathbf{A} \}_\omega \cdot [\mathbf{A}'_h]_T \\ &\quad - \sum_{F \in \mathcal{F}_h} \int_F \{ \mu^{-1} \nabla \times \mathbf{A}'_h \}_\omega \cdot [\mathbf{A}]_T + \sum_{F \in \mathcal{F}_h} \frac{\eta \gamma_{\mu, F}}{a_F} \int_F [\mathbf{A}]_T \cdot [\mathbf{A}'_h]_T + \varepsilon \int_{\Omega} \mathbf{A} \cdot \mathbf{A}'_h \\ &= T_1 + T_2 + T_3 + T_4 + T_5 \end{aligned}$$

Now let us bound these five terms individually:

$$\begin{aligned}
|T_1| &\stackrel{\text{(CS)}}{\leq} \left\| \mu^{-1/2} \nabla \times \mathbf{A} \right\|_{L^2(\Omega)^3} \left\| \mu^{-1/2} \nabla \times \mathbf{A}'_h \right\|_{L^2(\Omega)^3} \leq \|\mathbf{A}\|_{\text{SWIP}} \|\mathbf{A}'_h\|_{\text{SWIP}} \\
|T_2| &\stackrel{\text{Lemma 3}}{\leq} \zeta_2^{1/2} \|\mathbf{A}\|_{\text{SWIP},*} \|\mathbf{A}'_h\|_{\text{SWIP}} \\
|T_3| &\stackrel{\text{Lemma 3}}{\leq} \zeta_2^{1/2} \left\{ \sum_{T \in \mathcal{T}_h} h_T \left\| \mu^{-1/2} \nabla \times \mathbf{A}'_h \right\|_T \right\|_{L^2(\partial T)^3}^2 \right\}^{1/2} \|\mathbf{A}\|_{\text{SWIP}} \\
&\stackrel{\text{Lemma 1}}{\leq} C_{\text{tr}} \zeta_2^{1/2} \left\| \mu^{-1/2} \nabla \times \mathbf{A}'_h \right\|_{L^2(\Omega)^3} \|\mathbf{A}\|_{\text{SWIP}} \leq C_{\text{tr}} \zeta_2^{1/2} \|\mathbf{A}'_h\|_{\text{SWIP}} \|\mathbf{A}\|_{\text{SWIP}} \\
|T_4| &\stackrel{\text{(CS)}}{\leq} \eta |\mathbf{A}|_{j,\mu} |\mathbf{A}'_h|_{j,\mu} \leq \eta \|\mathbf{A}\|_{\text{SWIP}} \|\mathbf{A}'_h\|_{\text{SWIP}} \\
|T_5| &\stackrel{\text{(CS)}}{\leq} \left\| \varepsilon^{1/2} \mathbf{A} \right\|_{\text{SWIP}} \left\| \varepsilon^{1/2} \mathbf{A}'_h \right\|_{\text{SWIP}} \leq \|\mathbf{A}\|_{\text{SWIP}} \|\mathbf{A}'_h\|_{\text{SWIP}}
\end{aligned}$$

□

We can now finally combine the previous results into the following theorem:

Theorem 1 (Error Estimate). *Let $\mathbf{A} \in V^*$ be a solution of the strong formulation (1-2) (in the sense of distributions) and let $\mathbf{A}_h \in V_h \subseteq \mathbb{P}^k(\mathcal{T}_h)^3$ solve the variational formulation (6). Then there exist constants $C > 0$, $C_\eta > 0$, both independent of h , μ , ε such that for $\eta > C_\eta$,*

$$\|\mathbf{A} - \mathbf{A}_h\|_{\text{SWIP}} < C \inf_{\mathbf{v}_h \in V_h} \|\mathbf{A} - \mathbf{v}_h\|_{\text{SWIP},*}, \quad (16)$$

and the discrete problem (6) is well-posed. The constant C_η depends on ζ_2 and C depends on η , ζ_2 .

This theorem tells us that the total error is bounded by the best approximation error (w.r.t. suitable norms). Note that we didn't make any assumption on how the submeshes $\mathcal{T}_{h,1}$ and $\mathcal{T}_{h,2}$ meet at Γ . In order to get rates of convergence we will have to make additional assumptions about the approximation space V_h and the exact solution \mathbf{A} . This will be the topic of Section 4.

Proof (of Thm. 1). In this proof C denotes any arbitrary, positive constant, independent of h , μ that may have a different value every time it is used. Let us now pick an arbitrary $\mathbf{v}_h \in V_h$. Then, by the triangle inequality,

$$\|\mathbf{A} - \mathbf{A}_h\|_{\text{SWIP}} \leq \|\mathbf{A} - \mathbf{v}_h\|_{\text{SWIP}} + \|\mathbf{v}_h - \mathbf{A}_h\|_{\text{SWIP}} \quad (17)$$

This is almost the statement of Thm. 1. It remains to bound $\|\mathbf{A}_h - \mathbf{v}_h\|_{\text{SWIP}}$:

$$\begin{aligned}
\|\mathbf{A}_h - \mathbf{v}_h\|_{\text{SWIP}} &\stackrel{\text{Lemma 4}}{\leq} \frac{a_h^{\text{SWIP}}(\mathbf{A}_h - \mathbf{v}_h, \mathbf{A}_h - \mathbf{v}_h)}{\|\mathbf{A}_h - \mathbf{v}_h\|_{\text{SWIP}}} \\
&\leq C \sup_{\mathbf{w}_h \in V_h \setminus 0} \frac{a_h^{\text{SWIP}}(\mathbf{A}_h - \mathbf{v}_h, \mathbf{w}_h)}{\|\mathbf{w}_h\|_{\text{SWIP}}} \\
&\stackrel{\text{Lemma 2}}{=} C \sup_{\mathbf{w}_h \in V_h \setminus 0} \frac{a_h^{\text{SWIP}}(\mathbf{A} - \mathbf{v}_h, \mathbf{w}_h)}{\|\mathbf{w}_h\|_{\text{SWIP}}} \\
&\stackrel{\text{Lemma 5}}{\leq} C \sup_{\mathbf{w}_h \in V_h \setminus 0} \frac{\|\mathbf{A} - \mathbf{v}_h\|_{\text{SWIP},*} \|\mathbf{w}_h\|_{\text{SWIP}}}{\|\mathbf{w}_h\|_{\text{SWIP}}} \\
&\leq C \|\mathbf{A} - \mathbf{v}_h\|_{\text{SWIP},*}
\end{aligned}$$

Inserting this bound into (17) (which holds for arbitrary v_h) yields the assertion. Note that the bilinear form a_h^{SWIP} is coercive (Lemma 4) and bounded (finite dimensions). Thus, using Lax-Milgram it follows that the discrete problem is well posed. \square

Remark 1. Observe that for $\varepsilon \rightarrow 0$ the variational formulation (6) becomes ill-posed. To see this we observe that the $\|\cdot\|_{\text{SWIP}}$ norm “becomes” a semi-norm as $\varepsilon \rightarrow 0$. In order to study the behavior as $\varepsilon \rightarrow 0$ it is thus desirable to state the discrete coercivity (Lemma 4) w.r.t. a norm that does not depend on ε : We use that $\|\mathbf{A}_h\|_{\text{SWIP}}^2 \geq \varepsilon \|\mathbf{A}_h\|_{L^2}^2$ and thus Lemma (4) can be rewritten as

$$a_h^{\text{SWIP}}(\mathbf{A}_h, \mathbf{A}_h) \geq \varepsilon C_{\text{stab}} \|\mathbf{A}_h\|_{L^2}^2. \quad (18)$$

We see now clearly that the coercivity constant depends linearly on ε , i.e. the discrete problem becomes ill-posed as $\varepsilon \rightarrow 0$.

4 Rate of convergence for Edge Functions

In the following we will bound the best approximation error appearing in Theorem 1 for edge functions of the first kind. For this we assume, in addition to (9), that a_F is uniformly bounded from below in the sense that there exists a constant ζ_1 such that for all $h \in \mathcal{H}$, all $T \in \mathcal{T}_h$ and all $F \in \mathcal{F}_T$ we have

$$a_F \geq \zeta_1 h_T. \quad (19)$$

For the remainder of this section, let us choose $V_h = R^k(\Omega_1) \oplus R^k(\Omega_2) \subset \mathbb{P}^k(\mathcal{T}_h)^3$ where R^k is the space of k -th order edge functions ($k=1$ are the lowest order, $H(\mathbf{curl})$ conforming Whitney elements, cf. [9, Eq. (5.32)]). Because the sub-meshes $\mathcal{T}_{h,1}$, $\mathcal{T}_{h,2}$ are conforming, the spaces $R^k(\Omega_1)$, $R^k(\Omega_2)$ are $\mathbf{H}(\mathbf{curl})$ conforming. We can thus use the standard projection operator \mathbf{r}_h as it is defined in [9, Sect. 5.5] for edge functions on Ω_1 , Ω_2 to compose our global projection operator $\pi_h : V^* \rightarrow V_h$:

$$\mathbf{A} \mapsto \left(\mathbf{r}_h(\mathbf{A}|_{\Omega_1}), \mathbf{r}_h(\mathbf{A}|_{\Omega_2}) \right).$$

The following theorem then gives an upper bound for the best approximation error of Thm. 1:

Theorem 2. *Assume the exact solution of (1-2) is such that $\mathbf{A} \in \mathbf{H}(\mathbf{curl}, \Omega) \cap H^s(\Omega_1 \oplus \Omega_2)^3$, $\nabla \times \mathbf{A} \in H^s(\Omega_1 \oplus \Omega_2)$ with integer $1 \leq s \leq k$. Then*

$$\|\mathbf{A} - \pi_h \mathbf{A}\|_{\text{SWIP},*} < Ch^{s-1} \sum_{i=1}^2 \left(\|\mathbf{A}\|_{H^s(\Omega_i)} + \|\nabla \times \mathbf{A}\|_{H^s(\Omega_i)} \right)$$

where C depends on μ , ζ_1 but not on h . Moreover, C is independent of ε if $\varepsilon \leq \|\mu^{-1}\|_{L^\infty} = \mu_{\min}^{-1}$.

Remark 2. By combining Theorem 2 with Theorem 1 we see that for a sufficiently smooth exact solution \mathbf{A} , the total error $\|\mathbf{A} - \mathbf{A}_h\|_{\text{SWIP}} = O(h^{k-1})$ if k -th order edge functions are used. Theoretically it is possible that there exists another projector $\tilde{\pi}_h$ which would give a better rate of convergence, but numerical experiments show that Theorem 2 is sharp for $k = 1$ [1].

In order to prove the above theorem we will make use of two Lemmas to bound the face contributions:

Lemma 6. *Let $\mathcal{T}_{\mathcal{H}}$ be a sequence of shape regular, conforming, simplicial meshes. Suppose $\mathbf{u} \in H^s(T)^3$ and $\nabla \times \mathbf{u} \in H^s(T)^3$ for some integer $1 \leq s \leq k$. Then $\forall T \in \mathcal{T}_h \in \mathcal{T}_{\mathcal{H}}$*

$$\|\mathbf{u} - \mathbf{r}_h \mathbf{u}\|_{L^2(\partial T)^3} \leq Ch_T^{s-1/2} \left(\|\mathbf{u}\|_{H^s(T)^3} + \|\nabla \times \mathbf{u}\|_{H^s(T)^3} \right)$$

where C is independent of h_T , T .

For the proof of Lemma 6 we refer the reader to [9, Lemma 5.52] (which is proven element-wise).

Lemma 7. *Let $\mathcal{T}_{\mathcal{H}}$ be a sequence of shape regular, conforming, simplicial meshes. Assume $\mathbf{u} \in H^s(T)^3$ for some integer $1 \leq s \leq k$ and \mathbf{u} transforms such that it preserves the divergence, i.e. if $F : \hat{T} \rightarrow T$, $\hat{\mathbf{u}} \mapsto \mathbf{u}$ is an arbitrary mapping then \mathbf{u} transforms as*

$$\mathbf{u} \circ F = \frac{1}{|\det(dF)|} dF \hat{\mathbf{u}}. \quad (20)$$

Then the following estimate holds:

$$\|\mathbf{u} - \mathbf{w}_T \mathbf{u}\|_{L^2(\partial T)^3} \leq Ch_T^{s-1/2} \|\mathbf{u}\|_{H^s(T)^3} \quad \forall T \in \mathcal{T}_h \in \mathcal{T}_{\mathcal{H}}$$

where $\mathbf{w}_T : H^1(T)^3 \rightarrow D_k$ is the standard interpolation operator for k -th order Thomas-Raviart elements D_k [9, Sect. 5.4]. The constant C does not depend on h_T , T .

Proof. In order to simplify notation we will assume in this proof that $C > 0$ is an arbitrary constant independent of h , T that may have a different value every time it is used. We note that since $\mathbf{u} \in H^s(T)^3$, $\mathbf{w}_T \mathbf{u}$ is well defined by [9, Lemma 5.15]. Now split the integral over ∂T into its facet contributions:

$$\|\mathbf{u} - \mathbf{w}_T \mathbf{u}\|_{L^2(\partial T)^3}^2 = \sum_{F_T \in \mathbb{F}_T} \int_{F_T} |\mathbf{u} - \mathbf{w}_T \mathbf{u}|^2$$

Since our mesh contains only tetrahedrons we can find for every $F_T \in \mathbb{F}_T$ a linear transformation $\Phi_{T,F_T} : \hat{T} \rightarrow T$ which maps the reference element \hat{T} onto the actual element T such that the pre-image \hat{F}_T of facet F_T lies in the $x-y$ plane of \hat{T} :

$$\Phi_{T,F_T} : \hat{\mathbf{x}} \mapsto \mathbf{B}_{T,F_T} \hat{\mathbf{x}} + \mathbf{b}_{T,F_T}$$

where $\mathbf{B}_{T,F_T} \in \mathbb{R}^{3 \times 3}$. Now using the usual change of variables together with (20) we obtain

$$\begin{aligned} \int_{F_T} |\mathbf{u} - \mathbf{w}_T \mathbf{u}|^2 &= \int_{\hat{F}_T} |\det(\mathbf{B}_{T,F_T})^{-1} \mathbf{B}_{T,F_T} (\hat{\mathbf{u}} - \widehat{\mathbf{w}_T \mathbf{u}})|^2 |(\mathbf{B}_{T,F_T})_{:,1} \times (\mathbf{B}_{T,F_T})_{:,2}| \\ &= \frac{\text{area}(F_T)}{\text{area}(\hat{F}_T) |\det(\mathbf{B}_{T,F_T})|^2} \int_{\hat{F}_T} |\mathbf{B}_{T,F_T} (\hat{\mathbf{u}} - \mathbf{w}_{\hat{T}} \hat{\mathbf{u}})|^2 \\ &\leq Ch_T^2 |\det(\mathbf{B}_{T,F_T})|^{-2} \|\mathbf{B}_{T,F_T}\|^2 \|\hat{\mathbf{u}} - \mathbf{w}_{\hat{T}} \hat{\mathbf{u}}\|_{L^2(\hat{F}_T)}^2 \end{aligned} \quad (21)$$

Where $(\mathbf{B}_{T,F_T})_{:,i}$ denotes the i -th column of \mathbf{B}_{T,F_T} and we have used that $\widehat{\mathbf{w}_T \mathbf{u}} = \mathbf{w}_{\hat{T}} \hat{\mathbf{u}}$ [9, Lemma 5.22]. Now notice that $\hat{\mathbf{u}} - \mathbf{w}_{\hat{T}} \hat{\mathbf{u}} \in H^s(\hat{T})^3$ and thus we can use the trace inequality [9, Thm. 3.9]:

$$\|\hat{\mathbf{u}} - \mathbf{w}_{\hat{T}} \hat{\mathbf{u}}\|_{L^2(\hat{F}_T)}^3 \leq \|\hat{\mathbf{u}} - \mathbf{w}_{\hat{T}} \hat{\mathbf{u}}\|_{L^2(\partial \hat{T})}^3 \leq C \|\hat{\mathbf{u}} - \mathbf{w}_{\hat{T}} \hat{\mathbf{u}}\|_{H^1(\hat{T})}^3$$

Next notice that $\forall \phi \in \mathbb{P}^{k-1}(\hat{T})^3 \subset D_k(\hat{T})$, $\phi = \mathbf{w}_{\hat{T}} \phi$ by the definition of $\mathbf{w}_{\hat{T}}$. Therefore,

$$\|\hat{\mathbf{u}} - \mathbf{w}_{\hat{T}} \hat{\mathbf{u}}\|_{H^1(\hat{T})}^3 \leq \|(\mathbf{I} - \mathbf{w}_{\hat{T}})(\hat{\mathbf{u}} + \phi)\|_{H^1(\hat{T})}^3 \leq C \|\hat{\mathbf{u}} + \phi\|_{H^1(\hat{T})}^3$$

where we have used that $\mathbf{w}_{\hat{T}} : H^1(\hat{T})^3 \rightarrow D_k$ is a bounded operator, i.e. $\|\mathbf{w}_{\hat{T}} \hat{\mathbf{u}}\|_{H^1(\hat{T})}^3 \leq C \|\hat{\mathbf{u}}\|$ (cf. Proof of [9, Thm. 5.25]). Since ϕ is arbitrary we can use the Deny-Lions theorem [9, Thm. 5.5]

$$\|\hat{\mathbf{u}} - \mathbf{w}_{\hat{T}} \hat{\mathbf{u}}\|_{L^2(\hat{T})}^3 \leq C \inf_{\phi \in \mathbb{P}^{k-1}(\hat{T})^3} \|\hat{\mathbf{u}} + \phi\|_{H^1(\hat{T})}^3 \leq C |\hat{\mathbf{u}}|_{H^s(\hat{K})}^3 \quad (22)$$

Finally we have to map $|\hat{\mathbf{u}}|_{H^s(\hat{T})}^3$ back to the actual element T . For this observe that using (20),

$$\frac{\partial^\alpha}{\partial \hat{\mathbf{x}}^\alpha} \hat{\mathbf{u}} = \det(\mathbf{B}_{T,F_T}) \mathbf{B}_{T,F_T}^{-1} \frac{\partial^\alpha}{\partial \mathbf{x}^\alpha} (\mathbf{u} \circ \Phi_T)$$

where $\|\alpha\|_{\ell^1} = s$ is a multi-index. Therefore,

$$\begin{aligned} |\hat{\mathbf{u}}|_{H^s(\hat{T})}^2 &= \sum_{|\alpha|_{\ell^1} = s} \int_{\hat{T}} \left| \frac{\partial^\alpha \hat{\mathbf{u}}}{\partial \hat{\mathbf{x}}^\alpha} \right|^2 \leq |\det(\mathbf{B}_{T,F_T})|^2 \left\| \mathbf{B}_{T,F_T}^{-1} \right\|^2 \int_{\hat{T}} \left| \frac{\partial^\alpha (\mathbf{u} \circ \Phi_T)}{\partial \hat{\mathbf{x}}^\alpha} \right|^2 \\ &\leq C |\det(\mathbf{B}_{T,F_T})|^2 \left\| \mathbf{B}_{T,F_T}^{-1} \right\|^2 \|\mathbf{B}_{T,F_T}\|^{2s} |\det(\mathbf{B}_{T,F_T})|^{-1} |\mathbf{u}|_{H^s(T)}^2 \end{aligned} \quad (23)$$

where we have used [9, Lemma 5.9] in the last step. Now combining (21-23) gives:

$$\begin{aligned} \|\mathbf{u} - \mathbf{w}_T \mathbf{u}\|_{L^2(F_T)}^2 &\leq C |\det(\mathbf{B}_{T,F_T})|^{-1} h_T^2 \|\mathbf{B}_{T,F_T}\|^2 \left\| \mathbf{B}_{T,F_T}^{-1} \right\|^2 \|\mathbf{B}_{T,F_T}\|^{2s} |\mathbf{u}|_{H^s(T)}^2 \\ &\leq Ch_T^{2s-1} |\mathbf{u}|_{H^s(T)}^2 \end{aligned}$$

where we have used [9, Lemma 5.10] together with the fact that the mesh sequence is shape regular. Now summing over all facets $F_T \in \mathbb{F}_T$ yields the assertion. \square

Using these Lemmas we can finally give a bound for $\|\mathbf{A} - \pi_h \mathbf{A}\|_{\text{SWIP},*}$:

Proof (of Theorem 2). In order to simplify notation, C denotes in this proof an arbitrary, positive constant that is independent of h . Note that the interpolation operator $\mathbf{r}_h \left(\mathbf{A}|_{\Omega_1} \right)$ is well defined for $s \geq 1$ by the Sobolev Embedding Theorem and [9, Lemma 5.38]. Because the sub-meshes of Ω_1, Ω_2 are conforming, $\mathbf{r}_h \left(\mathbf{A}|_{\Omega_1} \right)$ is tangentially continuous across all inner, conforming faces. The same holds for Ω_2 and because $\mathbf{A} \in \mathbf{H}(\text{curl}; \Omega)$ the exact solution is also tangentially continuous across all inner faces. Therefore only jump terms across the faces $F \in \mathcal{F}_h^{\Gamma, b} := \mathcal{F}_h^b \cup \{F \in \mathcal{F}_h^i \mid F \subset \overline{\Omega_1} \cap \overline{\Omega_2}\}$ remain in the definition of the jump semi-norm $|\cdot|_{j, \mu}$. I.e. we have to bound

$$\begin{aligned} \|\mathbf{A} - \pi_h \mathbf{A}\|_{\text{SWIP},*}^2 &= \underbrace{\|\mu^{-1/2} \nabla \times (\mathbf{A} - \pi_h \mathbf{A})\|_{L^2(\Omega)^3}^2}_{:=T_1} + \underbrace{\|\varepsilon^{1/2} (\mathbf{A} - \pi_h \mathbf{A})\|_{L^2(\Omega)^3}^2}_{:=T_2} \\ &+ \underbrace{\sum_{F \in \mathcal{F}_h^{b, \Gamma}} \frac{\gamma_{\mu, F}}{a_F} \|\mathbf{A} - \pi_h \mathbf{A}\|_T^2}_{:=T_3} + \underbrace{\sum_{T \in \mathcal{T}_h} h_T \|\mu^{-1/2} \nabla \times (\mathbf{A} - \pi_h \mathbf{A})\|_{L^2(\partial T)^3}^2}_{:=T_4}, \end{aligned} \quad (24)$$

Since μ is piecewise constant on each $\Omega_{i, \mu} \in P_\Omega$ there are constants μ_{\min}, μ_{\max} such that $0 < \mu_{\min} < \mu < \mu_{\max}$. T_1 and T_2 are easily bounded using [9, Thm. 5.41]:

$$\begin{aligned} T_1 + T_2 &\leq \max(\mu_{\min}^{-1}, \varepsilon) \left[\|\nabla \times (\mathbf{A} - \pi_h \mathbf{A})\|_{L^2(\Omega)}^2 + \|(\mathbf{A} - \pi_h \mathbf{A})\|_{L^2(\Omega)}^2 \right] \\ &\leq Ch^{2s} \left[\sum_{i=1}^2 \left(\|\mathbf{A}\|_{H^s(\Omega_i)^3} + \|\nabla \times \mathbf{A}\|_{H^s(\Omega_i)^3} \right) \right]^2 \end{aligned}$$

The term T_3 is bounded using Lemma 6:

$$\begin{aligned} T_3 &\leq 2\mu_{\min}^{-1} \sum_{F \in \mathcal{F}_h^{b, \Gamma}} a_F^{-1} \left(\|\mathbf{A} - \pi_h \mathbf{A}\|_{T_1}^2 + \|\mathbf{A} - \pi_h \mathbf{A}\|_{T_2}^2 \right) \\ &\leq C \sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T \cap \mathcal{F}_h^{b, \Gamma}} h_T^{-1} \|\mathbf{A} - \pi_h \mathbf{A}\|_T^2 \\ &\leq C \sum_{T \in \mathcal{T}_h} \sum_{F \in \mathcal{F}_T \cap \mathcal{F}_h^{b, \Gamma}} h_T^{2s-2} \left(\|\mathbf{A}\|_{H^s(T)^3}^2 + \|\nabla \times \mathbf{A}\|_{H^s(T)^3}^2 \right) \\ &\leq Ch^{2s-2} \sum_{i=1}^2 \left(\|\mathbf{A}\|_{H^s(\Omega_i)^3}^2 + \|\nabla \times \mathbf{A}\|_{H^s(\Omega_i)^3}^2 \right) \end{aligned}$$

where we have used that $a_F \geq \zeta_1 h_T$.

In order to bound the term T_4 we first note that the global Thomas-Raviart interpolation operator $\mathbf{w}_h \left(\nabla \times \mathbf{A}|_{\Omega_i} \right)_{i=1,2}$ is well defined by [9, Lemma 5.15]. Thus, $\nabla \times \left[\mathbf{r}_h \left(\mathbf{A}|_{\Omega_i} \right) \right] =$

$\mathbf{w}_h \left(\nabla \times \mathbf{A}|_{\Omega_i} \right)$ by [9, Lemma 5.40] and we can bound T_4 as follows:

$$\begin{aligned} T_4 &\leq \mu_{\min}^{-1} \sum_{T \in \mathcal{T}_h} h_T \|\nabla \times \mathbf{A} - \mathbf{w}_h(\nabla \times \mathbf{A})\|_{L^2(\partial T)}^2 \\ &\leq C \sum_{T \in \mathcal{T}_h} h_T^{2s} \|\nabla \times \mathbf{A}\|_{H^s(T)}^3 \leq Ch^{2s} \sum_{i=1}^2 \|\nabla \times \mathbf{A}\|_{H^s(\Omega_i)}^3 \end{aligned}$$

where we have used Lemma 7 and the fact that $h_T \leq h$. \square

Remark 3. From the proof of Theorem 2 it is clear that for h sufficiently small the term T_3 dominates the other three terms and is thus responsible for the loss of one order of convergence as pointed out in Remark 2. Interestingly T_3 sums the jump terms only over the faces $\mathcal{F}_h^{b,\Gamma}$. This suggests that it suffices to use $(k+1)$ -th order edge functions in elements adjacent to $\mathcal{F}_h^{b,\Gamma}$ and k -th order edge functions everywhere else to achieve $O(h^k)$ order convergence. This can be implemented easily by using a hierarchical basis for the edge functions [10].

5 The local length scale a_F

So far we have assumed that the local length scale a_F fulfills (9), (19) in order to derive an a-priori error estimator, i.e. $0 < \varsigma_1 h_T \leq a_F \leq \varsigma_2 h_T$. We will now study the following, three concrete choices for a_F :

- $a_F^{(1)} := \frac{1}{2}(h_{T_1} + h_{T_2})$ if $F \in \mathcal{F}_h^i$ and $a_F^{(1)} = h_T$ for $F \in \mathcal{F}_h^b$, see [3][Remark 4.6]
- $a_F^{(2)} := \min(h_{T_1}, h_{T_2})$ if $F \in \mathcal{F}_h^i$ and $a_F^{(2)} = h_T$ for $F \in \mathcal{F}_h^b$, see [8]
- $a_F^{(3)} := h_F$ if $F \in \mathcal{F}_h$ see [2, 3]

where h_{T_1}, h_{T_2} are the diameters of the adjacent elements of face F and h_F is the diameter of face F . It turns out that for each choice of a_F we have to make additional assumptions on the mesh such that a_F fulfills (9), (19). So once we have chosen a concrete a_F we can think of ς_1, ς_2 as mesh dependent parameters. The important point is that the constants C in Theorems 1 and 2 depend on the constants $\sigma_{\max}, \varsigma_1, \varsigma_2$ but they do not depend in any other way on the shape of the underlying meshes. Hence, if we can assure that $\sigma_{\max}, \varsigma_1, \varsigma_2$ are independent of the way that $\mathcal{T}_{\mathcal{H},1}, \mathcal{T}_{\mathcal{H},2}$ intersect at the sliding interface Γ , we can be sure that there is an upper bound on the total error $\|\mathbf{A} - \mathbf{A}_h\|$ that is independent of the relative position of $\mathcal{T}_{\mathcal{H},1}$ to $\mathcal{T}_{\mathcal{H},2}$ and that tends to 0 as $h \rightarrow 0$.

Let us now discuss the precise conditions on the mesh for each choice of a_F ; For $a_F^{(1)}, a_F^{(2)}$ we require $\mathcal{T}_{\mathcal{H}}$ to be *quasi-uniform at the sliding interface*:

Definition 1. A mesh-sequence $\mathcal{T}_{\mathcal{H}}$ is said to be *quasi-uniform at Γ* if it is shape regular (3) and if there exists a constant $\sigma_1 > 0$ such that for all $h \in \mathcal{H}$, all $T \in \mathcal{T}_h^\Gamma := \{T \in \mathcal{T}_h \mid \partial T \cap \Gamma \neq \emptyset\}$:

$$h_T \geq \sigma_1 \max_{\tilde{T} \in \mathcal{T}_h^\Gamma} h_{\tilde{T}}. \quad (25)$$

Lemma 8. $a_F^{(1)}, a_F^{(2)}$ fulfill conditions (9), (19) if the mesh is quasi-uniform at Γ . Moreover the constants $\sigma_{\max}, \zeta_1, \zeta_2$ are independent of the way $\mathcal{T}_{\mathcal{H},1}, \mathcal{T}_{\mathcal{H},2}$ intersect at Γ .

Proof. $a_F^{(1)} \geq \frac{1}{2}h_{T_i}$ follows immediately from the definition for $i = 1, 2$. For the other direction we use (25) and get $a_F^{(1)} \leq \frac{1}{2}(1 + \sigma_1^{-1})h_{T_i}$. Moreover, $\sigma_1 h_{T_i} \leq a_F^{(2)} \leq h_{T_i}$. \square

The Lemma above asserts that the choices $a_F^{(1)}, a_F^{(2)}$ lead to a method that converges independent of the way that the two mesh-sequences $\mathcal{T}_{\mathcal{H},1}, \mathcal{T}_{\mathcal{H},2}$ intersect at Γ . In particular the faces can be very tiny “slivers” (i.e. triangles with high aspect ratio). But note that the choice of a_F determines the required minimum value of the penalty parameter (see Lemma 4).

By substituting $a_F^{(3)}$ into (19) we see that we need an estimate of the form $h_F \geq \zeta_1 h_T$ in order for Thm. 2 to hold. However if two meshes are sliding against each other such an estimate is not feasible since h_F can become arbitrarily small in comparison to h_T . In other words, the constant ζ_1 depends on the way $\mathcal{T}_{\mathcal{H},1}$ intersects with $\mathcal{T}_{\mathcal{H},2}$. Nevertheless using $a_F^{(3)}$ in the variational formulation 6 seems to work in practice (see below).

We study the behavior of the SWIP formulation for the three different choices of the local length scale a_F ; As in [1] we consider a 3D sphere with radius 1 that is split into two half-spheres which are then meshed separately (Fig. 2). We impose the analytical solution $\mathbf{A} = (\sin y, \cos z, \sin x)$ and choose $\mathbf{j}^i, \mathbf{g}_D$ such that they fulfill (1-2).




Fig. 2 The meshes for the two half spheres. The upper hemisphere is turned against the lower hemisphere by $\theta = 2.86$ degrees to create a non-conforming mesh.

Figure 3 shows the $\mathbf{H}(\text{curl})$ error for different angles of rotation for all three choices of a_F and for different mesh-sizes h . We can see that although the error depends slightly on the angle of rotation, it converges to zero in all three formulations as h is decreasing (see also [1]). Moreover we see that the choices $a_F^{(1)}, a_F^{(2)}$ yield similar results which are slightly better than the choice $a_F^{(3)}$.




Fig. 3 The relative $\mathbf{H}(\mathbf{curl})$ error vs. the rotation angle for three different choices of a_F and 4 different mesh-sizes: $h = 0.638174, 0.482025, 0.359644, 0.261798$. $\varepsilon = 10^{-6}$, $\eta = 50$, $\mu \equiv 1$.

6 The regularization parameter ε

So far we have looked at the regularized system (1-2) and in [1] it was shown that the proposed method yields the expected rates of convergence for $\varepsilon > 0$. However, genuine magnetostatics amounts to choosing $\varepsilon = 0$. We will consider two approaches to solve the system (1-2) with $\varepsilon = 0$: On the one hand we will try to set $\varepsilon = 0$ directly and on the other hand we will study the effect of choosing ε small enough such that the error due to regularization is negligible.

6.1 The case $\varepsilon = 0$

We note that by setting $\varepsilon = 0$, the boundary value problem (1-2) ceases to have a unique solution. Indeed the continuous $\mathbf{curl} - \mathbf{curl}$ operator has an infinite-dimensional kernel and the non-zero eigenvalues are well separated from 0 [9][Corollary 4.8]. If one uses $\mathbf{H}(\mathbf{curl})$ conforming edge functions of the first kind on a conforming mesh it can be shown that the discrete $\mathbf{curl} - \mathbf{curl}$

operator has a (finite-dimensional) kernel and that the discrete eigenvalues are well separated from it [9][Discrete Friedrichs inequality, Lemma 7.20]. I.e. edge functions of the first kind yield a spectrally accurate discretization of the **curl** – **curl** operator. From a theoretical point of view it remains unclear whether this property carries over to the SWIP formulation (6), cf. [14].

Therefore the spectrum of the a_h^{SWIP} bilinear form is investigated in a numerical experiment. The setup is very similar to the one in the previous section: The domain Ω consists of two half-spheres which can be rotated against each other by an angle θ . However this time we only assemble the matrix of the a_h^{SWIP} bilinear form with $\varepsilon = 0$, $a_F = a_F^{(3)}$ ¹ and compute its eigenvalues using the `eig` routine of MATLAB R2013a.




Fig. 4 The smallest/largest non-zero eigenvalue is plotted against the meshwidth for 50 different angles of rotation (dashed lines). For comparison the smallest/largest non-zero eigenvalue of a $\mathbf{H}(\mathbf{curl})$ conforming discretization based on second order edge functions is plotted as well. The angles are $\theta = 0.01n$ rad, $n \in 0, \dots, 49$ and R^2 edge functions were used to discretize a_h^{SWIP} .

Figure 4 shows the smallest and largest non-zero eigenvalues of the SWIP formulation for different mesh-widths h and different angles θ (dashed, blue lines) (a eigenvalue has been classified

¹ The choices $a_F^{(1)}$ and $a_F^{(2)}$ yield qualitatively the same results. In particular the smallest non-zero eigenvalues also tend to 0 as $\varepsilon \rightarrow 0$, cf. Fig. 5




Fig. 5 Smallest/Largest non-zero eigenvalues vs. the rotation angle θ for $h = 0.359644$. The discretization is based on R^2 edge functions.

as non-zero if its absolute value is greater than 10^{-12}). For comparison we have also plotted the eigenvalues of a standard $\mathbf{H}(\mathbf{curl})$ conforming discretization using second order edge functions on the conforming grid with $\theta = 0$ (green lines).

We see that the bandwidth of the SWIP eigenvalues is comparable to the bandwidth of the $\mathbf{H}(\mathbf{curl})$ conforming discretization for many angles. But we also observe that for some angles the lower end of the spectrum tends to zero. In order to better understand this phenomena we plotted the smallest/largest non-zero eigenvalues of the SWIP discretization against θ for one mesh-size (Fig. 5). We now see that the lower end of the spectrum deteriorates as $\theta \rightarrow 0$. I.e. we can expect spectral pollution for very small angles. This agrees with the observations of [13].

The previous considerations indicate that the a_h^{SWIP} bilinear form is not suitable to solve the Maxwell Eigenvalue Problem. However in this work we are concerned with the $\mathbf{curl} - \mathbf{curl}$ source problem (1-2). Although the Galerkin matrix becomes singular for $\varepsilon = 0$ we can in principle still solve the linear system if it is consistent, i.e. if the right-hand side lies in the range of the Galerkin matrix. Then the solution \mathbf{A}_h is not unique anymore, but $\mathbf{curl} \mathbf{A}_h$ is.

We attempt to solve the linear system of equations using the conjugate gradient (CG) method. In [15] it is shown that the CG method converges for consistent, symmetric positive semi-definite

problems and that its rate of convergence is determined by the non-zero eigenvalues. In particular, the number of CG iterations is related to the generalized condition number $\kappa = \frac{\lambda_{\max}}{\lambda_{\min}}$ where λ_{\min} is the smallest, non-zero eigenvalue of the system matrix. If again we take a look at Fig. 5 it becomes clear that $\kappa \rightarrow \infty$ as $\theta \rightarrow 0$. I.e. the number of CG iterations should increase as $\theta \rightarrow 0$.

This has been confirmed in a numerical experiment: We take the example from Sec. 5 with the same analytical solution and chose the right-hand side $\mathbf{j}^i = \nabla \times (\nabla \times \mathbf{A})$ ($\varepsilon = 0$, $\mu \equiv 1$). Table 1 provides the number of CG iterations required to reach the prescribed tolerance 10^{-6} . We see that without a preconditioner the computational cost for the angle $\theta = 10^{-6}$ is almost 6 times larger than for $\theta = 10^{-1}$. For comparison we also list the number of iterations needed when the multi-level ILU decomposition ILUPACK is employed [5] [6]². In this case the number of iterations also increases but the factor 6 is reduced to ≈ 3.15 .

Table 1 Number of CG iterations for $\theta \rightarrow 0$, $h = 0.359$; the discretization is based on R^2 edge functions.

θ [rad]	No Preconditioner	ILUPACK
10^{-1}	1118	135
10^{-2}	3705	214
10^{-3}	3731	320
10^{-4}	6102	426

Remark 4. Although the right-hand side \mathbf{j}^i chosen in the numerical experiment above is clearly divergence free, there is no guarantee that its discrete counterpart ℓ_h is so too. We have investigated this by splitting the right-hand side vector \mathbf{b} of the linear system into a part that lies in the kernel of matrix, $\tilde{\mathbf{b}}$, and into its orthogonal complement, $\tilde{\mathbf{b}}^\perp$. It turned out that for all angles $\|\tilde{\mathbf{b}}^\perp\|_2 / \|\mathbf{b}\|_2 \approx 10^{-9}$, which seems to be sufficient for CG to converge.

6.2 The case $0 < \varepsilon \ll 1$

If we set ε to zero we have to make sure that the right-hand side vector lies in the range of the system matrix in order for CG to converge. However this is a non-trivial task because we don't know a-priori the kernel of the system matrix.

Note 1. For $\mathbf{H}(\mathbf{curl})$ conforming discretizations, which fulfill the discrete sequence property, the kernel of the system matrix is known. Unfortunately this doesn't carry over to the SWIP formulation (6).

Let us therefore study a different approach: We choose ε so small that the error due to regularization becomes negligible. To make this more explicit we bound the total error between the discrete, regularized solution \mathbf{A}_h^ε and the exact solution of (1-2) with $\varepsilon = 0$, \mathbf{A}^0 , by two contributions:

² The parameters for ILUPACK are: type sol = 0, partitioning=3, flags=-1,-1, inv. droptol=5, threshold ILU=0.1, condest=1e-2, residual tol. = 5e-6

$$\|\nabla \times (\mathbf{A}_h^\varepsilon - \mathbf{A}^0)\|_{L^2(\Omega)} \leq \|\nabla \times (\mathbf{A}_h^\varepsilon - \mathbf{A}^\varepsilon)\|_{L^2(\Omega)} + \|\nabla \times (\mathbf{A}^\varepsilon - \mathbf{A}^0)\|_{L^2(\Omega)},$$

herein \mathbf{A}^ε is the exact solution of the regularized system (1-2). Clearly the second component is independent of the discretization and thus h , but it depends on ε for a given problem. Moreover, the first term depends on h but is independent of ε because the constant C of Thms. 1 and 2 is independent of ε (for ε small enough).

It is thus desirable to choose ε small such that $\|\nabla \times (\mathbf{A}^\varepsilon - \mathbf{A}^0)\|_{L^2(\Omega)^3} \ll \|\nabla \times (\mathbf{A}_h^\varepsilon - \mathbf{A}^\varepsilon)\|_{L^2(\Omega)^3}$. However, as $\varepsilon \rightarrow 0$ the discrete problem becomes ill-posed, cf. Remark 3.1, Sec. 6.1.

We try to circumvent this problem by two approaches:

- For small problems we use the Sparse Cholesky Decomposition of PARDISO [12] (Intel MKL Version 11.2) and solve the linear system of equations directly.
- For problems whose Cholesky Decomposition does not fit into memory we use the Conjugate Gradient Method together with ILUPACK [5, 6] as a preconditioner (using the settings of Sect. 6.1).

Note 2. We are only interested in the curl of the solution, i.e. the magnetic field \mathbf{B} . If we were to look at \mathbf{A} instead of $\nabla \times \mathbf{A}$ then $\|\mathbf{A}_h^\varepsilon - \mathbf{A}^\varepsilon\|_{L^2(\Omega)^3}$ would *not* be independent of ε as can be seen from Thm. 1.

Setup We consider the same setup as in the previous section (cf. Fig. 2) with the same analytic solution, $\mathbf{A}^0 = (\sin y, \cos z, \sin x)$. \mathbf{j}^i is chosen such that \mathbf{A}^0 fulfills (1-2) with $\varepsilon = 0$ and $\mu \equiv 1$.

We solve the system of linear equations using PARDISO for different values of ε (as in the previous section we choose $a_F = a_F^{(3)}$). Figure 6 shows the total error $\|\nabla \times (\mathbf{A}_h^\varepsilon - \mathbf{A}^0)\|_{L^2(\Omega)}$ as a function of ε for various mesh-sizes. We see that for $\varepsilon < 10^{-3}$ the discretization error $\|\nabla \times (\mathbf{A}_h^\varepsilon - \mathbf{A}^\varepsilon)\|_{L^2(\Omega)}$ clearly dominates the regularization error $\|\nabla \times (\mathbf{A}^\varepsilon - \mathbf{A}^0)\|_{L^2(\Omega)}$ whereas for $\varepsilon > 10^{-3}$ the discretization error starts to dominate the regularization error. This is what we can expect from the previous discussion.

Note 3. The same results are obtained if CG together with ILUPACK is used. For brevity we omit these results here.

We would like to point out that by using the direct solver PARDISO we were able to solve the resulting system of linear equations for ε as small as 10^{-10} and that the time needed to solve the problem seems to be independent of ε (see Table 2). A similar result holds for preconditioned CG with ILUPACK preconditioner where the system is solvable for arbitrary small ε (cf. Sec. 6.1) and the solution time seems to be independent of ε for ε small enough.

We can thus choose ε (almost) arbitrarily small without affecting the discretization error $\|\nabla \times (\mathbf{A}_h^\varepsilon - \mathbf{A}^\varepsilon)\|_{L^2(\Omega)}$ and incurring rising cost for solving the resulting linear systems of equations. In other words, the regularization error can be made arbitrarily small such that it becomes negligible for concrete applications.




Fig. 6 Relative L^2 -error of curl vs. ϵ for multiple mesh-sizes h . The meshes have been rotated against each other by $\theta = 0.057$ degrees and second order edge functions ($k = 2$) were used for discretization.

Table 2 Relative runtimes for $\epsilon \rightarrow 0$, $h = 0.359$; the discretization is based on R^2 edge functions and $\theta = 10^{-4}$ rad. The runtimes have been normalized with the runtime for $\epsilon = 10^{-1}$.

ϵ	PARDISO ^a	ILUPACK ^b
10^{-1}	1	1
10^{-2}	1.01	1.41
10^{-3}	1.01	1.42
10^{-4}	1.02	1.43
10^{-5}	0.98	1.42

^a Time includes cholesky factorization and back-substitution

^b Time includes ILU factorization and CG iterations

7 Conclusion and Outlook

We have proved a-priori error estimators for the interior penalty formulation of the regularized curl-curl source problem (1-2); If the solution is approximated by k -th order edge functions we can expect at least convergence of order $O(h^{k-1})$ (provided the exact solution is sufficiently smooth). In particular, for $k = 1$ no convergence was observed in a numerical experiment [1], which implies that our result is sharp. The reason for this is that R^k does not span the full polynomial space \mathbb{P}^k .

The bounds require the mesh to be *quasi-uniform* at the sliding interface but do *not* make any assumptions on how the sub-meshes abut at the sliding interface nor does the error estimate depend on it. This is confirmed by the numerical experiments in [1] where it is shown that the approximation is stable independent of the way the sub-meshes intersect.

Moreover the role of the regularization parameter ε has been investigated; For practical purposes one can choose ε (almost) arbitrarily small and solve the discrete problem with a direct solver or by using the preconditioned conjugate gradient method. The error due to regularization is then dominated by the discretization error of the regularized problem and is thus negligible.

Outlook The proof of Thm. 2 suggest that it suffices to use 2nd order edge functions solely in elements adjacent to the non-conforming interface, respectively boundary faces, to achieve $O(h)$ convergence. This would reduce the required number of unknowns drastically and will be presented in a future work.

References

1. Casagrande, R., Winkelmann, C., Hiptmair, R., Ostrowski, J.: DG Treatment of Non-Conforming Interfaces in 3D Curl-Curl Problems. SAM Report 2014-32, <http://www.sam.math.ethz.ch>, Zürich, (2014),
2. Stenberg, R.: Mortaring by a method of JA Nitsche. Computational mechanics (1998)
3. Di Pietro, D.A., Ern, A.: Mathematical aspects of discontinuous Galerkin methods. Springer, Heidelberg, London, New York (2012)
4. Brenner, S.C., Scott, L.R.: The Mathematical Theory of Finite Element Methods. Springer, New York (2008)
5. Bollhöfer, M., Saad, Y.: Multilevel preconditioners constructed from inverse-based ILU's. SIAMSciComp, **27.5**, 1627-1650, (2006)
6. Aliaga, J.I., Bollhöfer, M., Martin, A., Quintana-Orti, E.: ILUPACK. In: Encyclopedia of Parallel Computing pp. 917-926, Springer, New York Dordrecht Heidelberg London (2012)
7. Casagrande, R.: Sliding Interfaces for Eddy Current Simulations. Master's thesis, <http://www.sam.math.ethz.ch>, ETH Zürich, Zürich (2013)
8. Houston, P., Perugia, I., Schötzau, D.: Mixed Discontinuous Galerkin Approximation of the Maxwell Operator: Non-Stabilized Formulation. Journal of Scientific Computing, **22.1-3**, 315-346, (2005)
9. Monk, P.: Finite element methods for Maxwell's equations. Oxford University Press, New York (2003)
10. Bergot, M., Duruflé, M.: High-order optimal edge elements for pyramids, prisms and hexahedra. Journal of Computational Physics, **232**, 189-213, (2013)
11. Adams, R.A., Fournier, J.F.: Sobolev Spaces. Academic Press, Oxford (2003)
12. Schenk, O., Gärtner, K.: Solving unsymmetric sparse systems of linear equations with PARDISO. J. of Future Generation Computer Systems, 20(3):475-487, (2004)
13. Buffa, A., Houston, P., Perugia, I.: Discontinuous Galerkin computation of the Maxwell eigenvalues on simplicial meshes. J. of Comp. and Applied Mathematics, **204**, 317-333, (2007)
14. Buffa, A., Perugia, I.: Discontinuous Galerkin Approximation of the Maxwell Eigenproblem. SIAM Journal on Numerical Analysis, **44.5**, 2198-2226, (2006)
15. Kaasschieter, E.: Preconditioned conjugate gradients for solving singular systems. Journal of Comp. and Applied Mathematics, **24**, 265-275, (1988)

Recent Research Reports

Nr.	Authors/Title
2014-30	M. Hansen A new embedding result for Kondratiev spaces and application to adaptive approximation of elliptic PDEs
2014-31	F. Mueller and Ch. Schwab Finite elements with mesh refinement for elastic wave propagation in polygons
2014-32	R. Casagrande and C. Winkelmann and R. Hiptmair and J. Ostrowski DG Treatment of Non-Conforming Interfaces in 3D Curl-Curl Problems
2014-33	U. Fjordholm and R. Kappeli and S. Mishra and E. Tadmor Construction of approximate entropy measure valued solutions for systems of conservation laws.
2014-34	S. Lanthaler and S. Mishra Computation of measure valued solutions for the incompressible Euler equations.
2014-35	P. Grohs and A. Obermeier Ridgelet Methods for Linear Transport Equations
2014-36	P. Chen and Ch. Schwab Sparse-Grid, Reduced-Basis Bayesian Inversion
2014-37	R. Kaeppli and S. Mishra Well-balanced schemes for gravitationally stratified media
2014-38	D. Schoetzau and Ch. Schwab Exponential Convergence for hp-Version and Spectral Finite Element Methods for Elliptic Problems in Polyhedra
2014-39	P. Grohs and M. Sprecher Total Variation Regularization by Iteratively Reweighted Least Squares on Hadamard Spaces and the Sphere